Fatou's web and non-escaping endpoints

Vasiliki Evdoridou
Dept. of Mathematics and Statistics
The Open University

Workshop on Ergodic Theory and Holomorphic Dynamics
Vienna - September 29, 2015

Fatou's function

The function

$$f(z) = z + 1 + e^{-z}$$

was first studied by Fatou in 1926.
$F(f)$ consists of one invariant component (Baker domain) which contains the right half-plane.

Fatou's function

The function

$$f(z) = z + 1 + e^{-z}$$

was first studied by Fatou in 1926.
$F(f)$ consists of one invariant component (Baker domain) which contains the right half-plane.
$J(f)$ consists of an uncountable union of curves in the left half-plane (Cantor bouquet).

Fatou's function

The function
$$f(z) = z + 1 + e^{-z}$$
was first studied by Fatou in 1926.
$F(f)$ consists of one invariant component (Baker domain) which contains the right half-plane.
$J(f)$ consists of an uncountable union of curves in the left half-plane (Cantor bouquet).



The escaping set

The set

$$I(f) = \{z \in \mathbb{C} : f^n(z) \to \infty \text{ as } n \to \infty\}$$

is called the **escaping set**.

The escaping set

The set

$$I(f) = \{z \in \mathbb{C} : f^n(z) \to \infty \text{ as } n \to \infty\}$$

is called the **escaping set**.

The **fast escaping set**, $A(f) \subset I(f)$, consists of the points that go to infinity as quickly as possible under iteration.

$$A(f) = \{z \in \mathbb{C} : f^n(z) \to \infty \text{ as quickly as possible}\}.$$

The escaping set

For Fatou's function:

- $I(f)$ consists of the Baker domain and the curves in $J(f)$ except for some of their endpoints;

The escaping set

For Fatou's function:

- $I(f)$ consists of the Baker domain and the curves in $J(f)$ except for some of their endpoints;

- $A(f)$ consists of the curves in $J(f)$ except for some of their endpoints.

Spiders' webs

Rippon and Stallard showed that for many transcendental entire functions the escaping set has a structure called a **spider's web**.

Definition 1

A set E is an (infinite) spider's web if:

1) E is connected and
2) $\exists$ a sequence $(G_n), n \in \mathbb{N}$, of bounded, simply connected domains such that
    - $G_n \subset G_{n+1}, n \in \mathbb{N}$,
    - $\partial G_n \subset E, n \in \mathbb{N}$,
    - $\cup_{n \in \mathbb{N}} G_n = \mathbb{C}$.

Spiders' webs

Rippon and Stallard showed that for many transcendental entire functions the escaping set has a structure called a **spider's web**.

Definition 1

A set E is an (infinite) spider's web if:

1) E is connected and
2) $\exists$ a sequence (G_n), $n \in \mathbb{N}$, of bounded, simply connected domains such that
   - $G_n \subset G_{n+1}, n \in \mathbb{N}$,
   - $\partial G_n \subset E, n \in \mathbb{N}$,
   - $\cup_{n \in \mathbb{N}} G_n = \mathbb{C}$.



Spiders' webs

- Rippon and Stallard showed that when $I(f)$ **contains** a SW then it **is** a SW.
- In most examples we show that $A(f)$ is a SW which implies that $I(f)$ is a SW.
- There exists a complicated example of a function for which $I(f)$ is a SW whereas $A(f)$ is not, due to Rippon and Stallard.

Fatou's web

Theorem 2

Let $f(z) = z + 1 + e^{-z}$*. Then* $I(f)$ *is a SW.*

Fatou's web

Theorem 2

Let $f(z) = z + 1 + e^{-z}$. *Then* $I(f)$ *is a SW.*

Sketch of Proof.

Idea: Use a more general result which implies Theorem 2.

Fatou's web

Theorem 2

Let $f(z) = z + 1 + e^{-z}$. Then $I(f)$ is a SW.

Sketch of Proof.

Idea: Use a more general result which implies Theorem 2.

Let f be a t.e.f. and (a_n) be a positive sequence such that:

(1) $a_n \to \infty$ as $n \to \infty$,

(2) the disc $D(0, a_n)$ contains a periodic cycle of f, for all $n \in \mathbb{N}$.

Consider the set

$$I(f, (a_n)) = \{z \in \mathbb{C} : |f^n(z)| \geq a_n, n \in \mathbb{N}\}.$$

Fatou's web

Theorem 2

Let $f(z) = z + 1 + e^{-z}$. Then $I(f)$ is a SW.

Sketch of Proof.
Idea: Use a more general result which implies Theorem 2.
Let f be a t.e.f. and (a_n) be a positive sequence such that:

(1) $a_n \to \infty$ as $n \to \infty$,

(2) the disc $D(0, a_n)$ contains a periodic cycle of f, for all $n \in \mathbb{N}$.

Consider the set

$$I(f, (a_n)) = \{z \in \mathbb{C} : |f^n(z)| \geq a_n, n \in \mathbb{N}\}.$$

Theorem 3

Let f be a t.e.f. If (a_n) satisfies (1), (2) and $I(f, (a_n))^c$ has a bounded component, then $I(f)$ is a SW.

Fatou's web

Now we apply Theorem 3. Take

$$a_n = \frac{n+6}{2}, n \in \mathbb{N}.$$

Then

(1) $(n+6)/2 \to \infty$ as $n \to \infty$,

(2) $D(0, ((n+6)/2)) \supset D(0, 7/2) \supset \pm\pi i, n \in \mathbb{N}$, and

(3) All the components of $I(f, ((n+6)/2))^c$ are bounded.

Hence Theorem 3 $\Rightarrow$ Theorem 2. $\blacksquare$

Fatou's web



Endpoints

In 1988 Mayer showed that for the exponential family $f_a(z) = e^z + a, a < -1$, the set of all endpoints of $J(f_a)$ is totally disconnected whereas the union of the endpoints with ∞ is a connected set.
Alhabib and Rempe-Gillen recently showed that the same result holds for the set of escaping endpoints of $J(f_a)$.

Endpoints

In 1988 Mayer showed that for the exponential family $f_a(z) = e^z + a, a < -1$, the set of all endpoints of $J(f_a)$ is totally disconnected whereas the union of the endpoints with ∞ is a connected set.
Alhabib and Rempe-Gillen recently showed that the same result holds for the set of escaping endpoints of $J(f_a)$.

The Julia set for Fatou's function is also a Cantor bouquet and hence we can consider the set of endpoints of $J(f)$, which we denote by $E(f)$. Mayer's result holds also for Fatou's function.

Theorem 4

Let $f(z) = z + 1 + e^{-z}$. Then $E(f)$ is totally disconnected but $E(f) \cup \{\infty\}$ is connected.

The proof is based on a result of Barański.

Endpoints

The fact that $I(f)$ is a SW for Fatou's function leads to a result about the non-escaping endpoints of $J(f)$, $\hat{E}(f) = I(f) \setminus E(f)$.

Theorem 5

Let $f(z) = z + 1 + e^{-z}$. Then $\hat{E}(f) \cup \{\infty\}$ is totally disconnected.

Endpoints

The fact that $I(f)$ is a SW for Fatou's function leads to a result about the non-escaping endpoints of $J(f)$, $\hat{E}(f) = I(f) \setminus E(f)$.

Theorem 5

Let $f(z) = z + 1 + e^{-z}$. Then $\hat{E}(f) \cup \{\infty\}$ is totally disconnected.

Proof.

Suppose there is a non-trivial component of $\hat{E}(f) \cup \{\infty\}$. Since $I(f)$ is a SW, any non-escaping endpoint is separated from ∞ by a 'loop' in $I(f)$ and so this component must lie in $\hat{E}(f) \subset E(f)$. Since, by Theorem 4, $E(f)$ is totally disconnected,we obtain a contradiction. ∎

THANK YOU!