

# The Dolgopyat inequality for non-Markov maps in BV.

Henk Bruin (University of Vienna)

joint work with

Dalia Terhesiu (University of Exeter)

Vienna, May 2016

# Dolgopyat inequality for the twisted transfer operator

- ▶  $F$  is expanding non-Markov interval map;
- ▶  $\varphi$  is a piecewise  $C^1$  roof function;
- ▶  $\mathcal{L}$  is the transfer operator, with twisted version

$$\mathcal{L}_s v = \mathcal{L}(e^{s\varphi} v), \quad s = \sigma + ib.$$

**Theorem:** Under appropriate assumptions (to be discussed later) there exist  $A, b_0 \geq 1$  and  $\varepsilon, \gamma \in (0, 1)$  such that

$$\|\mathcal{L}_s^n\|_b \leq \gamma^n$$

for all  $|\sigma| < \varepsilon$ ,  $|b| > b_0$  and  $n \geq A \log |b|$ , where  $\|\cdot\|_b$  is a weighted version of the BV-norm.

## Previous results

The tool (cancellation mechanism) comes from Chernov and Dolgopyat's work to prove exponential mixing for certain Anosov flows.

- ▶ Baladi & Vallée [2005] for general setting of suspension semiflows over p.w.  $C^2$  Markov maps with p.w.  $C^1$  roof.
- ▶ Avila, Gouëzel & Yoccoz [2006] for Teichmüller flows.
- ▶ Araújo & Melbourne [2015] for suspension semiflows over p.w.  $C^{1+\alpha}$  Markov maps with p.w.  $C^1$  roof (to treat the Lorenz flow).
- ▶ Eslami [2015] stretched exponential mixing for skew-products on  $\mathbb{T}^2$  with non-Markov p.w.  $C^{1+\alpha}$  base map and p.w.  $C^1$  roof.
- ▶ Butterley & Eslami [2015] exponential mixing for skew-products on the torus with non-Markov base map with finitely many branches and p.w.  $C^2$  roof.

## The map $F$

Let  $F : Y \rightarrow Y$  be an AFU map for  $Y = [0, 1]$ , i.e.:

- ▶ **Uniformly expanding:**  $|F'| \geq \rho_0 > 1$ ,
- ▶ **Adler's distortion condition:**  $|F''|/|F'|^2$  uniformly bounded.
- ▶ possibly non-Markov, countably many branches, but with **Finite image partition:** Let  $\alpha$  be the partition into maximal intervals of continuity. Then

$X_1 := \cup\{\partial Fa : a \in \alpha\}$  is a finite set.

## The map $F$

Let  $F : Y \rightarrow Y$  be an AFU map for  $Y = [0, 1]$ , i.e.:

- ▶ **Uniformly expanding:**  $|F'| \geq \rho_0 > 1$ ,
- ▶ **Adler's distortion condition:**  $|F''|/|F'|^2$  uniformly bounded.
- ▶ possibly non-Markov, countably many branches, but with **Finite image partition:** Let  $\alpha$  be the partition into maximal intervals of continuity. Then

$$X_1 := \cup\{\partial F a : a \in \alpha\} \text{ is a finite set.}$$

Therefore  $F^n$  has a finite image partition too, and

$$X_n = \cup\{\partial F^n a : a \in \alpha_n\}, \quad \alpha_n = \bigvee_{i=0}^{n-1} F^{-i} \alpha$$

has cardinality  $\#X_n \leq n \#X_1$ .

# Roof function $\varphi$

Let  $\mathcal{H}_n$  be the collection of inverse branches of  $F^n$ .

Let  $\varphi : Y \rightarrow \mathbb{R}$  be  $C^1$  such that

- ▶  $\sup_{h \in \mathcal{H}_1} \sup_{x \in \text{dom}(h)} |(\varphi \circ h)'(x)| < \infty.$
- ▶ There is  $\varepsilon_0 > 0$  such that

$$\sup_{x \in Y} \sup_{h \in \mathcal{H}_1, x \in \text{dom}(h)} |h'(x)| e^{\varepsilon_0 \varphi \circ h(x)} < \infty.$$

# Roof function $\varphi$

Let  $\mathcal{H}_n$  be the collection of inverse branches of  $F^n$ .

Let  $\varphi : Y \rightarrow \mathbb{R}$  be  $C^1$  such that

- ▶  $\sup_{h \in \mathcal{H}_1} \sup_{x \in \text{dom}(h)} |(\varphi \circ h)'(x)| < \infty.$
- ▶ There is  $\varepsilon_0 > 0$  such that

$$\sup_{x \in Y} \sup_{h \in \mathcal{H}_1, x \in \text{dom}(h)} |h'(x)| e^{\varepsilon_0 \varphi \circ h(x)} < \infty.$$

This is used for “moving the contour to  $\Re s > 0$ ” (to prove exponential mixing). Without it, work on imaginary axis in renewal theory context to prove polynomial mixing.

## Transfer operator $\mathcal{L}$

The transfer operator associated to  $F$  is

$$\mathcal{L} : L^1(Y, \text{Leb}) \rightarrow L^1(Y, \text{Leb}).$$

For  $s = \sigma + ib \in \mathbb{C}$ , let  $\mathcal{L}_s$  be the twisted version of  $\mathcal{L}$ :

$$\mathcal{L}_s^n v = \sum_{h \in \mathcal{H}_n} e^{s\varphi_n \circ h} |h'|_v \circ h, \quad n \geq 1,$$

for  $\varphi_n = \sum_{i=0}^{n-1} \varphi \circ F^i$ .



# BV functions

Let  $\text{Var}_Y v$  be the total variation of  $v : Y \rightarrow \mathbb{C}$ .

For  $b \in \mathbb{R}$  define the norm

$$\|v\|_b = \frac{1}{1 + |b|} \text{Var}_Y v + \|v\|_1.$$

Throughout we will work with the Banach space

$$B = \{v : Y \rightarrow \mathbb{C} : \|v\|_b < \infty\}.$$

## Dolgopyat inequality

**Theorem:** Under the above + **additional assumptions**, including **UNI**, there exist  $A, b_0 \geq 1$  and  $\varepsilon, \gamma \in (0, 1)$  such that

$$\|\mathcal{L}_s^n\|_b \leq \gamma^n.$$

for all  $|\sigma| < \varepsilon$ ,  $|b| > b_0$  and  $n \geq A \log |b|$ .

## Dolgopyat inequality

**Theorem:** Under the above + **additional assumptions**, including **UNI**, there exist  $A, b_0 \geq 1$  and  $\varepsilon, \gamma \in (0, 1)$  such that

$$\|\mathcal{L}_s^n\|_b \leq \gamma^n.$$

for all  $|\sigma| < \varepsilon$ ,  $|b| > b_0$  and  $n \geq A \log |b|$ .

**Corollary:** For every  $\omega \in (0, 1)$  there exists  $b_0$  such that

$$\|(I - \mathcal{L}_s)^{-1}\|_b \leq |b|^\omega.$$

for all  $|\sigma| < \varepsilon$ ,  $|b| > b_0$  and  $n \geq A \log |b|$ .

## Additional assumptions

1. We use an iterate  $k$  large enough to kill various constants;
2. Let  $P_k$  be the image partition of  $F^k$ . Assume

$$\min_{p \in P_k} \text{Leb}(p) > C \rho_0^{-k/4},$$

where  $C$  depends the leading eigenfunction  $f_\sigma$  of  $\mathcal{L}_\sigma$ .

## Additional assumptions

1. We use an iterate  $k$  large enough to kill various constants;
2. Let  $P_k$  be the image partition of  $F^k$ . Assume

$$\min_{p \in P_k} \text{Leb}(p) > C \rho_0^{-k/4},$$

where  $C$  depends the leading eigenfunction  $f_\sigma$  of  $\mathcal{L}_\sigma$ .

This is trivially satisfied if  $F$  is Markov.

## Additional assumptions

1. We use an iterate  $k$  large enough to kill various constants;
2. Let  $P_k$  be the image partition of  $F^k$ . Assume

$$\min_{p \in P_k} \text{Leb}(p) > C \rho_0^{-k/4},$$

where  $C$  depends the leading eigenfunction  $f_\sigma$  of  $\mathcal{L}_\sigma$ .

This is trivially satisfied if  $F$  is Markov.

For the family  $x \mapsto \beta x + \alpha$ , it holds for Lebesgue-a.e.  $\alpha, \beta$ .

## Additional assumptions

1. We use an iterate  $k$  large enough to kill various constants;
2. Let  $P_k$  be the image partition of  $F^k$ . Assume

$$\min_{p \in P_k} \text{Leb}(p) > C \rho_0^{-k/4},$$

where  $C$  depends the leading eigenfunction  $f_\sigma$  of  $\mathcal{L}_\sigma$ .

This is trivially satisfied if  $F$  is Markov.

For the family  $x \mapsto \beta x + \alpha$ , it holds for Lebesgue-a.e.  $\alpha, \beta$ .

3. **UNI**: For some particular constant  $D > 0$ , and some fixed multiple  $n_0$  of  $k$ :

$$\forall p \in \mathcal{P}_k \exists h_1, h_2 \in \mathcal{H}_{n_0} \quad \inf_{x \in p} |\psi'(x)| \geq D$$

for  $\psi = \varphi_{h_{n_0}} \circ h_1 - \varphi_{h_{n_0}} \circ h_2$ .

## Line of proof

- ▶ Analyze **jump-sizes** and how **discontinuities are created and propagated**;
- ▶ Cancellation lemma within a **particular cone** of pairs  $(u, v)$ ;
- ▶ Invariance of the cone.
- ▶  $L^2$  contraction in the cone.
- ▶ From outside: **exponential contraction to the cone**
- ▶ **Version of** the Lasota-Yorke inequality.



## Jump-sizes

The non-Markov map  $F$  generates discontinuities at certain points  $x \in Y$  with **jump-size** defined as

$$\text{Size } v(x) := \lim_{\delta \rightarrow 0} \sup_{\xi, \xi' \in (x-\delta, x+\delta)} |v(\xi) - v(\xi')|.$$

**Definition:**  $v : Y \rightarrow \mathbb{C}$  has **exponentially decreasing jump-sizes** if

$$\text{Size } v(x) \leq C_0 \rho_0^{-j/4}$$

if  $x \in X_j \setminus X_{j-1}$  and  $v$  is continuous at every  $x \notin \cup_j X_j$ .

(Recall:  $|F'| \geq \rho_0$  and  $C_0$  is fixed in the proof.)

## Jump-sizes

For  $\lambda_\sigma$ ,  $f_\sigma$  eigenvalue resp. eigenfunction of  $\mathcal{L}_\sigma$ , let

$$\tilde{\mathcal{L}}_s v = \frac{1}{\lambda_\sigma f_\sigma} \mathcal{L}_s (f_\sigma v)$$

be the *normalized* version of  $\mathcal{L}_s$ ,  $s = \sigma + ib$ .

**Proposition:** Take  $k$  large such that the additional assumptions 1 & 2 hold, and  $n = 2k$ . If  $u, v$  with  $|v| \leq u$  have exponentially decreasing jump-sizes, then

$$\text{Size } \tilde{\mathcal{L}}_\sigma^n u(x), \text{ Size } \tilde{\mathcal{L}}_s^n v(x) \leq \frac{1}{4} \max_{p \in \mathcal{P}_k} \frac{\sup u|_p}{\inf u|_p} C \rho_0^{-j/4} \tilde{\mathcal{L}}_\sigma^n u(x)$$

for each  $x \in X_j \setminus X_{j-1}$ ,  $j > k$ .

## The cone

Define  $\text{Osc}_I v = \sup_{x,y \in I} |v(x) - v(y)|$  and

$$E_I(u) := \sum_{j>k} \rho_0^{-j/4} \sum_{x \in (X_j \setminus X_{j-1}) \cap I^\circ} \limsup_{\xi \rightarrow x} u(\xi)$$

as intended upper bound of the sum of jumps-sizes on  $I$ .

$$\text{Cone}_b := \left\{ (u, v) : 0 < u, 0 \leq |v| \leq u, \right.$$

$u, v$  have exponentially decreasing jump-sizes

and  $\text{Osc}_I v \leq C_1 |b| \text{Leb}(I) \sup u|_I + C_2 E_I(u)$

for all intervals  $I \subset$  single atom of  $\mathcal{P}_k$   $\left. \right\}$ .

( $C_1$  and  $C_2$  are fixed in the proof.)

# Invariance of the cone

**Lemma:** Assume  $|b| \geq 2$ ,  $n_0$  a large multiple of  $k$ . Then  $Cone_b$  is invariant under

$$(u, v) \mapsto (\tilde{\mathcal{L}}_\sigma^{n_0}(\chi u), \tilde{\mathcal{L}}_s^{n_0} v),$$

where  $\chi = \chi(b, u, v) \in C^1(Y, [0, 1])$  comes from the “cancellation lemma”.

## BV functions outside the cone.

BV functions can have discontinuities at  $x \notin \cup_j X_j$ , but their jump-sizes decrease exponentially under iteration of  $\mathcal{L}_s^{n_0}$ .

## BV functions outside the cone.

BV functions can have discontinuities at  $x \notin \cup_j X_j$ , but their jump-sizes decrease exponentially under iteration of  $\mathcal{L}_s^{n_0}$ .

**Proposition:** There exists  $\varepsilon \in (0, 1)$  such that for all  $s = \sigma + ib$ ,  $0 \leq \sigma < \varepsilon$ ,  $|b| \geq b_0$ , and all  $v \in \text{BV}$  satisfying

$$\text{Var}_Y v \leq C_3 |b|^2 \rho_0^{mn_0/4} \|v\|_1,$$

there exists a pair  $(u_{mn_0}, w_{mn_0}) \in \text{Cone}_b$  such that

$$\|\tilde{\mathcal{L}}_s^{mn_0} v - w_{mn_0}\|_\infty \leq 2C_4 \rho^{-mn_0} |b| \|v\|_\infty$$

where  $\|w_{mn_0}\|_\infty \leq \|v\|_\infty$ .

## Lasota-Yorke

The spaces  $(\text{BV}, L^1)$  form an adapted pair, but for unbounded roof function  $\varphi$ , the operator  $\mathcal{L}_s : L^1 \rightarrow L^1$  is **not** bounded when  $\Re(s) = \sigma > 0$ . Therefore, the usual Lasota-Yorke inequality fails.

**Proposition:** Choose  $k$  sufficiently large. Define






$$\Lambda_\sigma = \lambda_{2\sigma}^{1/2} / \lambda_\sigma \quad \lambda_\sigma \text{ leading eigenvalue of } \mathcal{L}_\sigma.$$

Then there exist  $\varepsilon > 0$  and  $c > 0$  such that for all  $s = \sigma + ib$  with  $|\sigma| < \varepsilon$  and  $b \in \mathbb{R}$ ,

$$\text{Var}_Y(\tilde{\mathcal{L}}_s^{nk} v) \leq \rho_0^{-nk/4} \text{Var}_Y v + c(1 + |b|) \Lambda_\sigma^{nk} (\|v\|_\infty \|v\|_1)^{1/2},$$

for all  $v \in \text{BV}(Y)$  and all  $n \geq 1$ .

# References

-  V. Araújo, I. Melbourne, Exponential decay of correlations for non-uniformly hyperbolic flows with a  $C^{1+\alpha}$  stable foliation. Preprint 2015 arXiv:1504.04316, to appear in *Annales Henri Poincaré*.
-  A. Avila, S. Gouëzel, J.-C. Yoccoz, Exponential mixing for the Teichmüller flow, *Publ. Math. Inst. Hautes Études Sci.* **104** (2006), 143–211.
-  V. Baladi, B. Vallée, Exponential decay of correlations for surface semi-flows without finite Markov partitions, *Proc. Amer. Math. Soc.* **133** (2005), 865–874.
-  O. Butterley, P. Eslami, Exponential mixing for skew products with discontinuities, Preprint 2015 arXiv:1405.7008, to appear in *Trans. of the AMS*.
-  P. Eslami, Stretched-exponential mixing for  $C^{1+\alpha}$  skew products with discontinuities, *Ergod. Th. & Dynam. Sys.*, published online July 2015.