

Matching for discontinuous interval maps;
its consequences and self-similarity of parameter
space.

Henk Bruin (University of Vienna)

joint with

Carlo Carminati (University of Pisa)

Stefano Marmi (Scuola Normale di Pisa)

Alessandro Profeti (University of Pisa)

explaining observations in a paper by


V. Botella-Soler, J. A. Oteo, J. Ros, and P. Glendinning

Leoben, March 2017

β -transformations

The β -transformation is defined as

$$x \mapsto \beta x \pmod{1}$$




For $|\beta| > 1$, T_β has an acip μ .

β -transformations

The β -transformation is defined as

$$x \mapsto \beta x \pmod{1}$$



For $|\beta| > 1$, T_β has an acip μ .






Figure: Density $\frac{d\mu}{dx}$ for $\beta = \frac{1}{2}(\sqrt{5} + 1)$ and $\beta = \sqrt[3]{7}$.

The density is **only** locally constant, if there is a Markov partition.

The map T_β


$$T_\beta(x) = \begin{cases} T_\beta^-(x) = x + 2 & \text{if } x \leq 0, \\ T_\beta^+(x) = \beta - 2x & \text{if } x \geq 0. \end{cases}$$



T_β preserves the $[\beta - \max\{2, \beta\}, \max\{2, \beta\}]$ and some iterate is uniformly expanding. Therefore T_β admits an acip.

The map T_β

$$T_\beta(x) = \begin{cases} T_\beta^-(x) = x + 2 & \text{if } x \leq 0, \\ T_\beta^+(x) = \beta - 2x & \text{if } x \geq 0. \end{cases}$$



T_β preserves the $[\beta - \max\{2, \beta\}, \max\{2, \beta\}]$ and some iterate is uniformly expanding. Therefore T_β admits an acip.





Figure: Invariant density for the T_β : left $\beta = \frac{1}{2}(\sqrt{5} + 1)$ right: $\beta = \sqrt[3]{7}$.

Not Markov but Matching

For the family T_β , there is no Markov partition in general, but something called **matching** takes can occur:

Definition: There is **matching** if there are iterates $\kappa_\pm > 0$ such that

$$T^{\kappa_-}(0^-) = T^{\kappa_+}(0^+) \text{ and derivatives } DT^{\kappa_-}(0^-) = DT^{\kappa_+}(0^+)$$

Not Markov but Matching

For the family T_β , there is no Markov partition in general, but something called **matching** takes can occur:

Definition: There is **matching** if there are iterates $\kappa_\pm > 0$ such that

$$T^{\kappa_-}(0^-) = T^{\kappa_+}(0^+) \text{ and derivatives } DT^{\kappa_-}(0^-) = DT^{\kappa_+}(0^+)$$

The **pre-matching partition** plays the role of Markov partition:

$$\{T^j(0^-)\}_{j=0}^{\kappa_- - 1} \cup \{T^j(0^+)\}_{j=0}^{\kappa_+ - 1};$$

Not Markov but Matching

For the family T_β , there is no Markov partition in general, but something called **matching** takes can occur:

Definition: There is **matching** if there are iterates $\kappa_\pm > 0$ such that

$$T^{\kappa_-}(0^-) = T^{\kappa_+}(0^+) \text{ and derivatives } DT^{\kappa_-}(0^-) = DT^{\kappa_+}(0^+)$$

The **pre-matching partition** plays the role of Markov partition:

$$\{T^j(0^-)\}_{j=0}^{\kappa_- - 1} \cup \{T^j(0^+)\}_{j=0}^{\kappa_+ - 1};$$

Theorem: If T has matching, then $\rho = \frac{d\mu}{dx}$ is constant on each element of the pre-matching partition.

Monotonicity of entropy

Numerical illustration for the metric entropy:





Figure: Entropy $h_\mu(T_\beta)$ for $\beta \in [4.6, 6]$ (l) and $\beta \in [5.29, 5.33]$ (r).

Monotonicity of entropy

Definition: The **matching index** is $\Delta = \kappa_+ - \kappa_-$.


Theorem: Topological and metric entropy are

$$h_\mu(T_\beta) \text{ and } h_{top}(T_\beta) \text{ are } \begin{cases} \text{increasing} & \text{if } \Delta < 0; \\ \text{constant} & \text{if } \Delta = 0; \\ \text{decreasing} & \text{if } \Delta > 0, \end{cases}$$

as function of β within matching intervals.

The α -continued fraction map T_α .


A generalization of the Gauß map stems from Nakada (and Natsui).



The α -continued fraction map T_α .

A generalization of the Gauß map stems from Nakada (and Natsui).


branches: $|\frac{1}{x}| + n$,
 $n \in \mathbb{Z}$, on $[-1, 1]$



The α -continued fraction map T_α .

A generalization of the Gauß map stems from Nakada (and Natsui).


branches: $|\frac{1}{x}| + n$,
 $n \in \mathbb{Z}$, on $[-1, 1]$



The α -continued fraction map T_α .

A generalization of the Gauß map stems from Nakada (and Natsui).

branches: $|\frac{1}{x}| + n$,
 $n \in \mathbb{Z}$, on $[-1, 1]$



The α -continued fraction map T_α .

A generalization of the Gauß map stems from Nakada (and Natsui).

branches: $|\frac{1}{x}| + n$,
 $n \in \mathbb{Z}$, on $[-1, 1]$




Figure: $T_\alpha : [\alpha - 1, \alpha] \rightarrow [\alpha - 1, \alpha]$, $x \mapsto |\frac{1}{x}| - \lfloor \frac{1}{x} + 1 - \alpha \rfloor$.

All of them have invariant densities (infinite if $\alpha = 0$).

The α -continued fraction map T_α .

A generalization of the Gauß map stems from Nakada (and Natsui).

branches: $|\frac{1}{x}| + n$,
 $n \in \mathbb{Z}$, on $[-1, 1]$




Figure: $T_\alpha : [\alpha - 1, \alpha] \rightarrow [\alpha - 1, \alpha]$, $x \mapsto |\frac{1}{x}| - \lfloor \frac{1}{x} + 1 - \alpha \rfloor$.

All of them have invariant densities (infinite if $\alpha = 0$).

Matching of the orbits of α and $\alpha - 1$ occurs for a.e. $\alpha \in [0, 1]$.

α -continued fractions and the Mandelbrot set





Figure: From a paper by Bonanno, Carminati, Isola and Tiozzo: [The non-matching set and the real antenna of Mandelbrot set](#)

Change of coordinates

For fixed slope $s > 1$, take:

$$Q_\gamma(x) = \begin{cases} x + 1, & x \leq \gamma \\ 1 + s(1 - x), & x > \gamma \end{cases}$$




For $s = 2$, Q_γ is conjugate to T_β above via

$$H \circ Q_\gamma = T_\beta \circ H \quad \text{with } H(x) = 2(x - \gamma), \beta := 2(1 + s)(1 - \gamma).$$

Change of coordinates

For fixed slope $s > 1$, take:

$$Q_\gamma(x) = \begin{cases} x + 1, & x \leq \gamma \\ 1 + s(1 - x), & x > \gamma \end{cases}$$



For $s = 2$, Q_γ is conjugate to T_β above via

$$H \circ Q_\gamma = T_\beta \circ H \quad \text{with } H(x) = 2(x - \gamma), \beta := 2(1 + s)(1 - \gamma).$$

Advantages of Q_γ :

- ▶ $x = 1$ is fixed for all $s \in \mathbb{R}$ and $\gamma < 1$;
- ▶ For integer $s \geq 2$, every point ps^{-m} , $p, m \in \mathbb{N}$, eventually maps to 1;
- ▶ Therefore matching occurs whenever $\gamma = ps^{-m}$;
- ▶ Matching occurs on an open dense set!

Matching is Lebesgue typical

Theorem: Q_γ has matching for Lebesgue-a.e. γ , but the set \mathcal{E} of non-matching parameters has Hausdorff dimension 1.


Matching is Lebesgue typical

Theorem: Q_γ has matching for Lebesgue-a.e. γ , but the set \mathcal{E} of non-matching parameters has Hausdorff dimension 1.

Let $g(x) := s(1-x) \bmod 1$ and $R : (0, 1) \rightarrow (0, 1)$ be the first return of $Q_\gamma^k(x)$ to $[0, 1)$.

Lemma:

$$R(x) = \begin{cases} g(x) & \text{if } x \in (0, \gamma) \\ g^2(x) & \text{if } x \in (\gamma, 1) \end{cases}$$



On the proof of “Matching is Lebesgue typical”


Lemma: For fixed $\gamma \in [0, 1]$, the following conditions are equivalent:

- (i) $g^k(\gamma) < \gamma$ for some $k \in \mathbb{N}$;
- (ii) matching holds for γ .

In other words, the bifurcation set is

$$\mathcal{E} = \{\gamma \in [0, 1] : g^k(\gamma) \geq \gamma \forall k \in \mathbb{N}\}.$$


On the proof of “Matching is Lebesgue typical”



Proof of the Theorem:

- ▶ Lebesgue measure is preserved by g , so the Ergodic Theorem implies that $\inf\{g^k(\gamma) : k \geq 1\} = 0$ for a.e. γ . The previous lemma gives that each such $\gamma \notin \mathcal{E}$.


On the proof of “Matching is Lebesgue typical”



Proof of the Theorem:

- ▶ Lebesgue measure is preserved by g , so the Ergodic Theorem implies that $\inf\{g^k(\gamma) : k \geq 1\} = 0$ for a.e. γ . The previous lemma gives that each such $\gamma \notin \mathcal{E}$.
- ▶ Define $K(t) := \{x \in [0, 1] : g^k(x) \geq t \forall k \geq 1\}$.


On the proof of “Matching is Lebesgue typical”



Proof of the Theorem:

- ▶ Lebesgue measure is preserved by g , so the Ergodic Theorem implies that $\inf\{g^k(\gamma) : k \geq 1\} = 0$ for a.e. γ . The previous lemma gives that each such $\gamma \notin \mathcal{E}$.
- ▶ Define $K(t) := \{x \in [0, 1] : g^k(x) \geq t \forall k \geq 1\}$.
- ▶ The Hausdorff dimension $\dim_H(K(t)) \rightarrow 1$ and $\dim_H(K(t) \cap [0, 1]) \rightarrow 1$ as $t \rightarrow 0$.

On the proof of “Matching is Lebesgue typical”



Proof of the Theorem:

- ▶ Lebesgue measure is preserved by g , so the Ergodic Theorem implies that $\inf\{g^k(\gamma) : k \geq 1\} = 0$ for a.e. γ . The previous lemma gives that each such $\gamma \notin \mathcal{E}$.
- ▶ Define $K(t) := \{x \in [0, 1] : g^k(x) \geq t \ \forall k \geq 1\}$.
- ▶ The Hausdorff dimension $\dim_H(K(t)) \rightarrow 1$ and $\dim_H(K(t) \cap [0, 1]) \rightarrow 1$ as $t \rightarrow 0$.
- ▶ Combine this with $\mathcal{E} \cap [0, t] \supset K(t)$.

Monotonicity

Recall the **Monotonicity Theorem** stated for Q_γ :

Topological and metric entropy are

$$h_\mu(Q_\gamma) \text{ and } h_{top}(Q_\gamma) \text{ are } \begin{cases} \text{decreasing} & \text{if } \Delta < 0; \\ \text{constant} & \text{if } \Delta = 0; \\ \text{increasing} & \text{if } \Delta > 0, \end{cases}$$

as function of γ within matching intervals.

Monotonicity

Recall the **Monotonicity Theorem** stated for Q_γ :

Topological and metric entropy are

$$h_\mu(Q_\gamma) \text{ and } h_{top}(Q_\gamma) \text{ are } \begin{cases} \text{decreasing} & \text{if } \Delta < 0; \\ \text{constant} & \text{if } \Delta = 0; \\ \text{increasing} & \text{if } \Delta > 0, \end{cases}$$

as function of γ within matching intervals.

The proof is based on the structure of the first return map R to a neighbourhood J of $z := Q_{\gamma^-}^{\kappa^-}(\gamma_-) = Q_{\gamma^+}^{\kappa^+}(\gamma_+)$ which is **nice**,

Monotonicity

Recall the **Monotonicity Theorem** stated for Q_γ :

Topological and metric entropy are

$$h_\mu(Q_\gamma) \text{ and } h_{top}(Q_\gamma) \text{ are } \begin{cases} \text{decreasing} & \text{if } \Delta < 0; \\ \text{constant} & \text{if } \Delta = 0; \\ \text{increasing} & \text{if } \Delta > 0, \end{cases}$$

as function of γ within matching intervals.

The proof is based on the structure of the first return map R to a neighbourhood J of $z := Q_{\gamma^-}^{\kappa^-}(\gamma_-) = Q_{\gamma^+}^{\kappa^+}(\gamma_+)$ which is **nice**, i.e.:

$$\text{orb}(\partial J) \cap J^\circ = \emptyset.$$


Monotonicity

Lemma: All branches of R are monotone onto, also the branches that contain a preimage $y \in Q_\gamma^{-N}(\gamma)$.

$R = Q_\gamma^\tau$ for first
return time $\tau : [0, 1) \rightarrow \mathbb{N}$

$$\tau(A_-) = N + \kappa_-$$

$$\tau(A_+) = N + \kappa_+$$



Corollary: R preserves Lebesgue measure m .

Monotonicity

Proof-sketch of the monotonicity theorem:

- ▶ $\int_J \tau \, dm h_m(R)$ increases by an amount proportional to $\eta := \Delta \times$ increased proportion of $|A_+|/|A_-|$.
- ▶ Abramov's Formula: $h_\mu(Q_\gamma) = \frac{1}{\int_J \tau \, dm} h_m(R)$.
- ▶ Therefore $h_\mu(Q_\gamma)$ decreases due to increase of η .

Monotonicity

Proof-sketch of the monotonicity theorem:

- ▶ $\int_J \tau \, dm h_m(R)$ increases by an amount proportional to $\eta := \Delta \times$ increased proportion of $|A_+|/|A_-|$.
- ▶ Abramov's Formula: $h_\mu(Q_\gamma) = \frac{1}{\int_J \tau \, dm} h_m(R)$.
- ▶ Therefore $h_\mu(Q_\gamma)$ decreases due to increase of η .
- ▶ Topological entropy is the exponential growth-rate of number of periodic point.
- ▶ As γ moves within a matching interval, periodic points in J don't change,

Monotonicity

Proof-sketch of the monotonicity theorem:

- ▶ $\int_J \tau \, dm h_m(R)$ increases by an amount proportional to $\eta := \Delta \times$ increased proportion of $|A_+|/|A_-|$.
- ▶ Abramov's Formula: $h_\mu(Q_\gamma) = \frac{1}{\int_J \tau \, dm} h_m(R)$.
- ▶ Therefore $h_\mu(Q_\gamma)$ decreases due to increase of η .
- ▶ Topological entropy is the exponential growth-rate of number of periodic point.
- ▶ As γ moves within a matching interval, periodic points in J don't change,
- ▶ but their period increases by Δ as A_+ absorbs them (when they previously belonged to A_-).

Monotonicity

Proof-sketch of the monotonicity theorem:

- ▶ $\int_J \tau \, dm h_m(R)$ increases by an amount proportional to $\eta := \Delta \times$ increased proportion of $|A_+|/|A_-|$.
- ▶ Abramov's Formula: $h_\mu(Q_\gamma) = \frac{1}{\int_J \tau \, dm} h_m(R)$.
- ▶ Therefore $h_\mu(Q_\gamma)$ decreases due to increase of η .
- ▶ Topological entropy is the exponential growth-rate of number of periodic point.
- ▶ As γ moves within a matching interval, periodic points in J don't change,
- ▶ but their period increases by Δ as A_+ absorbs them (when they previously belonged to A_-).
- ▶ Hence the topological entropy decreases accordingly.

Pseudo-centers

Motivation: Find exact formulas for matching intervals J and their matching indices Δ for slope $s = 2$ (also works for $2 \leq s \in \mathbb{N}$).

Pseudo-centers

Motivation: Find exact formulas for matching intervals J and their matching indices Δ for slope $s = 2$ (also works for $2 \leq s \in \mathbb{N}$).

Let \mathbb{Q}_{dyd} be the set of **dyadic rationals** in $(0, 1]$.

Definition The **pseudocenter** of an interval $J \subset (0, 1)$ is the (unique) dyadic rational $\xi \in \mathbb{Q}_{\text{dyd}}$ with minimal denominator.

Pseudo-centers

Motivation: Find exact formulas for matching intervals J and their matching indices Δ for slope $s = 2$ (also works for $2 \leq s \in \mathbb{N}$).

Let \mathbb{Q}_{dyd} be the set of **dyadic rationals** in $(0, 1]$.

Definition The **pseudocenter** of an interval $J \subset (0, 1)$ is the (unique) dyadic rational $\xi \in \mathbb{Q}_{\text{dyd}}$ with minimal denominator.

Definition

- ▶ For binary string u , let \check{u} be the bitwise negation of u .
- ▶ For $\xi \in \mathbb{Q}_{\text{dyd}} \setminus \{1\}$ and let w be the shortest **even** binary expansion of ξ and v be the shortest **odd** binary expansion of $1 - \xi$.

Pseudo-centers

Motivation: Find exact formulas for matching intervals J and their matching indices Δ for slope $s = 2$ (also works for $2 \leq s \in \mathbb{N}$).

Let \mathbb{Q}_{dyd} be the set of **dyadic rationals** in $(0, 1]$.

Definition The **pseudocenter** of an interval $J \subset (0, 1)$ is the (unique) dyadic rational $\xi \in \mathbb{Q}_{\text{dyd}}$ with minimal denominator.

Definition

- ▶ For binary string u , let \check{u} be the bitwise negation of u .
- ▶ For $\xi \in \mathbb{Q}_{\text{dyd}} \setminus \{1\}$ and let w be the shortest **even** binary expansion of ξ and v be the shortest **odd** binary expansion of $1 - \xi$.
- ▶ Define the interval $I_\xi := (\xi_L, \xi_R)$ containing ξ where,
- ▶ $\xi_L := \overline{.v\check{v}}$, $\xi_R := \overline{.w}$.
- ▶ Also define the “degenerate” interval $I_1 := (2/3, +\infty)$.

Pseudo-centers

In short: $I_\xi := (\xi_L, \xi_R)$ with $\xi_L := \overline{\check{v}v}$, $\xi_R := \overline{w}$.

Pseudo-centers

In short: $I_\xi := (\xi_L, \xi_R)$ with $\xi_L := \overline{.v\check{v}}$, $\xi_R := \overline{.w}$.

If $\xi = 1/2$ then $w = 10$, $v = 1$ and $\xi_L = \overline{.0\check{1}}$, $\xi_R = \overline{.1\check{0}}$.

$$(01) \quad w = u01 \Rightarrow \xi_L = \overline{.u001\check{u}110};$$

$$(11) \quad w = u11 \Rightarrow \xi_L = \overline{.u101\check{u}010};$$

$$(010) \quad w = u010 \Rightarrow \xi_L = \overline{.u00\check{u}11};$$

$$(110) \quad w = u110 \Rightarrow \xi_L = \overline{.u10\check{u}01}.$$

Pseudo-centers

In short: $I_\xi := (\xi_L, \xi_R)$ with $\xi_L := \overline{.v\bar{v}}$, $\xi_R := \overline{.w}$.

If $\xi = 1/2$ then $w = 10$, $v = 1$ and $\xi_L = \overline{.0\bar{1}}$, $\xi_R = \overline{.1\bar{0}}$.

$$(01) \quad w = u01 \Rightarrow \xi_L = \overline{.u001\check{u}110};$$

$$(11) \quad w = u11 \Rightarrow \xi_L = \overline{.u101\check{u}010};$$

$$(010) \quad w = u010 \Rightarrow \xi_L = \overline{.u00\check{u}11};$$

$$(110) \quad w = u110 \Rightarrow \xi_L = \overline{.u10\check{u}01}.$$

ξ	ξ_R	ξ_L
$\frac{1}{2} = .10$	$\frac{2}{3} = \overline{.10}$	$\frac{1}{3} = \overline{.01}$
$\frac{1}{4} = .01$	$\frac{1}{3} = \overline{.01}$	$\frac{2}{9} = \overline{.001110}$
$\frac{7}{32} = .001110$	$\frac{2}{9} = \overline{.001110}$	$\frac{7}{33} = \overline{.0011011001}$
$\frac{3}{16} = .0011$	$\frac{1}{5} = \overline{.0011}$	$\frac{2}{11} = \overline{.0010111010}$
$\frac{9}{64} = .001001$	$\frac{1}{7} = \overline{.001}$	$\frac{4334}{16383} = \overline{.00100011101110}$
$\frac{1}{8} = .0010$	$\frac{2}{15} = \overline{.0010}$	$\frac{1}{9} = \overline{.000111}$

Pseudo-centers

Theorem:

- ▶ All matching intervals have the form I_ξ , where $\xi \in \mathbb{Q}_{\text{dyd}}$ are precisely the pseudo-centers of the components of $[0, \frac{2}{3}] \setminus \mathcal{E}$.
- ▶ The matching index is

$$\Delta(\xi) = \frac{3}{2}(|w|_0 - |w|_1),$$

where $|w|_a$ is the number of symbols a in w (the shortest **even** binary expansion of ξ).

Pseudo-centers

Theorem:

- ▶ All matching intervals have the form I_ξ , where $\xi \in \mathbb{Q}_{\text{dyd}}$ are precisely the pseudo-centers of the components of $[0, \frac{2}{3}] \setminus \mathcal{E}$.
- ▶ The matching index is

$$\Delta(\xi) = \frac{3}{2}(|w|_0 - |w|_1),$$

where $|w|_a$ is the number of symbols a in w (the shortest **even** binary expansion of ξ).

Proposition: If $g^k(\gamma) \geq \frac{1}{6}$, then $|w_0| = |w_1|$. In particular, all matching intervals in $(\frac{1}{6}, \frac{2}{3})$ have matching index $\Delta = 0$.

Pseudo-centers





Figure: Entropies $h_{top}(T_\beta)$ and $h_\mu(T_\beta)$ for $\beta \in [0, 6.5]$.

Remark: This proposition explains constant entropy on all matching intervals in $(\frac{1}{6}, \frac{2}{3})$. A **no devil's staircase** argument would give:

$$h_\mu(Q_\gamma) = \log\left(\frac{1 + \sqrt{5}}{2}\right) \quad \text{and} \quad h_{top}(Q_\gamma) = \frac{2}{3} \log 2,$$

for all $\gamma \in [\frac{1}{6}, \frac{2}{3}]$.


Pseudo-centers (period doubling)

Pseudo-center $\xi = .w$ (even expansion) and $1 - \xi = .v$ (odd exp).

The **matching interval** is $I_\xi = [\xi_L, \xi_R]$ for $\xi_L = .\check{v}v$ and $\xi_R = .\bar{w}$.


Pseudo-centers (period doubling)

Pseudo-center $\xi = .w$ (even expansion) and $1 - \xi = .v$ (odd exp).
The **matching interval** is $I_\xi = [\xi_L, \xi_R]$ for $\xi_L = .\check{v}\overline{v}$ and $\xi_R = .\overline{w}$.
But ξ_L is also the right end-point of I_{ξ_1} for $\xi_1 = .\check{v}v$. We call this
“**period doubling**”. It repeats countably often, converging to ξ_∞ .



Pseudo-centers (period doubling)

Pseudo-center $\xi = .w$ (even expansion) and $1 - \xi = .v$ (odd exp).
 The **matching interval** is $I_\xi = [\xi_L, \xi_R]$ for $\xi_L = .\check{v}\bar{v}$ and $\xi_R = .\bar{w}$.
 But ξ_L is also the right end-point of I_{ξ_1} for $\xi_1 = .\check{v}v$. We call this
 “**period doubling**”. It repeats countably often, converging to ξ_∞ .



Lemma: The pseudo-center of the next period doubling can be obtained from the previous using the substitution:

$$\chi : \begin{array}{ll} w \mapsto \check{v}v & \check{w} \mapsto v\check{v} \\ v \mapsto vw & \check{v} \mapsto \check{v}\check{w} \end{array} .$$

Thus the limit ξ_∞ has s -adic expansion


$$\xi_\infty = .\check{v}\check{w}v\check{v}vw\check{v}\check{w}vw\check{v}\check{v}\check{w}\dots$$

Pseudo-centers (tuning windows)

Pseudo-center $\xi = .w$ (even expansion) and $1 - \xi = .v$ (odd exp).

The **matching interval** is $I_\xi = [\xi_L, \xi_R]$ for $\xi_L = .\check{v}\check{v}$ and $\xi_R = .\check{w}$.

The **tuning interval** is $T_\xi = [\xi_T, \xi_R]$ for $\xi_T = .\check{v}\check{w}$.




Pseudo-centers (tuning windows)

Pseudo-center $\xi = .w$ (even expansion) and $1 - \xi = .v$ (odd exp).

The **matching interval** is $I_\xi = [\xi_L, \xi_R]$ for $\xi_L = .\check{v}\check{v}$ and $\xi_R = .\bar{w}$.

The **tuning interval** is $T_\xi = [\xi_T, \xi_R]$ for $\xi_T = .\check{v}\check{w}$.




Theorem: Let $K(\xi_T) = \{x : g^k(x) \geq \xi_T \forall k\}$.

Then $x \in K(\xi_T) \cap T_\xi$ if and only if

$$x = .\sigma_1\sigma_2\sigma_3\sigma_4\dots$$

for $\sigma_1 \in \{w, \check{v}\}$, $\sigma_j \in \{w, v, \check{w}, \check{v}\}$
describing a path in the diagram.




Pseudo-centers (tuning windows)

Pseudo-center $\xi = .w$ (even expansion) and $1 - \xi = .v$ (odd exp).

The **matching interval** is $I_\xi = [\xi_L, \xi_R]$ for $\xi_L = .\check{v}\check{v}$ and $\xi_R = .\bar{w}$.

The **tuning interval** is $T_\xi = [\xi_T, \xi_R]$ for $\xi_T = .\check{v}\check{w}$.




Theorem: Let $K(\xi_T) = \{x : g^k(x) \geq \xi_T \forall k\}$.

Then $x \in K(\xi_T) \cap T_\xi$ if and only if


$$x = .\sigma_1\sigma_2\sigma_3\sigma_4\dots$$

for $\sigma_1 \in \{w, \check{v}\}$, $\sigma_j \in \{w, v, \check{w}, \check{v}\}$
describing a path in the diagram.

If $\Delta(\xi) = 0$, all matching in T_ξ is neutral.




Shape of the entropy function



Question: We know that $\gamma \mapsto h(\mu_\gamma)$ is Hölder. Is $\gamma \mapsto h_{top}(Q_\gamma)$ Hölder?


Shape of the entropy function



Question: We know that $\gamma \mapsto h(\mu_\gamma)$ is Hölder. Is $\gamma \mapsto h_{top}(Q_\gamma)$ Hölder?

Conjecture: The neutral tuning windows are exactly the plateaus of (topological and metric) entropy.

Shape of the entropy function




Question: We know that $\gamma \mapsto h(\mu_\gamma)$ is Hölder. Is $\gamma \mapsto h_{top}(Q_\gamma)$ Hölder?

Conjecture: The neutral tuning windows are exactly the plateaus of (topological and metric) entropy.

Conjecture: The shape of the entire entropy function (i.e., pattern of increase/decrease) is repeated in every tuning window T_ξ with $\Delta(\xi) > 0$, and reversed in every tuning window T_ξ with $\Delta(\xi) < 0$.

Shape of the entropy function








Question: We know that $\gamma \mapsto h(\mu_\gamma)$ is Hölder. Is $\gamma \mapsto h_{top}(Q_\gamma)$ Hölder?

Conjecture: The neutral tuning windows are exactly the plateaus of (topological and metric) entropy.

Conjecture: The shape of the entire entropy function (i.e., pattern of increase/decrease) is repeated in every tuning window T_ξ with $\Delta(\xi) > 0$, and reversed in every tuning window T_ξ with $\Delta(\xi) < 0$.

End of the Show

References

-  C. Bonanno, C. Carminati, S. Isola, G. Tiozzo, *Dynamics of continued fractions and kneading sequences of unimodal maps*, Discrete Contin. Dyn. Syst. **33** (2013), no. 4, 1313–1332.
-  V. Botella-Soler, J. A. Oteo, J. Ros, P. Glendinning, *Families of piecewise linear maps with constant Lyapunov exponents*, J. Phys. A: Math. Theor. **46** 125101
-  C. Carminati, G. Tiozzo, *Tuning and plateaux for the entropy of α -continued fractions*, Nonlinearity **26** (2013), no. 4, 1049–1070.
-  H. Nakada, *Metrical theory for a class of continued fraction transformations and their natural extensions*, Tokyo J. Math. **4** (1981), 399–426
-  H. Nakada, R. Natsui, *The non-monotonicity of the entropy of α -continued fraction transformations*, Nonlinearity, **21** (2008), 1207–1225.