Maximal Entropy Random Networks with Given Degree Distribution

Michel Bauer Denis Bernard

Vienna, Preprint ESI 1180 (2002)

July 2, 2002

Supported by the Austrian Federal Ministry of Education, Science and Culture Available via $\rm http://www.esi.ac.at$

Maximal entropy random networks with given degree distribution

Michel Bauer¹ and Denis Bernard²

Service de Physique Théorique, CEA/DSM/SPhT, Unité de recherche associée au CNRS CEA-Saclay, 91191 Gif-sur-Yvette cedex, France

Abstract

Using a maximum entropy principle to assign a statistical weight to any graph, we introduce a model of random graphs with arbitrary degree distribution in the framework of standard statistical mechanics. We compute the free energy and the distribution of connected components. We determine the size of the percolation cluster above the percolation threshold. The conditional degree distribution on the percolation cluster is also given. We briefly present the analogous discussion for oriented graphs, giving for example the percolation criterion.

1 Introduction

The statistical properties of networks, either biological, social or technological, have received a lot of attention recently both experimentally and theoretically, See eg. refs. [2, 6]. One of the most studied features of those networks is the degree distribution, which describes the probability for the vertices to have $0, 1, \cdots$ neighbors. One striking observation is that, in many examples, the degree distribution is large so that the probability to have nneighbors decreases slowly with n. Several models (static or evolving) predict

¹Email: bauer@spht.saclay.cea.fr

²Member of the CNRS; Email : dbernard@spht.saclay.cea.fr

such a behavior. More generally, they contain enough tunable parameters to reproduce almost any degree distribution.

However, the static models are in general not conveniently defined within the language of statistical mechanics (see ref.[2], which motivated our interest in this question). This is for instance the case with the most intuitive proposal [5]: generate independently half edges for each vertex, with the appropriate distribution, and then join the half edges at random. This makes it rather easy to generate random graphs, but does not assign in a simple way a probability to any given simple graph : it is formally complicated to eliminate multiple edges. Another proposal made in [4] has some formal technical similarity with our work but really leads to a different model.

It is moreover obvious, if not always apparent in the literature, that the knowledge of the degree distribution leaves many statistical properties of the graphs undetermined, even if one insists that all vertices are equivalent. This arbitrariness is a problem, because most of the time the models used to fit the behavior of say a communication network are just ingenious constructions : they are not derived from clear basic principles. Such principles may be out of our reach at the moment, and so is a classification of all random graph models with certain apriori properties. Consequently, we propose to use maximum entropy as a criterion to build a model that does not make any a priori bias, incorporating what we know – in this case the degree distribution – but nothing else. Comparison with real networks is a way to get evidence for other striking features that might be overlooked today.

The maximal entropy principle is applied here to deal with constraints on the degree distribution but it can clearly be engineered to deal with other constraints.

This paper is organized as follows :

- Section 2 starts with the main definitions, goes on with a quick reminder on the Molloy-Reed model [1] and continues with the definition of the maximum entropy model. We use it to reformulate the standard Erdös-Renyi random graph model. Then we derive a few general identities valid for the maximal entropy model, and study the distribution of connected components. Our model is a close cousin of the Molloy-Reed model and we make the connexion precise below. Finally we discuss the possibility of numerical simulations.

- Section 3 studies the thermodynamical limit when the number N of sites is large, but the number of edges scales like N, hence the name finite connectivity limit for this regime. We derive the equations that determine

all physical quantities in this regime : free energy, distribution of the number of edges incident at a vertex, ... We then study the connected components, derive the criterion for the existence of a percolation cluster and the formula for its size. Finally, we study the distribution of the number of edges incident at a vertex in the percolation cluster.

- Section 4 analyzes the generalization to oriented graphs, ending with the criterion for the existence of a percolation cluster and the formula for its size.

2 The model

2.1 General definitions

In the following, we shall concentrate on labeled simple unoriented graphs, or equivalently on symmetric 0-1 matrices, with vanishing main diagonal : the matrix element (i, j) is 1 if vertices i and j are connected by an edge and 0 else. So we use the same letter G to denote the graph and its adjacency matrix with matrix elements $G_{i,j}$. In the sequel, unless otherwise stated, the term graph refers to labeled simple unoriented graph. The number of edges of a graph G is denoted by E(G) and the number of vertices by V(G).

The row-sum $\hat{G}_i = \sum_j G_{i,j}$ is the number of neighbors of site *i*. The degree distribution of *G* is the sequence $\tilde{G}_k = \#\{i \text{ such that } \hat{G}_i = k\}$, so that \tilde{G}_0 is the number of isolated points of *G*, \tilde{G}_1 is the number of vertices of *G* with exactly one neighbor, and so on.

Not every integer sequence can appear as the degree distribution of a graph on N vertices : $\tilde{G}_k = 0$ for $k \ge N$, $\sum_k \tilde{G}_k = N$ and $\sum_k k \tilde{G}_k$ is even because this number counts twice the number of edges of G, i.e. $\sum_{i,j} G_{i,j}$. There are other less obvious constraints. We call the sequences that appear as degree distribution of a graph on N vertices N-admissible. There is a relatively simple family of inequalities that characterizes N-admissible sequences, but for instance the (asymptotic) counting of N-admissible sequences is still unknown.

2.2 The Molloy–Reed model

Before we introduce our model, let us describe the method of Molloy and Reed [1] which can be interpreted as a kind of microcanonical version of our model. The idea is quite elegant : for any integer N fix an N-admissible sequence $\{m_{N,k}\}_{k\geq 0}$ and take as probability space the set $\mathcal{G}_{\{m_{N,k}\}}$ of graphs with degree distribution $\{m_{N,k}\}_{k\geq 0}$, endowed with the uniform (counting) probability. By construction, in $\mathcal{G}_{\{m_{N,k}\}}$, the probability that vertex $i \in [1, N]$ has k neighbors is exactly $m_{N,k}/N$.

Molloy and Reed show that if the sequence $\{m_{N,k}/N\}$ converges (uniformly) to a probability distribution $\{\pi_k\}, (\sum_k \pi_k = 1)$, under one technical assumption, the space $\mathcal{G}_{\{m_{N,k}\}}$ converges in an appropriate sense to a random graph ensemble $\mathcal{G}_{\{\pi_k\}}$ on which standard questions can be formulated and answered :

- the probability in $\mathcal{G}_{\{\pi_k\}}$ that a given vertex has k neighbors is - not surprisingly $-\pi_k$,

- Molloy and Reed give a criterion for the presence or absence of a giant component.

Heuristic arguments [5] show that the 'intuitive' model (which does not in general lead to simple graphs) namely "generate independently half edges for each vertex, with distribution $\{\pi_k\}$ and then join the half edges at random", has the same thermodynamical – large N – properties as the Molloy–Reed model.

2.3 The maximum entropy model

To start with, we fix an integer $N \ge 1$, and a probability distribution $\{\pi_{N,k}\}$ $(\sum_{k=1}^{N-1} \pi_{N,k} = 1)$. We want to look for a probability distribution $\{p_G\}$ on the set of graphs on N vertices such that for any vertex i, $\sum_{G;\hat{G}_i=k} p_G = \pi_{N,k}$ where, here and below, the notation means that the sum is restricted to graphs such that \hat{G}_i , the number of neighbors of vertex i in G, equals k. With words, we look for a probability distribution $\{p_G\}$ on the set of graphs on N vertices such that the probability that vertex i has k neighbors is $\pi_{N,k}$. As explained in the introduction, this requirement is far from fixing the probability distribution.

We also want this probability distribution to have no other bias. So we look for a distribution $\{p_G\}$ with maximal entropy ¹.

¹Notice that, if no constraint is imposed, the uniform counting measure has maximal entropy. This measure can be described as follows: the probability of an edge between vertices i and j is 1/2 independently of the presence or absence of any other edge.

Hence we want to maximize $\sum_G p_G \log p_G$ under the constraints

$$\sum_{G;\,\hat{G}_i=k} p_G = \pi_{N,k}$$

which we implement as Lagrange multipliers. The extremum conditions for

$$\sum_{G} p_{G}(\log p_{G}+1) - \lambda(\sum_{G} p_{G}-1) - \sum_{i,k} \lambda_{i,k}(\sum_{G;\hat{G}_{i}=k} p_{G}-\pi_{N,k})$$

are

$$p_G = e^{\lambda + \sum_i \lambda_{i,\hat{G}_i}}, \ \sum_G p_G = 1 \text{ and } \sum_{G; \hat{G}_i = k} p_G = \pi_{N,k}.$$

It is not obvious to us that these equations always have a solution and that this solution is unique and symmetric i.e. $\lambda_{i,k}$ does not depend on i^2 . But as usual in statistical mechanics, we can reverse the logic : we start from an arbitrary sequence of positive numbers $t_k = e^{\lambda_k}$ and define p_G :

$$p_G \equiv e^{\lambda + \sum_i \lambda_{\hat{G}_i}} = e^{\lambda} \prod_k t_k^{\tilde{G}_k} \tag{1}$$

with suitably adjusted λ so as to ensure $\sum_G p_G = 1$.

We define the weight w_G of a graph as

$$w_G \equiv \prod_k t_k^{\tilde{G}_k} = \prod_i t_{\hat{G}_i}$$

and the partition function as the sum of weights

$$Z_N \equiv \sum_G w_G = \sum_G \prod_k t_k^{\tilde{G}_k}$$

Hence $\lambda = -\log Z_N$ and $p_G = w_G/Z_N$.

By construction the probability distribution $\pi_{N,k}$ that vertex *i* has *k* neighbors is *i*-independent. Recall that \tilde{G}_k is the number of vertices with *k* neighbors in *G* so that

$$N \pi_{N,k} = \sum_{i} \sum_{G; \hat{G}_{i} = k} p_{G} = \sum_{G} \tilde{G}_{k} p_{G} = \frac{1}{Z_{N}} \sum_{G} \tilde{G}_{k} w_{G}.$$

But $\tilde{G}_k w_G = \partial w_G / \partial \lambda_k$ so

$$\pi_{N,k} = \frac{1}{N} \frac{\partial \log Z_N}{\partial \lambda_k} = \frac{1}{N} t_k \frac{\partial \log Z_N}{\partial t_k}$$

 $^{^2 \}mathrm{In}$ the large N limit, some spontaneous symmetry breaking might occur. We shall not pursue this questions here.

2.4 The Erdös–Renyi model revisited

As a first application, but also as a preparation to section 3, let us reinterpret the standard Erdös-Renyi random graph model [7] in our framework. Recall that in the Erdös-Renyi model, the edges are described by independent binomial variables, each edge being drawn with probability p. Recall that E(G) denotes the number of edges of a graph G. The probability of the graph G is simply $p^{E(G)}(1-p)^{N(N-1)/2-E(G)}$ which we rewrite as $(1-p)^{N(N-1)/2} \left(\frac{p}{1-p}\right)^{E(G)}$. Now $2E(G) = \sum_k k\tilde{G}_k$. So letting $t_k \equiv \left(\frac{p}{1-p}\right)^{k/2}$ shows that the Erdös-Renyi model is the maximal entropy model such that the probability that a vertex has k neighbors is $\binom{N-1}{k}p^k(1-p)^{N-1-k}$. The average number of neighbors is (N-1)p. In the large N limit, an interesting regime occurs when this number is kept fixed, so that $p \sim \alpha/N$ and α is the control parameter. The important observation is that if p scales like N^{-1} , the parameters t_k scale like $N^{-k/2}$. In section 3, we shall see that indeed generically these scaling relations ensure that $\log Z_N$ scales like N, as any "good" free energy should.

2.5 Useful relations

We establish a few formulæ which will be central in the following discussion.

The sequence $\{Z_N\}_{N\geq 1}$ satisfies a first order functional recursion relation that will prove useful in the subsequent analysis. We define the formal Laurent series $H(\omega, t_0, \dots, t_k, \dots) = \sum_k t_k \omega^{-k}$.

Suppose $N \geq 2$. Then Z_N is the constant (i.e. of degree 0) term in the ω -expansion of the product

$$H(\omega, t_0, \cdots, t_k, \cdots) Z_{N-1}(t_0 + \omega t_1, \cdots, t_j + \omega t_{j+1}, \cdots).$$

The ω -expansion of the product is well-defined because both factors involve at most a finite number of terms of positive degree.

The proof of this relation goes as follows. If G' is a graph on N-1 vertices $1, \dots, N-1$, it can be completed to a graph G on N vertices in the following ways : add vertex N and $k = 0, \dots, N-1$ edges emerging from N. Attach these edges to any k distinct vertices of G'. There is a simple relation between the weights of G and G' because one vertex of degree k has been added (this is taken care of by the term t_k in H), and k vertices in $1, \dots, N-1$ have seen their degree increased by 1 so, in $w_{G'} = \prod_{1}^{N-1} t_{\hat{G}'_{1}}, k$

of the factors t_j are replaced by t_{j+1} (this is taken care of by replacing all t_j 's in Z_{N-1} by $t_j + \omega t_{j+1}$ and expanding to order k in ω). Note that the relation is also true for N = 1 if we make the natural choice $Z_0 = 1$.

We rewrite this result as a (formal) contour integral 3 :

$$Z_N(t_0,\cdots,t_j,\cdots) = \oint \frac{d\omega}{\omega} \sum_k t_k \omega^{-k} Z_{N-1}(t_0+\omega t_1,\cdots,t_j+\omega t_{j+1},\cdots).$$

The same argument, based on enumerating the ways the point N can be linked to the remaining part of the graph under the condition that $\hat{G}_N = k$, shows that

$$Z_N \pi_{N,k} = t_k \oint \frac{d\omega}{\omega} \omega^{-k} Z_{N-1}(t_0 + \omega t_1, \cdots, t_j + \omega t_{j+1}, \cdots),$$

to be compared with the formula of Section 2.3. These two formulæ for $\pi_{N,k}$ are not so trivially equivalent because they involve different rearrangements of the sum of weights.

2.6 Component distribution

We study the distribution of sizes of connected components.

Define W_n by

$$W_n = \sum_{G; V(G)=n}^c w_G,$$

where $\sum_{i=1}^{c}$ denotes the sum over connected graphs. Observe that if G splits as a disjoint union of two subgraphs G_1 and G_2 (G contains no edge joining a vertex of G_1 to a vertex of G_2), the weight of G factorizes : $w_G = w_{G_1} w_{G_2}$. So the total weight of graphs G of size N that are the disjoint union of k_1 connected components of size 1 (i.e. isolated points), k_2 connected components of size $2, \dots, k_n$ connected components of size n, \dots (so $\sum_{n \ge 1} nk_n = N$) is

$$\frac{N!}{\prod_n k_n! n!^{k_n}} \prod_n W_n^{k_n}.$$

The combinatorial factor just counts the number of ways to split the N vertices of G in packets of the right size. Summing over all possible k_n 's

³The symbol \oint denotes the contour integral $\frac{1}{2i\pi}\int$ along small contour surrounding the origin.

gives back Z_N :

$$Z_N = \sum_{k_n \ge 0, \sum_{n \ge 1} n k_n = N} \frac{N!}{\prod_n k_n! n!^{k_n}} \prod_n W_n^{k_n}.$$

This formula allows to view Z_N not as a function of the t_k 's but as a function of the W_n 's, and using this interpretation, we see that, denoting by $C_m(G)$ the number of connected components of size m in the graph G, the average number of components of size m in the random graph model is

$$\sum_{G} p_G C_m(G) = W_m \frac{\partial \log Z_N}{\partial W_m}.$$

So $mW_m \frac{\partial \log Z_N}{\partial W_m}$ is the average number of sites belonging to components of size m, and summing over m we should have $\sum_m mW_m \frac{\partial \log Z_N}{\partial W_m} = N$. This is simply the statement that Z_N is a homogeneous function of degree N in the W_n 's if W_n is assigned degree n.

This can be rephrased in compact form. Introduce a (complex or formal)⁴ variable z and define $Z \equiv \sum_{N\geq 0} \frac{z^N}{N!} Z_N$, the z-generating function for the Z_N 's. Replacing Z_n by its expression in terms of the W_n 's, we get the (well-known) fact that $Z = e^W$ where $W = \sum_{n\geq 1} \frac{z^n}{n!} W_n$. Conversely, one retrieves Z_N by

$$Z_N = N! \oint \frac{dz}{z} z^{-N} e^{\sum_{n \ge 1} \frac{z^n}{n!} W_n}.$$
 (2)

The average number of components of size n in the random graph is thus

$$W_n \frac{\partial \log Z_N}{\partial W_n} = \frac{N!}{n!(N-n)!} W_n \frac{Z_{N-n}}{Z_N}.$$

Similarly, the average number of times a given graph g of size $n \leq N$ appears as a connected component in the random graph G of size N is

$$\frac{N!}{n!(N-n)!} w_g \frac{Z_{N-n}}{Z_N}.$$

⁴In this section, some of the computations we make require that the t_k 's satisfy some properties so as to ensure that the series we write have a finite domain of convergence. For instance, we could assume that only a finite (though arbitrarily large) number of t_k 's are non vanishing. Alternatively we could work with formal power series.

2.7 Discussion

A crucial observation is that the weight of a graph depends only on its degree distribution, as in the Molloy–Reed model. But whereas in the Molloy–Reed model the weight of a graph is 0 unless it has the correct degree distribution, the degree distribution fluctuates in our model. So our model is a canonical description of a random graph model with given "number of edges distribution tion at a vertex", and the Molloy–Reed model a microcanonical one.

That the two models turn out to be equivalent in some large N limit is maybe not surprising. However, note that contrary to standard statistical mechanics (when only a few quantities, for instance energy and number of particles, fluctuate in the canonical description but are fixed in the microcanonical one) the constraint hypersurface of the microcanonical model has a codimension that gets larger and larger as N grows.

Finally, let us observe that the maximum entropy model is well suited for standard thermodynamical simulations, namely heatbath algorithms or metroplolis algorithms. This is because contrary to the 'naive' model or the Molloy–Reed model the phase space has a simple structure.

3 Finite connectivity limit

3.1 General analysis

As suggested at the end of section 2.3 by the special case of the Erdös–Renyi model, we shall show that a thermodynamic limit occurs in the large N limit if t_k scales like $N^{-k/2}$. Note that in this case w_G , the weight of G scales like $N^{-E(G)}$, where as before E(G) stands for the number of edges of G.

The starting point of the analysis will be the functional equation established in section 2.5:

$$Z_N(t_0,\cdots,t_j,\cdots) = \oint \frac{d\omega}{\omega} \sum_k t_k \omega^{-k} Z_{N-1}(t_0+\omega t_1,\cdots,t_j+\omega t_{j+1},\cdots).$$

We set $t_k = \tau_k N^{-k/2}$ and define $F_N(\tau_{\cdot}) \equiv \frac{1}{N} \log Z_N(t_{\cdot})$. Substituting $\omega N^{-1/2}$ for ω leads after a few manipulations to

$$e^{NF_{N}(\tau_{.})} = \oint \frac{d\omega}{\omega} \sum_{k} \tau_{k} \omega^{-k} e^{(N-1)F_{N-1}((\tau_{.} + \frac{\omega}{N}\tau_{.} + 1)(1-1/N)^{k/2})}.$$

This equation still involves no approximation. Now we make the usual thermodynamical hypothesis, namely that $NF_N(\tau) - (N-1)F_{N-1}(\tau)$ has a limit, say $F(\tau)$ when $N \to \infty$. This implies in particular that $F_N(\tau)$ converges to $F(\tau)$. The above equation has then a large N limit. To see it clearly, we rewrite it as

$$e^{NF_{N}(\tau_{\cdot}) - (N-1)F_{N-1}(\tau_{\cdot})} = \oint \frac{d\omega}{\omega} \sum_{k} \tau_{k} \omega^{-k} e^{(N-1)[F_{N-1}((\tau_{\cdot} + \frac{\omega}{N}\tau_{\cdot} + 1)(1-1/N)^{k/2}) - F_{N-1}(\tau_{\cdot})]}$$

In the large N limit, this leads to the equation

$$1 = \bar{y} \sum_{k} \tau_k \frac{\bar{x}^k}{k!},\tag{3}$$

where we have defined

$$\bar{y} \equiv e^{-F - \frac{1}{2} \sum_{k} k \tau_k \frac{\partial F}{\partial \tau_k}} , \quad \bar{x} \equiv \sum_{k} \tau_{k+1} \frac{\partial F}{\partial \tau_k}$$

On can take an analogous limit of other relations in 2.5 to obtain the more detailed equations for the degree distribution,

$$\pi_k = \tau_k \frac{\partial F}{\partial \tau_k} = \bar{y} \tau_k \frac{\bar{x}^k}{k!}.$$
(4)

Eq.(3) ensures that this distribution is correctly normalized, $\sum_k \pi_k = 1$.

The parameter \bar{x} possesses a simple interpretation. We start from the relation $\frac{\partial F}{\partial \tau_k} = \bar{y}\tau_k \frac{\bar{x}^k}{k!}$, multiply it by $\bar{x}\tau_{k+1}$ and sum over k to get

$$\bar{x}^2 = \bar{y} \sum_k \tau_{k+1} \frac{\bar{x}^{k+1}}{k!} = \sum_k k \pi_k,$$
 (5)

so that \bar{x}^2 is the first moment of the distribution π_k for the number of edges incident at a vertex.

We can summarize quite compactly our results as follows: Let us introduce the function $V(x) \equiv \sum_k \tau_k \frac{x^k}{k!}$, which we call the potential for reasons which will be clear in a moment. If all our previous formulae are to make sense, this function should have a positive radius of convergence. Let us also define

$$\mathcal{F}(y,x) \equiv -1 - \log y - \frac{x^2}{2} + yV(x). \tag{6}$$

Then (\bar{y}, \bar{x}) is a critical point for \mathcal{F} , thanks to eqs.(3,5), and F is the corresponding critical value.

It is not true that these equations for (\bar{y}, \bar{x}) always have a single solution. It is not difficult to find examples with no solution at all. We can interpret this by saying that in that case there is no thermodynamic limit in our sense. More troublesome is the case when there are several solutions. The most naive requirement would be that the physical solution is to take the couple (\bar{y}, \bar{x}) that leads to the absolute maximum F_{max} for F because the factor $e^{NF_{max}}$ will be the dominant contribution to Z. We shall meet such a behaviour in one of the examples of Section 3.6, and make a few comments there.

For most of the paper, we shall simply assume that if there is more than one extremum, we have picked the correct one.

3.2**Connected** components

In section 2.6, we gave a formula for Z_N in terms of connected components. This formula has also an interesting limiting form in the thermodynamic limit, but we shall wait until the next section to derive it. For the time being, recall that W_n is the sum of the weights of connected graphs on n vertices. We have shown that the average number of components of size n in the random graph is $\frac{N!}{n!(N-n)!}W_n \frac{Z_{N-n}}{Z_N}$. Now, we split $W_n = \sum_{l \ge 0} W_{n,l}$ as a sum of contributions corresponding to

connected graphs with $l = 0, 1, \cdots$ (independent) loops⁵. If G is a connected graph with L (independent) loops, E edges and V vertices, an old theorem of Euler says that L = E - V + 1 (in particular trees, i.e. connected graphs without loops, have E = V-1) so $W_{n,l}$ is simply the homogeneous component of degree 2(n+l-1) in the $t'_k s$. If we set $t_k = \tau_k N^{-k/2}$ we see that $W_{n,l}(t_k) =$ $N^{1-n-l}W_{n,l}(\tau)$. We define $T_n \equiv W_{n,0}(\tau)$.

When $N \to \infty$ for fixed *n* and τ_k 's we find that $\frac{N!}{n!(N-n)!}W_n \sim N\frac{T_n}{n!}$, meaning that trees dominate. In the thermodynamic limit, we find as before that $\frac{Z_{N-n}}{Z_N}(t_{-}) \sim e^{-n(F+\frac{1}{2}\sum_k k\tau_k \frac{\partial F}{\partial \tau_k})} \equiv \bar{y}^n$. So in the thermodynamic limit, the average number of components of size

n in the random graph is $N\frac{T_n}{n!}\bar{y}^n$. The number of points in components of

⁵Or closed circuits in the mathematical literature.

size n in the random graph is

$$C_n \equiv Nn \frac{T_n}{n!} \bar{y}^n,$$

and the total fraction of sites occupied by finite components is

$$Q \equiv \sum_{n} n \frac{T_n}{n!} \bar{y}^n.$$

If this number is 1, we can consistently interpret the random graph model as a random forest model in the thermodynamical limit. However, if this number is < 1, this means that a finite fraction of points is not in finite components, and there is a percolation cluster in the system.

3.3 Tree distribution

We would like to find a closed formula for the generating function

$$T(y) \equiv \sum_{n} n \frac{T_n}{n!} y^n.$$

The first observation comes from an analogy with a baby quantum field theory. The asymptotic expansion of the integral

$$I = \frac{1}{(2\pi\hbar)^{1/2}} \int_{-\infty}^{+\infty} e^{(-x^2/2 + yV(x))/\hbar}$$

in powers of the τ_k 's has a useful reinterpretation. Namely

$$I \sim \sum_{G} \frac{1}{A(G)} \hbar^{L(G) - C(G)} \prod_{k} (y\tau_k)^{\tilde{G}_k},$$

where the sum is over Feynman graphs⁶ with an arbitrary number of vertices, A(G) is the order of the automorphism group of G (for a precise definition see e.g. [3]) and C(G) the number of connected components of G. Again by a factorization argument for the weights the connected contributions exponentiate, and

$$\hbar \log I \sim \sum_{G}^{c} \frac{1}{A(G)} \hbar^{L(G)} \prod_{k} (y\tau_{k})^{\tilde{G}_{k}}.$$

⁶Warning : Feynman graphs are essentially general graphs i.e. not necessarily simple !

In the classical (small \hbar) limit, on the one hand graphs with L(G) = 0 dominate. Though Feynman graphs are not necessarily simple, loopless Feynman graphs are just ordinary trees. On the other hand, I can be calculated in the limit \hbar small by the saddle point approximation, leading to the identity between formal power series:

$$T(y) \equiv \sum_{\text{loopless}G}^{c} \frac{1}{A(G)} \prod_{k} (y\tau_k)^{\tilde{G}_k} = S(\tilde{x})$$
(7)

with $S(x) = -x^2/2 + yV(x) = -x^2/2 + y\sum_k \tau_k x^k/k!$ and \tilde{x} is the formal power series of y and the τ_k 's for which S is extremal, $\tilde{x} = y\sum_k \tau_{k+1}\tilde{x}^k/k! = yV'(\tilde{x})$.

Hence, the expansion of $-\tilde{x}^2/2 + yV(\tilde{x})$ in a formal power series of y with $\tilde{x} = yV'(\tilde{x})$ yields $\sum_n y^n T_n/n!$. From $T(y) = -\tilde{x}^2/2 + yV(\tilde{x})$ and the stationnarity condition, we also infer that $T'(y) = V(\tilde{x})$.

The expansion of T(y) is convergent for small y if V(x) has a non vanishing radius of convergence. Note that if $\tau_1 = 0$, the solution $\tilde{x} = 0$ has to be chosen, because it leads to the correct $T(y) = \tau_0 y$ (trees on two or more vertices have leaves, so they count 0 if $\tau_1 = 0$).

We now make the general assumption that $\tau_1 \neq 0$, and V(x) has a nonvanishing radius of convergence. Let us study the inversion of the relation y = x/V'(x). This can be obtained via the Lagrange formula. By Cauchy's residue formula

$$\tilde{x}(y) = \oint x \, \frac{(x/V'(x))'}{x/V'(x) - \tilde{x}/V'(\tilde{x})} dx$$

where the x contour has index 1 with respect to \tilde{x} . Replacing $\tilde{x}/V'(\tilde{x})$ by y, the y-expansion yields

$$\tilde{x}(y) = \sum_{n \ge 0} y^n \oint x \, \frac{(x/V'(x))'}{(x/V'(x))^{n+1}} dx.$$

One can use integration by parts to get:

$$\tilde{x}(y) = \sum_{n \ge 1} \frac{y^n}{n!} \left(\frac{d^{n-1}}{dx^{n-1}} \left(\frac{dV}{dx} \right)^n \right)_{|x=0}.$$

This is clearly a series with non-negative coefficients. As a consequence, its radius of convergence is given by the first singularity on the positive real axis, at the point $y_m = x_m/V'(x_m)$ corresponding to the unique maximal value of the concave function x/V'(x). This maximum might have two origins : either

 x_m is a singular point of V, or the derivative of x/V'(x), which is positive for small x, vanishes at x_m . This is equivalent to $V'(x_m) - x_m V''(x_m) = 0$

The explicit form of T_n can similarly be obtained via the Lagrange formula:

$$T(y) = \tau_0 y + \sum_{n \ge 1} \frac{y^n}{n!} \left(\frac{d^{n-2}}{dx^{n-2}} \left(\frac{dV}{dx} \right)^n \right)_{|x=0},$$

a classical formula which can also be proved by purely combinatorial arguments, giving an independent argument for the fact that the classical limit of quantum field theory is described by trees.

3.4 Percolation

The analysis of the previous section shows that the series

$$\sum_n n(x/V'(x))^{n-1} T_n/n!$$

converges for any positive x in the domain of convergence of V, and that, in this domain, its sum is equal to V(x) for $x < x_m$.

For $x > x_m$ the series is still convergent. However there is a (unique) number $x^* < x_m$ such that $x^*/V'(x^*) = x/V'(x)$ and, since the series only involves the ratio x/V'(x), we have:

$$\sum_{n} n(x/V'(x))^{n-1} T_n/n! = V(x^*) < V(x), \quad \text{for} \quad x > x_m.$$

The percolation question can be rephrased as follows. Is the relevant solution \bar{x} of the system $\bar{y}V'(\bar{x}) = \bar{x}$, $\bar{y}V(\bar{x}) = 1$ such that $\bar{x} \leq x_m$ or not? Indeed, we know that the fraction of points in finite clusters is $\sum_n n \frac{T_n}{n!} \bar{y}^n$. Substituting $\bar{y} = \bar{x}/V'(\bar{x})$, this series sums to $\bar{y}V(\bar{x}) = 1$ if $\bar{x} \leq x_m$ but to $\bar{y}V(x^*(\bar{x})) < 1$ if $\bar{x} > x_m$. The condition $\bar{x} \leq x_m$ is equivalent to the condition $V'(\bar{x}) - \bar{x}V''(\bar{x}) \leq 0$. This can be transcribed in term of the probability distribution $\{\pi_k\} : \bar{y}\bar{x}V'(\bar{x}) = \sum_k k\pi_k \equiv \langle k \rangle$ and $\bar{y}\bar{x}^2V''(\bar{x}) = \sum_k k(k-1)\pi_k \equiv$ $\langle k(k-1) \rangle$ (we use brackets to denote averages of the distribution $\{\pi_k\}$).

Thus, the percolation criterion is that there is a percolation cluster in the system if and only if $\langle 2k - k^2 \rangle < 0$. This is precisely the criterion given in ref.[1]. The relative size of the giant component $Q_{\infty} = 1 - \sum_{n} n \frac{T_n}{n!} \bar{y}^n$ is then:

$$Q_{\infty} = 1 - \bar{y}V(\bar{x}^*) = 1 - \sum_k \pi_k (\bar{x}^*/\bar{x})^k$$
(8)

where \bar{x}^* is the smallest x solution of $\bar{y} = x/V'(x)$. This is again in agreement with the result of ref.[1]. Close to the percolation threshold, and for a generic potential V, the size of the giant component increases linearly with $\langle k^2 - 2k \rangle$:

$$Q_{\infty} \simeq \frac{\langle 2k \rangle \langle k^2 - 2k \rangle}{\langle k(k-1)(k-2) \rangle}$$

This formula is not valid when the probability distribution π_k has no third moment. Then the grows of the giant component close to the transition can exhibit a different critical behavior. We shall give an example of this situation in the examples of section 3.6.

Let us analyse in more details what happens if $\bar{x} = x_m$. We know that there is no percolation cluster. Now, if the radius of convergence of V is strictly larger than x_m , close to $y_m = x_m/V'(x_m)$, T'(y) has a square root branch point. This implies that the contribution of points in components of size n in the system decreases algebraically as $C_n \sim Nn^{-3/2}$ for large n. In the physics language, this is interpreted as a critical point and 3/2 as a critical exponent. Note that even in this case, the distribution $\{\pi_k\}$ is still decreasing at least exponentially at large k.

To observe other critical points, with different critical exponents, the radius of convergence of V has to be exactly $\bar{x} = x_m$, which requires some fine tuning. In that case, both C_n and π_k decrease algebraically. Assume that V has a leading singularity at $\bar{x} = x_m$ locally of the form $(\bar{x} - x)^{\gamma}$, with $\gamma > 2$ to ensure the existence of $\langle k \rangle$ and $\langle k^2 \rangle$. Generically, $y - y_m$ is linear in $x - x_m$ so that yT'(y) = V(x) has a leading singularity of the form $(y - y_m)^{\gamma}$ and both π_k and $C_k/N = kT_k \bar{y}^k/k!$ decrease as $k^{-\gamma-1}$. We shall give an example below.

If there is no percolation cluster, we can treat the large N limit from another point of view. We start from eq.(2) and in the contour integral giving Z_N , we change variables and replace $z \to Nz$, leading to

$$Z_N = \frac{N!}{N^N} \oint \frac{dz}{z} z^{-N} e^{N \sum_{l \ge 0} N^{-l} \sum_{n \ge 1} \frac{z^n}{n!} W_{n,l}(\tau)}.$$

For fixed n, the connected graphs with loops $(l \ge 1)$ are suppressed by inverse powers of N. However, in the sum over the size of connected components, terms up to n = N make a contribution to the contour integral, and it might happen that for large n and N related by some condition connected components of size n of with loops make a finite contributions to $\frac{1}{N} \log Z_N$. However, if there is no percolation cluster, we may safely neglect $l \ge 1$ and get an accurate approximation to the leading exponential behavior of Z_N in the large N limit.

Under appropriate conditions, the contour integral for Z_N can be deformed to pass through a dominant saddle point. Then the free energy is given by the saddle point approximation. We see that $Z_N \sim e^{NF(\tau_i)}$ with $F(\tau_i) = -1 - \log \bar{z} + T(\bar{z})$, \bar{z} being the the saddle point maximizing $-1 - \log z + T(z)$. This equation is what one gets from eq.(6) when $\bar{y} = \bar{z}$ and \bar{x} is seen as a function of $\bar{y} = \bar{x}/V'(\bar{x})$. This gives yet another proof of the dominance of trees and the Lagrange inversion formula.

3.5 Conditional degree distributions

We now present formulas for the degree distributions, denoted $\pi_k^{(n)}$, for vertices within clusters of size n. We are particularly interested in the degree distribution $\pi_k^{(\infty)}$ in the giant component when it exists.

From the last formula of Section 2.6, the average number of vertices of degree k belonging to a component of size n is:

$$C_n(k) = \frac{N!}{n!(N-n)!} \left(t_k \frac{\partial W_n}{\partial t_k} \right) \frac{Z_{N-n}}{Z_N}$$

In the thermodynamic limit, $t_k = N^{-k/2} \tau_k$, $N \to \infty$, this becomes

$$\frac{C_n(k)}{N} = \frac{\bar{y}^n}{n!} (\tau_k \frac{\partial T_n}{\partial \tau_k}).$$

By definition, the degree distribution within components of size n is $C_n(k)$ divided by the average number of points in components of size n so that $\pi_k^{(n)} = C_n(k)/C_n$, with $C_n/N = nT_n\bar{y}^n/n!$ in the thermodynamic limit. Hence:

$$\pi_k^{(n)} = \frac{1}{n} \tau_k \frac{\partial \log T_n}{\partial \tau_k}.$$

Notice that these distributions are normalized, $\sum_k \pi_k^{(n)} = 1$, since the T_n 's are homogeneous polynomials in the τ_i of degree n if each τ_i is assigned degree one.

Assume now that the percolation criterion is satisfied so that a giant component exists. The number of vertices of degree k in the giant component

are: $C_{\infty}(k) = N\pi_k - \sum_n C_n(k)$. In the thermodynamic limit,

$$C_{\infty}(k)/N = \pi_k - \sum_n (\tau_k \partial_{\tau_k} T_n) \frac{\bar{y}^n}{n!} = \pi_k - (\tau_k \partial_{\tau_k} T)(\bar{y})$$

But $T(y) = -\tilde{x}/2 + yV(\tilde{x})$ with the extremum condition $\tilde{x} = yV'(\tilde{x})$ so that $(\partial_{\tau_k}T)(y) = y(\tilde{x})^k/k!$. Using $\pi_k = \bar{y}\tau_k \frac{\bar{x}^k}{k!}$, we get, with \bar{x}^* defined as in Section 3.4:

$$\frac{C_{\infty}(k)}{N} = \pi_k \left(1 - \left(\frac{\bar{x}^*}{\bar{x}}\right)^k \right)$$

or equivalently,

$$\pi_k^{(\infty)} = \frac{\pi_k}{Q_\infty} \left(1 - \left(\frac{\bar{x}^*}{\bar{x}}\right)^k \right) \tag{9}$$

since $\pi_k^{(\infty)}$ is the ratio between $C_{\infty}(k)$ and the number of points in the giant cluster, which is NQ_{∞} . As it should, $\pi_k^{(\infty)}$ is correctly normalized: $\sum_k \pi_k^{(\infty)} =$ 1, and vanishes at k = 0 (there is no isolated vertex in the giant component). There is a crossover value $k_c = \log(\bar{x}/\bar{x}^*)^{-1}$ above which $\pi_k^{(\infty)}$ is exponentially close to π_k/Q_{∞} . Close to the transition the ratio $\pi_k^{(\infty)}/\pi_k$ goes to $k/\langle k \rangle$. The formula for $\pi_k^{(\infty)}$ has a simple probabilistic interpretation : as it is the conditional probability that a vertex has k neighbors given that it is in the giant component, it can be written as the quotient of $\pi_{k,\infty}$, the probability to have k neighbors and be in the percolation cluster, by Q_{∞} . We read from eq.(9) that $\pi_{k,\infty} = \pi_k - \pi_k \left(\frac{\bar{x}^*}{\bar{x}}\right)^k$. Hence $\pi_k \left(\frac{\bar{x}^*}{\bar{x}}\right)^k$ is the probability for a vertex to have k neighbors and to be in a finite component. This suggests that when a new point is added to the graph, the probability that it connects to k other vertices none of them in the giant component is $\pi_k \left(\frac{\bar{x}^*}{\bar{x}}\right)^k$: for each new edge, the penalty for avoiding the giant component is $\frac{\bar{x}^*}{\bar{x}}$

3.6 Reconstruction, with examples

The maximal entropy graph distribution can be reconstructed form the data of the degree distribution π_k , $\sum_k \pi_k = 1$. We set $H(s) = \sum_k \pi_k s^k$.

Given π_k , \bar{x} is defined as the positive square root of $\langle k \rangle = \sum_k k \pi_k$, and $\bar{y}\tau_k$ as $\pi_k k!/\bar{x}^k$. This yields $\bar{y}V(x) = \sum_k \pi_k (x/\bar{x})^k = H(x/\bar{x})$. The coefficient \bar{y} appears then as a normalization factor which may be choosen at will, eg. we could set $\bar{y} = 1$.

The tree distribution $T(y) = \sum_n T_n y^n / n!$ is then reconstructed, as a formal series, from $T'(y) = V(\tilde{x})$ with $\tilde{x} = yV'(\tilde{x})$. It is clear that $T_n \bar{y}^n$ is independent of the choosen normalization for \bar{y} .

The fraction of site occupied by the finite size components is $Q = \bar{y}T'(\bar{y})$. By construction, \bar{x} is solution to $\bar{x}/V'(\bar{x}) = \bar{y}$. The giant component exists when there are two solutions to the above equation in the interval $[0, \bar{x}]$. We denote by \bar{x}^* the smallest of them. The fraction of site occupied by the giant component is $Q_{\infty} = 1 - \bar{y}V(\bar{x}^*)$. Equivalently, one can look for a solution $0 < s^* < 1$ to the equation H'(s) = sH'(1), and if there is one, $Q_{\infty} = 1 - H(s^*)$.

Let us illustrate this reconstruction on a few simple examples.

- 1. Poissonian degree distribution: $\pi_k = e^{-\alpha} \alpha^k / k!$. This is the Erdös-Renyi model. We have $\bar{x} = \alpha^{1/2}$, and $V(x) = \exp x\bar{x}$, choosing $\bar{y} = e^{-\alpha}$. The tree distribution is $T'(y) = \exp \tilde{x}$ with $\tilde{x}e^{-\tilde{x}} = \alpha y$. The giant component exists for $\alpha > 1$ when the equation $\tilde{x}e^{-\tilde{x}} = \alpha e^{-\alpha}$ admits two solutions α^* and α with $\alpha^* < 1 < \alpha$. Its relative size is $Q_{\infty} = 1 - e^{-\alpha}T'(e^{-\alpha}) = 1 - \alpha^*/\alpha$.
- 2. Geometric degree distribution: $\pi_k = (1-p)p^k$. Then $\bar{x}^2 = p/(1-p)$ and $\bar{y}V(x) = 1/(1+\bar{x}^2-x\bar{x})$. The extremum relation x = yV'(x) is a cubic equation: $\bar{y}x(1+\bar{x}^2-x\bar{x})^2 = y\bar{x}$. The percolation transition is at $\bar{x}^2 = 1/2$ (p = 1/3) and the relative size of the giant component is $Q_{\infty} = 1 - (\bar{x}^*/\bar{x})^{1/2}$ with $\bar{x}\bar{x}^* = \frac{1}{2}(\bar{x}^2 + 2 - \bar{x}\sqrt{4+\bar{x}^2})$.

This example confronts us with the ambiguity problem alluded to a long time ago.

Changing p into 1-p leads to replace \bar{x} by $1/\bar{x}$. This changes $\bar{y}V(x) = 1/(1 + \bar{x}^2 - x\bar{x})$ into itself up to an irrelevant multiplicative factor. To state things in a slightly different way, the extremum conditions yV'(x) = x and yV(x) = 1 have two solutions, and one leads to the geometric distribution with parameter p and the other one to the geometric distribution with parameter 1-p. Of course, the real result of the computation of Z_N will make a definite choice. The criterion of the maximum for F leads to choose $\inf(p, 1-p)$, and this is also consistent with continuity starting from p = 0, a random graph made of isolated points.

However, the formulas obtained before for, say, the size of the giant

component, coincide with the ones from ref.[1] even for p > 1/2. This situation requires clarification. Maybe this is the point when the canonical and the microcanonical approaches finally diverge and stop being equivalent.

3. An example of a scale free distribution: $H(s) = \sum \pi_k s^k \equiv \tau_0 + \tau_1 s + \tau_2 s^2/2 + \tau (1 - \beta s + \beta (\beta - 1) s^2/2 - (1 - s)^\beta)$, where $2 < \beta < 3$ and τ_0, τ_1, τ_2 and τ are nonnegative parameters subject to the condition $\tau_0 + \tau_1 s + \tau_2 s^2/2 + \tau (\beta - 1)(\beta - 2)/2 = 1$ to ensure that the π_k 's are correctly normalized. The π_k 's decrease like $\pi_k \sim \frac{-\tau}{\Gamma(-\beta)} k^{-\beta-1}$.

Then $\bar{x}^2 = H'(s = 1) = \tau_1 + \tau_2 + \tau \beta(\beta - 2)$ from which the potential $\bar{y}V(x)$ is recovered as usual.

There is a percolation cluster if and only if the equation H'(s) = sH'(1)has a solution $s^* < 1$. So, we look for the solutions of $(1-s)(\tau\beta - \tau_1) = \tau\beta(1-s)^{\beta-1}$. If $\langle k^2 - 2k \rangle = \tau\beta - \tau_1$ is negative, there is no percolation cluster, but if it is positive, $1 - s^* = (1 - \frac{\tau_1}{\tau\beta})^{1/(\beta-2)}$. The size of the giant component is $1 - H(s^*) \sim (1 - s^*)H'(1)$, and

$$Q_{\infty} \sim \langle k^2 - 2k \rangle^{1/(\beta - 2)}$$

close to the threshold. This is an example when the growth of the giant component close to the threshold is nonlinear as a function of $\langle k^2 - 2k \rangle$.

The number of points in components of size k is reconstructed from T'(y) = V(x) with x = yV'(x). Below the threshold, this leads to a singularity $T'_{sing} \sim (\bar{y} - y)^{\beta}$, which implies that $C_n \sim n^{-\beta-1}$. Above the threshold, the radius of convergence r of T is larger than \bar{y} , leading to C_n 's that decrease exponentially as $C_n \sim n^{-3/2}(\bar{y}/r)^n$.

4 The case of oriented graphs

It is not difficult to modify the previous arguments to deal with maximum entropy oriented graphs with given "in-out" degree distributions. We give the percolation criterion, omitting all details.

The first result is that for such models, each vertex with k outgoing and l incoming vertices contributes a fixed multiplicative factor, say $t_{k,l}$, to the

weight of a graph. The generalization of the recursion formula for Z_N reads

$$Z_N(t_{i,j}) = \oint \frac{d\omega_+}{\omega_+} \frac{d\omega_-}{\omega_-} \sum_{k,l} t_{k,l} \omega_+^{-k} \omega_-^{-l} Z_{N-1}(t_{i,j} + \omega_+ t_{i,j+1} + \omega_- t_{i+1,j}).$$

The large N finite connectivity limit is obtained by letting $N \to \infty$ while keeping $\tau_{k,l} = t_{k,l} N^{(k+l)/2}$ fixed. Defining

$$V(x_{+}, x_{-}) \equiv \sum_{k,l} \tau_{k,l} \frac{x_{+}^{k}}{k!} \frac{x_{-}^{l}}{l!},$$

a straightforward adaptation of the argument in section 3.1 leads to the fact that the free energy F is the value of

$$\mathcal{F}(y,x) \equiv -1 - \log y - x_+ x_- + y V(x_+,x_-).$$

at the point $(\bar{y}, \bar{x}_+, \bar{x}_-)$ where it is maximum :

$$\bar{x_{+}} = \bar{y}(\partial_{x_{-}}V)(\bar{x}_{+}, \bar{x}_{-}), \quad \bar{x_{-}} = \bar{y}(\partial_{x_{+}}V)(\bar{x}_{+}, \bar{x}_{-}), \quad 1 = \bar{y}V(\bar{x}_{+}\bar{x}_{-}).$$
(10)

The analysis of the first two equations is a bit more involved than the analysis of the single implicit equation for the oriented case. We can view the pair of equations $x_+ = y\partial_{x_-}V$ and $x_- = y\partial_{x_+}V$ in the following way. It defines a function y over the curve C in the positive quadrant of the (x_+, x_-) plane given by $x_-\partial_{x_-}V = x_+\partial_{x_+}V$. This curve is smooth as long as V is well-defined. For instance, one can take $x \equiv x_+\partial_{x_+}V = x_-\partial_{x_-}V$ as an analytic parameter on it. Then y is a smooth convex function of x, and all properties of the non-oriented case are true for y(x) : y is a good analytic parameter on C for small y, but there is a singularity if the convex function y(x) has a maximum. To be more explicit, taking differentials we see that

$$\begin{pmatrix} dx/x \\ dx/x \end{pmatrix} = \begin{pmatrix} 1 + x_{+}\partial_{x_{+}}^{2}V/\partial_{x_{+}}V & x_{-}\partial_{x_{+}}\partial_{x_{-}}V/\partial_{x_{+}}V \\ x_{+}\partial_{x_{-}}\partial_{x_{+}}V/\partial_{x_{-}}V & 1 + x_{-}\partial_{x_{-}}^{2}V/\partial_{x_{-}}V \end{pmatrix} \begin{pmatrix} dx_{+}/x_{+} \\ dx_{-}/x_{-} \end{pmatrix},$$

and

$$\begin{pmatrix} dy/y \\ dy/y \end{pmatrix} = \begin{pmatrix} 1 - x_{+}\partial_{x_{-}}\partial_{x_{+}}V/\partial_{x_{-}}V & -x_{-}\partial_{x_{-}}^{2}V/\partial_{x_{-}}V \\ -x_{+}\partial_{x_{+}}^{2}V/\partial_{x_{+}}V & 1 - x_{-}\partial_{x_{+}}\partial_{x_{-}}V/\partial_{x_{+}}V \end{pmatrix} \begin{pmatrix} dx_{+}/x_{+} \\ dx_{-}/x_{-} \end{pmatrix}.$$

A simple computation shows that the determinant of the 2 by 2 matrix in the first relation is always strictly positive, but that the determinant of the 2 by 2 matrix in the second relation is positive for small y but can change sign. This happens if y'(x) vanishes. So the discussion of the non-oriented case carries over word for word. It is consistent to write $\bar{y} = y(\bar{x}), \bar{x}_{\pm} = x_{\pm}(\bar{x})$ and $\bar{x}^*(\bar{x})$ for the smallest x such that $y(\bar{x}) = y(x)$. We set $\bar{x}_{\pm}^* \equiv x_{\pm}(\bar{x}^*)$. There is a percolation cluster if and only if the second determinant is < 0 at $(\bar{y}, \bar{x}_{\pm}, \bar{x}_{\pm})$. This is equivalent to

$$(\partial_{x_-}V - x_+\partial_{x_-}\partial_{x_+}V)(\partial_{x_+}V - x_-\partial_{x_+}\partial_{x_-}V) - x_+x_-\partial_{x_-}^2V\partial_{x_+}^2V$$

being < 0 at that point. Then the fraction of sites in the percolation cluster is

$$Q_{\infty} = 1 - \bar{y}V(\bar{x}_{+}^{*}, \bar{x}_{-}^{*}).$$

To get the percolation criterion, we just have to rephrase the vanishing of the determinant in terms of the probability distribution π_{k_+,k_-} that a vertex of the random graph has k_+ outgoing and k_- incoming vertices. The explicit formula is

$$\pi_{k_+,k_-} = \bar{y}\tau_{k_+,k_-} \frac{\bar{x}_+^{k_+} \bar{x}_-^{k_-}}{k_+! k_-!}.$$
(11)

By construction $\sum_{k,l} k \pi_{k,l} = \sum_{k,l} l \pi_{k,l} \equiv \langle k \rangle$ since any graph has the same number of outgoing and incoming edges. The parameters \bar{x}_+ and \bar{x}_- are constrained by the relation

$$\bar{x}_+\bar{x}_- = \langle k \rangle$$

The percolation criterion reads:

$$(\langle k \rangle - \langle k_+ k_- \rangle)^2 - \langle k_+^2 - k_+ \rangle \langle k_-^2 - k_- \rangle < 0.$$
 (12)

Given the distribution $\pi_{k,l}$, the potential V can be reconstructed via eq.(11). As in the oriented case, the parameter \bar{y} is an arbitrary normalization factor. The product of the parameters \bar{x}_{\pm} is determined by $\bar{x}_{\pm}\bar{x}_{-} = \langle k \rangle$. The ratio $\bar{x}_{\pm}/\bar{x}_{-}$ can be choosen at will since there is a natural invariance in eq.(10). Namely if \bar{x}_{\pm} solves the extremum condition for the potential $V(x_{\pm}, x_{\pm})$, so does $\hat{x}_{\pm} = \lambda^{\pm 1} \bar{x}_{\pm}$ for the potential $W(x_{\pm}, x_{\pm}) = V(\lambda x_{\pm}, \lambda^{-1} x_{\pm})$ and leaves $\pi_{k,l}$ invariant. Again this finds its origin in the fact that any graph has the same number of outgoing and incoming edges.

Acknowledgments: We thank the E. Schrödinger Institut in Vienna for hospitality during the completion of this work.

References

- M. Molloy and B. Reed, Random Struct. Algorithms 6, 161, (1995); and Comb. Proba. Comput. 7, 295, (1998).
- S. N. Dorogovtsev and J.F.F. Mendes, *Evolution of Networks*, Adv. Phys. 51, 1079, (2002).
- [3] C. Itzykson and J.-B. Zuber, *Quantum Field Theory*, McGraw-Hill,(1980).
- [4] Z. Burda, J. D. Correia and A. Krzywicki, Phys. Rev. E 64 (2001) 046118.
- [5] M. E. Newman, S. H. Strogatz and D. Watts, Phys. Rev. E 64 (2001) 026118.
- [6] R. Albert and A.-L. Barabási, Statistical mechanics of complex networks, Reviews of Modern Physics 74, 47 (2002).
- [7] P. Erdös and A. Rényi, On the evolution of random graphs, Publ. Math. Inst. Hungar. Acad. Sci. 5 (1960), 17-61.