

Spherically Symmetric Solutions of Nonlinear Schrödinger Equations

Roman Cherniha

Vienna, Preprint ESI 187 (1995)

January 23, 1995

Supported by Federal Ministry of Science and Research, Austria
Available via WWW.ESI.AC.AT

SPHERICALLY SYMMETRIC SOLUTIONS OF NONLINEAR SCHRODINGER EQUATIONS

CHERNIHA ROMAN

(Institute of Mathematics, Ukrainian Academy of Sciences,
Tereshchenkivs'ka str.3, Kyiv-4, Ukraine)

Introduction

At the present time there are no efficient analytic methods for solving nonlinear partial differential equations (PDEs). Construction particular exact solutions to these equations remains an important problem. Finding exact solutions that have physical interpretation is of fundamental importance. Soliton solutions seem to be just as needed, since they describe the movement of a traveling solitary wave.

In many papers (see, for example, [1–5]) there are constructed and investigated soliton-like solutions of the nonlinear Schrodinger equation (NSchE)

$$i\Psi_t + \Delta\Psi + \lambda|\Psi|^{2d}\Psi = 0, \quad (1)$$

where $\Psi_t = \partial\Psi/\partial t$, $\Delta = \partial^2/\partial(x_1)^2 + \dots + \partial^2/\partial(x_n)^2$, $\Psi = \Psi(t, \vec{x})$, $\vec{x} = (x_1, \dots, x_n)$, $\lambda = const$, $d \in \mathfrak{R}$. But almost all of these solutions are solutions to the one-dimensional NSchEs ($n=1$) or their simple generalization to the n -dimensional case. For example, the well-known soliton solution [1]

$$\Psi = (\lambda/2)^{1/2} \alpha_1 \frac{\exp[(-iv/2)(x_1 + (v/2 - 2(\alpha_1)^2/v)t)]}{\cosh(\alpha_1 x_1 + \alpha_1 vt)} \quad (2)$$

to the NSchE

$$i\Psi_t + \Psi_{x_1 x_1} + \lambda|\Psi|^2\Psi = 0 \quad (3)$$

can be generalized to the n -dimensional case by substituting $\alpha_1 x_1 \rightarrow \alpha_1 x_1 + \dots + \alpha_n x_n$, $\alpha_a, v \in \mathfrak{R}$, $a = 1, \dots, n$. In the case of many-dimensional NSchEs it is of great importance to find spherically symmetric solutions (SSSs). In the present paper (Section 2) we construct soliton-like SSSs to the NSchEs of the form

$$\Delta\Psi + \lambda h_1(|\vec{x}|^2)|\Psi|^{2d}\Psi + \alpha h_2(|\vec{x}|^2)\bar{\Psi} = 0, \quad (4)$$

where h_1 and h_2 are arbitrary differentiable functions, λ and α are constants, $d \in \mathfrak{R}$, $\Psi = \Psi(\vec{x})$. and $|\vec{x}|^2 = x_1^2 + \dots + x_n^2$.

Note that nonstationary NSchEs of the form

$$i\Psi_{1,t} + \Delta\Psi_1 + \lambda h_1(|\vec{x}|^2)|\Psi_1|^{2d}\Psi_1 = 0, \Psi_{1,t} = \partial\Psi_1/\partial t \quad (5)$$

are reduced, by the substitution

$$\Psi_1(t, \vec{x}) = \exp(-i\alpha t)\Psi(\vec{x}), \quad (6)$$

to Eq.(4) for $h_2 = 1$. Consequently, the solutions of the NSchEs (5) can be obtained from those of the stationary NSchE (4) by the substitution (6).

In Section 3 the ansatze are presented for obtaining periodic solutions and solutions with singularity to Eq.(4).

Soliton-like Solutions of the Nonlinear Schrodinger Equations (4)

Since we are constructing spherically symmetric solutions (SSSs), we will always have

$$\Psi(\vec{x}) = \phi(|\vec{x}|^2). \quad (7)$$

Substituting the ansatz (7) in Eq.(4), we obtain the following nonlinear ordinary differential equation (ODE):

$$4\omega\phi_{\omega\omega} + 2n\phi_{\omega} + \lambda h_1(\omega)|\phi|^{2d}\phi + \alpha h_2(\omega)\phi = 0, \omega = |\vec{x}|^2 \quad (8)$$

(indices denote derivatives with respect to ω). It is easily seen that the substitution

$$\phi = \rho(\omega)\exp(iW(\omega)), \quad (9)$$

where ρ and W are real functions, reduces the ODE (8) to the following nonlinear system:

$$W_{\omega} = c\omega^{-n/2}\rho^{-2}, c \in \Re, \quad (10a)$$

$$4\omega\rho_{\omega\omega} + 2n\rho_{\omega} + \lambda h_1(\omega)\rho^{2d+1} - 4c^2\omega^{1-n}\rho^{-3} + \alpha h_2(\omega)\rho = 0. \quad (10b)$$

In the special case where

$$d = -2, h_1(\omega) = \omega^{1-n}, \lambda = 4c^2, h_2(\omega) = 1,$$

Eq.(10b) reduces to the Bessel equation, and we obtain its general solution in the form [6]

$$\rho(\omega) = \omega^{1/2-n/4}Z_{\nu}((\alpha\omega)^{1/2}), \nu = n/2 - 1, \quad (11)$$

$$W(\omega) = c \int \omega^{-1}[Z_{\nu}((\alpha\omega)^{1/2})]^{-2}d\omega + c_0,$$

where Z_{ν} are cylindrical functions.

Thus, the formulas (7), (9), (11) give the family of the SSSs to the NSchE

$$\Delta\Psi + \lambda|\vec{x}|^{2-2n}|\Psi|^{-4}\Psi + \alpha\Psi = 0. \quad (12)$$

In the general case the system (10) cannot be integrated. For this reason we will construct particular exact solutions to this system. Consider the following generalization of soliton-like solutions:

$$\rho(\omega) = af(\omega)[\cosh(g(\omega))]^{-1/d}, \omega = |\vec{x}|^2, \quad (13)$$

where f and g are arbitrary twice differentiable real functions except for a finite number of points, and a is an arbitrary constant.

It is easily seen that the soliton-like solution (2) can be obtained from the formula (13) for $n=1$ and $d=1$, if we apply the Galilei transformations (for details see [3]).

After substituting the ansatz (13) in Eq.(10b) for $c=0$, we obtain the following relation

$$\begin{aligned} &4\omega\rho[f_{\omega\omega} - d^{-1}(2f_{\omega}g_{\omega} + fg_{\omega\omega})\tanh(g(\omega)) + d^{-1}f(g_{\omega})^2[(1 + d^{-1}) \\ &(\tanh(g(\omega)))^2 - 1]] + 2n\rho[f_{\omega} - d^{-1}fg_{\omega}\tanh(g(\omega))] + \\ &+\lambda a^{2d}h_1(\omega)f^{2d+1}\rho(\cosh(g(\omega)))^{-2} + \alpha h_2(\omega)f\rho = 0. \end{aligned} \quad (14)$$

Let the coefficient multiplying $\tanh(g(\omega))$ in the relation (14) be zero. Then we obtain the equation

$$4\omega(2f_\omega g_\omega + fg_{\omega\omega}) + 2nf g_\omega = 0,$$

whence it follows

$$g_\omega = c_2\omega^{-n/2}f^{-2} \quad (15)$$

If we introduce now the condition

$$4\omega f_{\omega\omega} + 2nf_\omega = 0, \quad (16)$$

then we obtain

$$f = c_1\omega^{1-n/2} + c_0, n \neq 2 \quad (17)$$

$$f = c_1 \ln \omega + c_0, n = 2.$$

So, for the functions f and g of the form (15) and (17), the relation (14) can be reduced to

$$4d^{-1}(g_\omega)^2[(1+d^{-1})(\tanh(g(\omega)))^2 - 1] + \quad (14^1)$$

$$+\lambda a^{2d}h_1(\omega)f^{2d}(\cosh(g(\omega)))^{-2} + \alpha h_2(\omega) = 0, f \neq 0.$$

It is evident that the relation (14¹) turns into identity only under the following conditions:

$$h_1(\omega) = 4(1+d^{-1})c^2(\lambda d)^{-1}a^{-2d}\omega^{1-n}f^{-4-2d}, d \neq -1,$$

$$h_2(\omega) = -4c^2(\alpha d^2)^{-1}\omega^{1-n}f^{-4}.$$

Consequently, we have proved the following theorem.

THEOREM 1. Let a NSchE be of the form (4). Then

$$\Psi(|\vec{x}|^2) = af(|\vec{x}|^2)[\cosh(g(|\vec{x}|^2))]^{-1/d} \quad (18)$$

is the soliton-like SSS of this equation, if

$$f = c_1|\vec{x}|^{2-n} + c_0, n \neq 2,$$

$$f = c_1 \ln |\vec{x}|^2 + c_0, n = 2,$$

$$g = c_2 \int \omega^{-n/2} f^{-2} d\omega, \omega = |\vec{x}|^2, \quad (19)$$

$$h_1 = (1+d)b^2\lambda^{-1}a^{-2d}|\vec{x}|^{2-2n}f^{-4-2d},$$

$$h_2 = -b^2\alpha^{-1}|\vec{x}|^{2-2n}f^{-4},$$

where $b^2 = 4(c_2)^2/d^2, \alpha\lambda \neq 0, d > 0, c_0, c_1, c_2 \in \mathfrak{R}$

Remark 1. In case $d < 0$ the formula (18) gives the solution of the NSchE (4), but it is not a soliton-like solution.

Consider some examples, which are corollaries of Theorem 1.

Example 1. The case $n = 1(\vec{x} = x_1)$. For $c_1 = 0$ (see (19)) we obtain the well-known soliton-like solution

$$\Psi = a[\cosh(d\alpha^{1/2}x_1 + c)]^{-1/d} \quad (20)$$

of the stationary NSchE

$$\Psi_{x_1 x_1} + \lambda |\Psi|^{2d} \Psi - \alpha \Psi = 0$$

(indices denote derivatives with respect to x_1). Here and subsequently, $a^{2d} = (1+d)\alpha/\lambda$, $\alpha > 0$, $\lambda > 0$, $c \in \mathfrak{R}$.

From the solution (20), using the formula (6) for $\alpha \rightarrow -\alpha$ and the Galilei transformations [3], it is easily to obtain the following soliton solution:

$$\Psi = a \frac{\exp[(-iv/2)(x_1 + (v/2 - 2\alpha/v)t)]}{[\cosh(d\alpha^{1/2}(x_1 + vt) + c)]^{1/d}}$$

to the nonstationary NSchE (5) for $h_1 = 1$.

If $c_0 = 0$ (see (19)), then we obtain the solution

$$\Psi = ax_1 [\cosh(d\alpha^{1/2}/x_1 + c)]^{-1/d} \quad (21)$$

to the NSchE

$$\Psi_{x_1 x_1} + \lambda |x_1|^{-4-2d} |\Psi|^{2d} \Psi - \alpha |x_1|^{-4} \Psi = 0.$$

Example 2. The case $n = 2$ ($\vec{x} = (x_1, x_2)$). For $c_1 = 0$ (see (19)) we obtain the soliton-like solution

$$\Psi = a [\cosh(d\alpha^{1/2} \ln |\vec{x}| + c)]^{-1/d} \quad (22)$$

of the stationary NSchE

$$\Delta_2 \Psi + (\lambda/|\vec{x}|^2) |\Psi|^{2d} \Psi - (\alpha/|\vec{x}|^2) \Psi = 0,$$

d being a positive constant.

Example 3. The case $n = 3$ ($\vec{x} = (x_1, x_2, x_3)$). For $c_1 = 0$ (see (19)) we find the soliton-like solution

$$\Psi = a [\cosh(d(\alpha/|\vec{x}|^2)^{1/2} + c)]^{-1/d} \quad (23)$$

of the stationary NSchE

$$\Delta_3 \Psi + (\lambda/|\vec{x}|^4) |\Psi|^{2d} \Psi - (\alpha/|\vec{x}|^4) \Psi = 0,$$

d being a positive constant.

In case $c_0 = 0$ we obtain the soliton-like SSS

$$\Psi = a |\vec{x}|^{-1} [\cosh(d(\alpha |\vec{x}|^2)^{1/2} + c)]^{-1/d} \quad (24)$$

of the stationary NSchE

$$\Delta_3 \Psi + \lambda |\vec{x}|^{2d} |\Psi|^{2d} \Psi - \alpha \Psi = 0,$$

d being a positive constant.

From the solution (24) we can obtain (see (6)) the solution

$$\Psi_1 = a |\vec{x}|^{-1} \exp(i\alpha t) [\cosh(d(\alpha |\vec{x}|^2)^{1/2} + c)]^{-1/d}. \quad (25)$$

to the nonstationary NSchE

$$i\Psi_{1,t} + \Delta_3 \Psi_1 + \lambda (|\vec{x}| |\Psi_1|)^{2d} \Psi_1 = 0. \quad (26)$$

Note that for the solutions (24)-(25) possessing discontinuity at the point $|\vec{x}|^2 = 0$ the integrals are well defined in the vicinity of $\vec{x} = 0$, since we work in the space \mathfrak{R}^3 .

Example 4. The case $n = 4(\vec{x} = (x_1, x_2, x_3, x_4))$. For $c_1 = 0$ (see (19)) we find the soliton-like solution

$$\Psi = a[\cosh(d\alpha^{1/2}/(2|\vec{x}|^2) + c)]^{-1/d} \quad (27)$$

of the NSchE

$$\Delta_4\Psi + (\lambda/|\vec{x}|^6)|\Psi|^{2d}\Psi - (\alpha/|\vec{x}|^6)\Psi = 0,$$

d being a positive constant.

In case $c_0 = 0$ we obtain the soliton-like SSS

$$\Psi = a|\vec{x}|^{-2}[\cosh(d\alpha^{1/2}|\vec{x}|^2/2 + c)]^{-1/d}$$

of the stationary NSchE

$$\Delta_4\Psi + \lambda|\vec{x}|^{2+4d}|\Psi|^{2d}\Psi - \alpha|\vec{x}|^2\Psi = 0 \quad (28)$$

d being a positive constant.

Observe that Eq.(28) is a nonlinear generalization of the harmonic oscillator equation

$$\Delta\Psi - \alpha|\vec{x}|^2\Psi = 0.$$

Remark 2. The $\vec{x} = 0$ is not a point of discontinuity to the solutions (21)-(23),(27) for $d \neq 0$. So, the integrals are well defined in the vicinity of $\vec{x} = 0$.

Theorem 1 admits generalization, if we reject the condition (16).

THEOREM 2. The soliton-like solution (18) is a solution of the NSchE of the form (4), if

$$g = c_2 \int \omega^{-n/2} f^{-2} d\omega, \omega = |\vec{x}|^2, \quad (29a)$$

$$h_1 = (1 + d)b^2\lambda^{-1}a^{-2d}|\vec{x}|^{2-2n}f^{-4-2d}, \quad (29b)$$

$$h_2 = -(\alpha f)^{-1}[b^2|\vec{x}|^{2-2n}f^{-3} + 4\omega f_{\omega\omega} + 2nf_{\omega}], \quad (29c)$$

where $f(|\vec{x}|^2)$ is an arbitrary twice differentiable real functions except for a finite number of points, $b^2 = 4(c_2)^2/d^2$, $\alpha\lambda \neq 0$, $d > 0$, $c_2 \in \mathfrak{R}$

The proof of Theorem 2 is based on the relation (14) and is analogous to that of Theorem 1.

It is evident that in the case

$$4\omega f_{\omega\omega} + 2nf_{\omega} + b^2\omega^{1-n}f^{-3} + \alpha_1 f = 0 \quad (30)$$

the function $h_2(\omega) = \alpha_1/\alpha = const$ (see (29c)). On the other hand, Eq.(30) coincides with Eq.(10b) (for $\lambda = 0, h_2 = 1$) up to the sign of the term $\omega^{1-n}f^{-3}$. Evidently, in the case $n = 1, 3, 5, \dots$ the substitution $f(\omega) = f^1(v), \omega = -v$ reduces Eq.(30) to the form

$$4v(f^1)_{vv} + 2n(f^1)_v - b^2v^{1-n}(f^1)^{-3} - \alpha_1 f^1 = 0. \quad (31)$$

(indices denote derivatives with respect to v). Eq.(31) coincides with Eq.(10b) (for $\lambda = 0, h_2 = 1$) to within the notation. Now it is easily seen that any complex solution $\phi(v)$ ($Im\phi \neq 0$) of the Bessel equation

$$4v\phi_{vv} + 2n\phi_v - \alpha_1\phi = 0 \quad (32)$$

generates the function $f^1(v) = Re\phi$. So, with the help of f^1 , we can find a soliton-like SSS of the NSchE (4) for $h_2 = \alpha_1/\alpha$.

Consequently, we have proved the following theorem.

THEOREM 3. The soliton-like solution (18) is a solution of the NSchE of the form

$$\Delta\Psi + \lambda h_1(|\vec{x}|^2)|\Psi|^{2d}\Psi + \alpha_1\Psi = 0,$$

if $f = Re(-\omega)^{1/2-n/4}Z_\nu((\alpha_1\omega)^{1/2})$, $Im(-\omega)^{1/2-n/4}Z_\nu((\alpha_1\omega)^{1/2}) \neq 0$, and the functions g and h are determined with the help of the formulas (29a), (29b). Here $Z_\nu, \nu = |n/2 - 1|$ are cylindrical functions and $n=1,3,5\dots$

Discussion

We have thus proved Theorems 1,2 and 3 that allow us to construct soliton-like SSSs of the form (18) to the NSchE (4). This is well illustrated by the examples 1-4. In particular cases, where we have solutions to the NSchE (4) for $h_2 = const$, exact solutions can be found to the nonstationary NSchE (5). The well-known soliton-like solutions of the (1+1)-dimensional Schrodinger equation with a power nonlinearity, for example Zakharov-Shabat's solution, can be easily obtained from the formula (18) by the Galilean transformations.

It should be observed that the ansatz (13) is a particular case of the ansatz for Lie's solutions (see, for example, [7-9]) and of the ansatz for non-Lie's solutions [9,10,11]. But here we determine first the 'outer' function $F = (cosh)^{-1/d}$ and then obtain a system of ODEs for the unknown functions f and g . According to the papers [7-10] we find first the functions f and g , whereupon we find the unknown function F .

As concerns the direct method [10,11], it does not provide a clear algorithm for obtaining the 'outer' function F and it does not connect this function with the Lie solutions.

Here, for obtaining the function F , we have used the form of the Lie solutions constructed in [3]. Moreover, using the other Lie solutions from the papers [3], we can find the new ones to the NSchE (4). For example we can obtain periodic SSSs of the form

$$\Psi(|\vec{x}|^2) = af(|\vec{x}|^2)[cos(g(|\vec{x}|^2))]^{-1/d}, \quad (33)$$

and solutions with a singularity for $d > 0$ of the form

$$\Psi(|\vec{x}|^2) = af(|\vec{x}|^2)[sinh(g(|\vec{x}|^2))]^{-1/d}. \quad (34)$$

For ansatze (33)-(34) we can obtain the theorems that are analogous to Theorem 1,2 and 3.

Thus, a wide class of the Lie solutions to nonlinear Schrodinger equation enables us to determine a structure of exact solutions to a generalization of this equation.

Appendix

The NSchE (26) for $d = 2$ has a wide Lie symmetry. In fact, using the Lie method [7-9] we can prove that in this case Eq. (26) is invariant under the projective transformations

$$t' = t/(1 - pt), \quad x' = x/(1 - pt), \quad p \in \mathfrak{R}, \quad (35a)$$

$$\Psi'_1 = \Psi_1(1 - pt)^{3/2} \exp \frac{ip|\vec{x}|^2}{4(1 - pt)}, \quad (35b)$$

the scale transformations

$$t' = m^2t, \quad x' = mx, \quad \Psi'_1 = m^{-3/2} \exp(iq)\Psi_1, \quad m, q \in \mathfrak{R},$$

the translations in variable t and rotations in the space \mathfrak{R}^3 .

Using the transformations (35) to the solutions (25) for $d = 2$, we obtain the following soliton-like SSS

$$\Psi_1 = a[|\vec{x}|^2(1 - pt)]^{-1/2} \exp \frac{i(4\alpha t - p|\vec{x}|^2)}{4(1 - pt)} \left[\cosh \left(\frac{2(\alpha|\vec{x}|^2)^{1/2}}{1 - pt} + c \right) \right]^{-1/2} \quad (36)$$

of the NSchE

$$i\Psi_{1,t} + \Delta_3\Psi_1 + \lambda(|\vec{x}|\Psi_1)^4\Psi_1 = 0.$$

Note that the solution (36) is not a solution of the form (6).

Acknowledgments

The author wishes to express his thanks to Professor W.Thirring for stimulating discussion and advice and the ESI, where most of this work was carried out, for hospitality. I also acknowledge financial support by the Austrian Academic Exchange Service and Ukrainian DKNT.

References

- [1] Zakharov V E and Shabat A B 1971 Zhurn.Eksper.Teor.Fiz.(JETP) -v.61-p. 118-134 (in Russian).
- [2] Ablowitz M J and Segur H J 1981 Solitons and the Inverse Scattering Transform. -SIAM, Philadelphia, PA.
- [3] Fushchych W I and Cherniha R M 1989 Ukr.Mat.Zhurn. (Ukrainian Math. J.) -v.41.- p.1349-1357; p.1687-1694 (Journal transl. in the USA).
- [4] Cherniha R M 1995 Ukr.Fiz.Zhurn. (Ukrainian Phys.J.) -v.40. -N 5.(in Ukrainian).
- [5] Lieb E H and Thirring W E The Eigenvalues of the Schrodinger Hamilton (in Press)
- [6] Kamke E. 1959 Differentialgleichungen Lösungsmethoden und Lösungen. -Leipzig.
- [7] Bluman A and Cole B 1974 Similarity Methods for Differential Equations.- Berlin: Springer.
- [8] Ovsyannikov L V 1978 The Group Analysis of Differential Equations.-Moscow: Nauka (in Russian).
- [9] Fushchych W I, Shtelen W M and Serov M I 1993 Symmetry Analysis and Exact Solutions of Equations of Nonlinear Mathematical Physics -Dordrecht: Kluwer Publishers (Russian version- Kyiv : Naukova Dumka, 1989).
- [10] Clarkson P A and Kruskal M D 1989 J. Math. Phys.- v.30- p.2201-2213.
- [11] Clarkson P.A. 1992 Nonlinearity. - v.5. - p.453-472.