Compactness of the $\overline{\partial}$ -Neumann Operator on Weighted (0,q)-Forms

Friedrich Haslinger

Vienna, Preprint ESI 2291 (2010)

December 21, 2010

Supported by the Austrian Federal Ministry of Education, Science and Culture Available online at $\rm http://www.esi.ac.at$

Compactness of the $\overline{\partial}$ - Neumann operator on weighted (0,q)- forms

Friedrich Haslinger

Abstract. As an application of a characterization of compactness of the $\overline{\partial}$ -Neumann operator we derive a sufficient condition for compactness of the $\overline{\partial}$ -Neumann operator on (0, q)-forms in weighted L^2 -spaces on \mathbb{C}^n .

Mathematics Subject Classification (2000). Primary 32W05; Secondary 32A36. 35J10.

Keywords. $\overline{\partial}$ -Neumann problem, Sobolev spaces, compactness.

1. Introduction

In this paper we continue the investigations of [12] and [11] concerning existence and compactness of the canonical solution operator to $\overline{\partial}$ on weighted L^2 -spaces over \mathbb{C}^n .

Let $\varphi:\mathbb{C}^n\longrightarrow\mathbb{R}^+$ be a plurisubharmonic \mathcal{C}^2 -weight function and define the space

$$L^2(\mathbb{C}^n,\varphi)=\{f:\mathbb{C}^n\longrightarrow\mathbb{C}\ :\ \int_{\mathbb{C}^n}|f|^2\,e^{-\varphi}\,d\lambda<\infty\},$$

where λ denotes the Lebesgue measure, the space $L^2_{(0,q)}(\mathbb{C}^n, \varphi)$ of (0,q)-forms with coefficients in $L^2(\mathbb{C}^n, \varphi)$, for $1 \leq q \leq n$. Let

$$(f,g)_{\varphi} = \int_{\mathbb{C}^n} f \,\overline{g} e^{-\varphi} \, d\lambda$$

denote the inner product and

$$||f||_{\varphi}^2 = \int_{\mathbb{C}^n} |f|^2 e^{-\varphi} \, d\lambda$$

the norm in $L^2(\mathbb{C}^n, \varphi)$.

F. Haslinger

We consider the weighted $\overline{\partial}$ -complex

$$L^{2}_{(0,q-1)}(\mathbb{C}^{n},\varphi) \xrightarrow[\overline{\partial}]{\overline{\partial}_{\varphi}} L^{2}_{(0,q)}(\mathbb{C}^{n},\varphi) \xrightarrow[\overline{\partial}]{\overline{\partial}_{\varphi}} L^{2}_{(0,q+1)}(\mathbb{C}^{n},\varphi),$$

where for (0,q)-forms $u = \sum_{|J|=q}' u_J d\overline{z}_J$ with coefficients in $\mathcal{C}_0^{\infty}(\mathbb{C}^n)$ we have

$$\overline{\partial}u = \sum_{|J|=q}' \sum_{j=1}^n \frac{\partial u_J}{\partial \overline{z}_j} \, d\overline{z}_j \wedge d\overline{z}_J,$$

and

$$\overline{\partial}_{\varphi}^* u = -\sum_{|K|=q-1}' \sum_{k=1}^n \delta_k u_{kK} \, d\overline{z}_K,$$

where $\delta_k = \frac{\partial}{\partial z_k} - \frac{\partial \varphi}{\partial z_k}$.

There is an interesting connection between $\overline{\partial}$ and the theory of Schrödinger operators with magnetic fields, see for example [5], [2], [8] and [6] for recent contributions exploiting this point of view.

The complex Laplacian on (0, q)-forms is defined as

$$\Box_{\varphi} := \overline{\partial} \,\overline{\partial}_{\varphi}^* + \overline{\partial}_{\varphi}^* \overline{\partial},$$

where the symbol \Box_{φ} is to be understood as the maximal closure of the operator initially defined on forms with coefficients in \mathcal{C}_0^{∞} , i.e., the space of smooth functions with compact support.

 \Box_{φ} is a selfadjoint and positive operator, which means that

$$(\Box_{\varphi}f, f)_{\varphi} \ge 0$$
, for $f \in dom(\Box_{\varphi})$.

The associated Dirichlet form is denoted by

$$Q_{\varphi}(f,g) = (\overline{\partial}f, \overline{\partial}g)_{\varphi} + (\overline{\partial}_{\varphi}^* f, \overline{\partial}_{\varphi}^* g)_{\varphi}, \qquad (1.1)$$

for $f, g \in dom(\overline{\partial}) \cap dom(\overline{\partial}_{\varphi}^*)$. The weighted $\overline{\partial}$ -Neumann operator $N_{\varphi,q}$ is - if it exists - the bounded inverse of \Box_{φ} .

We indicate that a (0,1)-form $f = \sum_{j=1}^n f_j \, d\overline{z}_j$ belongs to $dom(\overline{\partial}_{\varphi}^*)$ if and only if

$$\sum_{j=1}^{n} \left(\frac{\partial f_j}{\partial z_j} - \frac{\partial \varphi}{\partial z_j} f_j \right) \in L^2(\mathbb{C}^n, \varphi)$$

and that forms with coefficients in $\mathcal{C}_0^{\infty}(\mathbb{C}^n)$ are dense in $dom(\overline{\partial}) \cap dom(\overline{\partial}_{\varphi}^*)$ in the graph norm $f \mapsto (\|\overline{\partial}f\|_{\varphi}^2 + \|\overline{\partial}_{\varphi}^*f\|_{\varphi}^2)^{\frac{1}{2}}$ (see [10]).

We consider the Levi - matrix

$$M_{\varphi} = \left(\frac{\partial^2 \varphi}{\partial z_j \partial \overline{z}_k}\right)_{jk}$$

of φ and suppose that the sum s_q of any q (equivalently: the smallest q) eigenvalues of M_φ satisfies

$$\liminf_{|z| \to \infty} s_q(z) > 0. \tag{1.2}$$

We show that (2.2) implies that there exists a continuous linear operator

$$N_{\varphi,q}: L^2_{(0,q)}(\mathbb{C}^n, \varphi) \longrightarrow L^2_{(0,q)}(\mathbb{C}^n, \varphi),$$

such that $\Box_{\varphi} \circ N_{\varphi,q} u = u$, for any $u \in L^2_{(0,q)}(\mathbb{C}^n, \varphi)$.

If we suppose that that the sum s_q of any q (equivalently: the smallest q) eigenvalues of M_{φ} satisfies

$$\lim_{|z| \to \infty} s_q(z) = \infty. \tag{1.3}$$

Then the $\overline{\partial}$ -Neumann operator $N_{\varphi,q}: L^2_{(0,q)}(\mathbb{C}^n, \varphi) \longrightarrow L^2_{(0,q)}(\mathbb{C}^n, \varphi)$ is compact. This generalizes results from [12] and [11], where the case of q = 1 was

This generalizes results from [12] and [11], where the case of q = 1 was handled.

Finally we discuss some examples in \mathbb{C}^2 .

2. The weighted Kohn-Morrey formula

First we compute

$$(\Box_{\varphi}u, u)_{\varphi} = \|\overline{\partial}u\|_{\varphi}^{2} + \|\overline{\partial}_{\varphi}^{*}u\|_{\varphi}^{2}$$

for $u \in dom(\Box_{\varphi})$. We obtain

$$\begin{split} \|\overline{\partial}u\|_{\varphi}^{2} + \|\overline{\partial}_{\varphi}^{*}u\|_{\varphi}^{2} &= \sum_{|J|=|M|=q}' \sum_{j,k=1}^{n} \epsilon_{jJ}^{kM} \int_{\mathbb{C}^{n}} \frac{\partial u_{J}}{\partial \overline{z}_{j}} \overline{\frac{\partial u_{M}}{\partial \overline{z}_{k}}} e^{-\varphi} d\lambda \\ &+ \sum_{|K|=q-1}' \sum_{j,k=1}^{n} \int_{\mathbb{C}^{n}} \delta_{j} u_{jK} \overline{\delta_{k} u_{kK}} e^{-\varphi} d\lambda, \end{split}$$

where $\epsilon_{jJ}^{kM} = 0$ if $j \in J$ or $k \in M$ or if $k \cup M \neq j \cup J$, and equals the sign of the permutation $\binom{kM}{jJ}$ otherwise. The right-hand side of the last formula can be rewritten as

$$\sum_{|J|=q}' \sum_{j=1}^n \left\| \frac{\partial u_J}{\partial \overline{z}_j} \right\|_{\varphi}^2 + \sum_{|K|=q-1}' \sum_{j,k=1}^n \int_{\mathbb{C}^n} \left(\delta_j u_{jK} \overline{\delta_k u_{kK}} - \frac{\partial u_{jK}}{\partial \overline{z}_k} \overline{\frac{\partial u_{kK}}{\partial \overline{z}_j}} \right) e^{-\varphi} d\lambda,$$

see [18] Proposition 2.4 for the details. Now we mention that for $f, g \in \mathcal{C}_0^{\infty}(\mathbb{C}^n)$ we have

$$\left(\frac{\partial f}{\partial \overline{z}_k}, g\right)_{\varphi} = -(f, \delta_k g)_{\varphi}$$

and hence

$$\left(\left[\delta_j, \frac{\partial}{\partial \overline{z}_k}\right] u_{jK}, u_{kK}\right)_{\varphi} = -\left(\frac{\partial u_{jK}}{\partial \overline{z}_k}, \frac{\partial u_{kK}}{\partial \overline{z}_j}\right)_{\varphi} + (\delta_j u_{jK}, \delta_k u_{kK})_{\varphi}$$

Since

$$\left[\delta_j, \frac{\partial}{\partial \overline{z}_k}\right] = \frac{\partial^2 \varphi}{\partial z_j \partial \overline{z}_k}$$

we get

$$\|\overline{\partial}u\|_{\varphi}^{2} + \|\overline{\partial}_{\varphi}^{*}u\|_{\varphi}^{2} = \sum_{|J|=q}' \sum_{j=1}^{n} \left\|\frac{\partial u_{J}}{\partial\overline{z}_{j}}\right\|_{\varphi}^{2} + \sum_{|K|=q-1}' \sum_{j,k=1}^{n} \int_{\mathbb{C}^{n}} \frac{\partial^{2}\varphi}{\partial z_{j}\partial\overline{z}_{k}} u_{jK}\overline{u}_{kK} e^{-\varphi} d\lambda.$$

$$(2.1)$$

Formula (2.1) is a version of the Kohn -Morrey formula, compare [18] or [16].

Proposition 2.1. Let $1 \le q \le n$ and suppose that the sum s_q of any q (equivalently: the smallest q) eigenvalues of M_{φ} satisfies

$$\liminf_{|z| \to \infty} s_q(z) > 0. \tag{2.2}$$

Then there exists a uniquely determined bounded linear operator

$$N_{\varphi,q}: L^2_{(0,q)}(\mathbb{C}^n, \varphi) \longrightarrow L^2_{(0,q)}(\mathbb{C}^n, \varphi),$$

such that $\Box_{\varphi} \circ N_{\varphi,q} u = u$, for any $u \in L^2_{(0,q)}(\mathbb{C}^n, \varphi)$.

Proof. Let $\mu_{\varphi,1} \leq \mu_{\varphi,2} \leq \cdots \leq \mu_{\varphi,n}$ denote the eigenvalues of M_{φ} and suppose that M_{φ} is diagonalized. Then, in a suitable basis,

$$\sum_{|K|=q-1}' \sum_{j,k=1}' \frac{\partial^2 \varphi}{\partial z_j \partial \overline{z}_k} u_{jK} \overline{u}_{kK} = \sum_{|K|=q-1}' \sum_{j=1}'' \mu_{\varphi,j} |u_{jK}|^2$$
$$= \sum_{J=(j_1,\dots,j_q)}' (\mu_{\varphi,j_1} + \dots + \mu_{\varphi,j_q}) |u_J|^2$$
$$\ge s_q |u|^2$$

It follows from (2.1) that there exists a constant C > 0 such that

$$u\|_{\varphi}^{2} \leq C(\|\overline{\partial}u\|_{\varphi}^{2} + \|\overline{\partial}_{\varphi}^{*}u\|_{\varphi}^{2})$$

$$(2.3)$$

for each (0,q)-form $u \in \operatorname{dom}(\overline{\partial}) \cap \operatorname{dom}(\overline{\partial}_{\varphi}^*)$. For a given $v \in L^2_{(0,q)}(\mathbb{C}^n,\varphi)$ consider the linear functional L on $\operatorname{dom}(\overline{\partial}) \cap \operatorname{dom}(\overline{\partial}_{\varphi}^*)$ given by $L(u) = (u,v)_{\varphi}$. Notice that $\operatorname{dom}(\overline{\partial}) \cap \operatorname{dom}(\overline{\partial}_{\varphi}^*)$ is a Hilbertspace in the inner product Q_{φ} . Since we have by 2.3

$$|L(u)| = |(u, v)_{\varphi}| \le ||u||_{\varphi} ||v||_{\varphi} \le CQ_{\varphi}(u, u)^{1/2} ||v||_{\varphi}$$

Hence by the Riesz reprentation theorem there exists a uniquely determined (0, q)-form $N_{\varphi,q}v$ such that

$$(u,v)_{\varphi} = Q_{\varphi}(u,N_{\varphi,q}v) = (\overline{\partial}u,\overline{\partial}N_{\varphi,q}v)_{\varphi} + (\overline{\partial}_{\varphi}^{*}u,\overline{\partial}_{\varphi}^{*}N_{\varphi,q}v)_{\varphi},$$

from which we immediately get that $\Box_{\varphi} \circ N_{\varphi,q}v = v$, for any $v \in L^2_{(0,q)}(\mathbb{C}^n, \varphi)$. If we set $u = N_{\varphi,q}v$ we get again from 2.3

$$\|\overline{\partial}N_{\varphi,q}v\|_{\varphi}^{2} + \|\overline{\partial}_{\varphi}^{*}N_{\varphi,q}v\|_{\varphi}^{2} = Q_{\varphi}(N_{\varphi,q}v, N_{\varphi,q}v) = (N_{\varphi,q}v, v)_{\varphi} \le \|N_{\varphi,q}v\|_{\varphi} \|v\|_{\varphi}$$

4

Compactness of the $\overline{\partial}$ - Neumann operator

$$\leq C_1(\|\overline{\partial}N_{\varphi,q}v\|_{\varphi}^2 + \|\overline{\partial}_{\varphi}^*N_{\varphi,q}v\|_{\varphi}^2)^{1/2} \|v\|_{\varphi}$$

hence

$$(\|\overline{\partial}N_{\varphi,q}v\|_{\varphi}^{2} + \|\overline{\partial}_{\varphi}^{*}N_{\varphi,q}v\|_{\varphi}^{2})^{1/2} \le C_{2}\|v\|_{\varphi}$$

and finally again by 2.3

$$\|N_{\varphi,q}v\|_{\varphi} \le C_3(\|\overline{\partial}N_{\varphi,q}v\|_{\varphi}^2 + \|\overline{\partial}_{\varphi}^*N_{\varphi,q}v\|_{\varphi}^2)^{1/2} \le C_4\|v\|_{\varphi}$$

where $C_1, C_2, C_3, C_4 > 0$ are constants. Hence we get that $N_{\varphi,q}$ is a continuous linear operator from $L^2_{(0,q)}(\mathbb{C}^n, \varphi)$ into itself (see also [13] or [4]).

3. Compactness of $N_{\varphi,q}$

We use a characterization of precompact subsets of L^2 -spaces, see [1]:

A bounded subset \mathcal{A} of $L^2(\Omega)$ is precompact in $L^2(\Omega)$ if and only if for every $\epsilon > 0$ there exists a number $\delta > 0$ and a subset $\omega \subset \subset \Omega$ such that for every $u \in \mathcal{A}$ and $h \in \mathbb{R}^n$ with $|h| < \delta$ both of the following inequalities hold:

(i)
$$\int_{\Omega} |\tilde{u}(x+h) - \tilde{u}(x)|^2 dx < \epsilon^2$$
, (ii) $\int_{\Omega \setminus \overline{\omega}} |u(x)|^2 dx < \epsilon^2$. (3.1)

In addition we define an appropriate Sobolev space and prove compactness of the corresponding embedding, for related settings see [3], [14], [15].

Definition 3.1. Let

$$\mathcal{W}_q^{Q_{\varphi}} = \{ u \in L^2_{(0,q)}(\mathbb{C}^n, \varphi) : \|\overline{\partial}u\|_{\varphi}^2 + \|\overline{\partial}_{\varphi}^*u\|_{\varphi}^2 < \infty \}$$

with norm

$$\|u\|_{Q_{\varphi}} = (\|\overline{\partial}u\|_{\varphi}^2 + \|\overline{\partial}_{\varphi}^*u\|_{\varphi}^2)^{1/2}.$$

Remark 3.2. $\mathcal{W}_q^{Q_{\varphi}}$ coincides with the form domain $dom(\overline{\partial}) \cap dom(\overline{\partial}_{\varphi}^*)$ of Q_{φ} (see [9], [10]).

Proposition 3.3. Let φ be a plurisubharmonic C^2 - weight function. Let $1 \leq q \leq n$ and suppose that the sum s_q of any q (equivalently: the smallest q) eigenvalues of M_{φ} satisfies

$$\lim_{|z| \to \infty} s_q(z) = \infty. \tag{3.2}$$

Then $N_{\varphi,q}: L^2_{(0,q)}(\mathbb{C}^n, \varphi) \longrightarrow L^2_{(0,q)}(\mathbb{C}^n, \varphi)$ is compact.

Proof. For (0,q) forms one has by (2.1) and Proposition 2.1 that

$$\|\overline{\partial}u\|_{\varphi}^{2} + \|\overline{\partial}_{\varphi}^{*}u\|_{\varphi}^{2} \ge \int_{\mathbb{C}^{n}} s_{q}(z) |u(z)|^{2} e^{-\varphi(z)} d\lambda(z).$$

$$(3.3)$$

We indicate that the embedding

$$j_{\varphi,q}: \mathcal{W}_q^{Q_{\varphi}} \hookrightarrow L^2_{(0,q)}(\mathbb{C}^n, \varphi)$$

F. Haslinger

is compact by showing that the unit ball of $\mathcal{W}_q^{Q_{\varphi}}$ is a precompact subset of $L^2_{(0,q)}(\mathbb{C}^n, \varphi)$, which follows by the above mentioned characterization of precompact subsets in L^2 - spaces with the help of Gårding's inequality to verify (3.1) (i)(see for instance [7] or [4]) and to verify (3.1) (ii) : we have

$$\int_{\mathbb{C}^n \setminus \mathbb{B}_R} |u(z)|^2 e^{-\varphi(z)} \, d\lambda(z) \le \int_{\mathbb{C}^n \setminus \mathbb{B}_R} \frac{s_q(z)|u(z)|^2}{\inf\{s_q(z) : |z| \ge R\}} e^{-\varphi(z)} d\lambda(z),$$

which implies by (3.3) that

$$\int_{\mathbb{C}^n \setminus \mathbb{B}_R} |u(z)|^2 e^{-\varphi(z)} \, d\lambda(z) \le \frac{\|u\|_{Q_{\varphi}}^2}{\inf\{s_q(z) \, : \, |z| \ge R\}} < \epsilon,$$

if R is big enough, see [11] for the details.

This together with the fact that $N_{\varphi,q} = j_{\varphi,q} \circ j_{\varphi,q}^*$, (see [18]) gives the desired result.

Remark 3.4. If q = 1 condition (3.2) means that the lowest eigenvalue $\mu_{\varphi,1}$ of M_{φ} satisfies

$$\lim_{|z| \to \infty} \mu_{\varphi,1}(z) = \infty.$$
(3.4)

This implies compactness of $N_{\varphi,1}$ (see [11]).

Examples: a) We consider the plurisubharmonic weight function $\varphi(z, w) = |z|^2 |w|^2 + |w|^4$ on \mathbb{C}^2 . The Levi matrix of φ has the form

$$\left(\begin{array}{cc} |w|^2 & \overline{z}w\\ \overline{w}z & |z|^2 + 4|w|^2 \end{array}\right)$$

and the eigenvalues are

$$\mu_{\varphi,1}(z,w) = \frac{1}{2} \left(5|w|^2 + |z|^2 - \sqrt{9|w|^4 + 10|z|^2|w|^2 + |z|^4} \right)$$
$$= \frac{16|w|^4}{2 \left(5|w|^2 + |z|^2 + \sqrt{9|w|^4 + 10|z|^2|w|^2 + |z|^4} \right)},$$

and

$$\mu_{\varphi,2}(z,w) = \frac{1}{2} \left(5|w|^2 + |z|^2 + \sqrt{9|w|^4 + 10|z|^2|w|^2 + |z|^4} \right).$$
that (3.4) fails since even

It follows that (3.4) fails, since even

$$\lim_{|z|\to\infty} |z|^2 \mu_{\varphi,1}(z,0) = 0,$$

but

$$s_2(z,w) = \frac{1}{4}\Delta\varphi(z,w) = |z|^2 + 5|w|^2,$$

hence (3.2) is satisfied for q = 2.

b) In the next example we consider decoupled weights. Let $n \ge 2$ and

$$\varphi(z_1, z_2, \dots, z_n) = \varphi(z_1) + \varphi(z_2) + \dots + \varphi(z_n)$$

be a plurisubharmonic decoupled weight function and suppose that $|z_{\ell}|^2 \Delta \varphi_{\ell}(z_{\ell}) \rightarrow +\infty$, as $|z_{\ell}| \rightarrow \infty$ for some $\ell \in \{1, \ldots, n\}$. Then the $\overline{\partial}$ -Neumann operator $N_{\varphi,1}$ acting on $L^2_{(0,1)}(\mathbb{C}^n, \varphi)$ fails to be compact (see [12], [9], [17]).

Finally we discuss two examples in \mathbb{C}^2 : for $\varphi(z_1, z_2) = |z_1|^2 + |z_2|^2$ all eigenvalues of the Levi matrix are 1 and $N_{\varphi,1}$ fails to be compact by the above result on decoupled weights, for the weightfunction $\varphi(z_1, z_2) = |z_1|^4 + |z_2|^4$ the eigenvalues are $4|z_1|^2$ and $4|z_2|^2$ and $N_{\varphi,1}$ fails to be compact again by the above result, whereas $N_{\varphi,2}$ is compact by 3.3.

References

- R.A. Adams and J.J.F. Fournier, *Sobolev spaces*. Pure and Applied Math. Vol. 140, Academic Press, 2006.
- [2] B. Berndtsson, $\overline{\partial}$ and Schrödinger operators. Math. Z. 221 (1996), 401–413.
- [3] P. Bolley, M. Dauge and B. Helffer, Conditions suffisantes pour l'injection compacte d'espace de Sobolev à poids. Séminaire équation aux dérivées partielles (France), Université de Nantes 1 (1989), 1–14.
- [4] So-Chin Chen and Mei-Chi Shaw, Partial differential equations in several complex variables. Studies in Advanced Mathematics, Vol. 19, Amer. Math. Soc., 2001.
- [5] M. Christ, On the ∂ equation in weighted L² norms in C¹. J. of Geometric Analysis 1 (1991), 193–230.
- [6] M. Christ and S. Fu, Compactness in the ∂-Neumann problem, magnetic Schrödinger operators, and the Aharonov-Bohm effect. Adv. Math. 197 (2005), 1–40.
- [7] G.B. Folland, Introduction to partial differential equations. Princeton University Press, Princeton, 1995.
- [8] S. Fu and E.J. Straube, Semi-classical analysis of Schrödinger operators and compactness in the of Neumann problem. J. Math. Anal. Appl. 271 (2002), 267–282.
- [9] K. Gansberger, Compactness of the ∂-Neumann operator. Dissertation, University of Vienna, 2009.
- [10] K. Gansberger and F. Haslinger, Compactness estimates for the ∂̄- Neumann problem in weighted L² - spaces. Complex Analysis (P. Ebenfelt, N. Hungerbühler, J.J. Kohn, N. Mok, E.J. Straube, eds.), Trends in Mathematics, Birkhäuser (2010), 159–174.
- [11] F. Haslinger, Compactness for the ∂̄- Neumann problem- a functional analysis approach. Collectanea Math. (to appear), arXiv: 0912.4406.
- [12] F. Haslinger and B. Helffer, Compactness of the solution operator to $\overline{\partial}$ in weighted L^2 spaces. J. of Functional Analysis **255** (2008), 13–24.
- [13] L. Hörmander, An introduction to complex analysis in several variables. North Holland, 1990.
- [14] J. Johnsen, On the spectral properties of Witten Laplacians, their range projections and Brascamp-Lieb's inequality. Integral Equations Operator Theory 36 (2000), 288– 324.

F. Haslinger

- [15] J.-M. Kneib and F. Mignot, Equation de Schmoluchowski généralisée. Ann. Math. Pura Appl. (IV) 167 (1994), 257–298.
- [16] J.D. McNeal, L² estimates on twisted Cauchy-Riemann complexes. 150 years of mathematics at Washington University in St. Louis. Sesquicentennial of mathematics at Washington University, St. Louis, MO, USA, October 3–5, 2003. Providence, RI: American Mathematical Society (AMS). Contemporary Mathematics **395** (2006), 83–103.
- [17] G. Schneider, Non-compactness of the solution operator to ∂ on the Fock-space in several dimensions. Math. Nachr. 278 (2005), 312–317.
- [18] E. Straube, The L^2 -Sobolev theory of the $\overline{\partial}$ -Neumann problem. ESI Lectures in Mathematics and Physics, EMS, 2010.

Friedrich Haslinger Fakultät für Mathematik Universität Wien Nordbergstr. 15 A-1090 Wien Austria e-mail: friedrich.haslinger@univie.ac.at