

# Representation Theory of Groups - Blatt 1

11:30-12:15, Seminarraum 9, Oskar-Morgenstern-Platz 1, 2.Stock

[http://www.mat.univie.ac.at/~gagt/rep\\_theory2016](http://www.mat.univie.ac.at/~gagt/rep_theory2016)

Goulnara Arzhantseva

[goulnara.arzhantseva@univie.ac.at](mailto:goulnara.arzhantseva@univie.ac.at)

Martin Finn-Sell

[martin.finn-sell@univie.ac.at](mailto:martin.finn-sell@univie.ac.at)

We begin by recalling a few notions from linear algebra. Let  $V$  and  $W$  be  $k$ -vector spaces.

- The set  $\text{Hom}(V, W)$  of all linear maps from  $V$  to  $W$  is a vector space with pointwise operations;
- The tensor product  $V \otimes W$  is the universal vector space for bilinear maps: that is  $V \otimes W$  is the quotient of the free vector space on the set  $V \times W$  under the equivalence relation generated by:

$$(v_1, w) + (v_2, w) = (v_1 + v_2, w)$$

$$(v, w_1) + (v, w_2) = (v, w_1 + w_2)$$

$$\lambda(v, w) = (\lambda v, w) = (v, \lambda w).$$

We denote elements of  $V \otimes W$  by  $v \otimes w$ , moreover if  $\{e_i\}_i$  and  $\{f_j\}_j$  are bases for  $V$  and  $W$  respectively,  $\{e_i \otimes f_j\}_{i,j}$  is a basis for  $V \otimes W$ . Finally, as an example, one should show that  $\mathbb{C}^n \otimes \mathbb{C}^m \cong \mathbb{C}^{nm}$  as finite dimensional vector spaces.

- Let  $\{e_i\}_{i=1}^{\dim(V)}$  be a basis of  $V$ . The dual space to  $V$ , denoted  $V^*$ , is the vector space of all linear functions  $V \rightarrow k$  with pointwise operations, which is spanned by the linear extensions of the functions  $f_i(e_i) = 1$ . If  $V$  has an inner product, and  $\{e_i\}_i$  is orthonormal with respect to that inner product, then  $V^*$  also has an inner product such that  $\{f_i\}$  are orthonormal. If  $V$  is finite dimensional, then  $V \cong V^*$ , but otherwise not.

**Question 1.** Let  $V$  and  $W$  be finite dimensional  $k$ -vector spaces for some field  $k$ . Show that  $\text{Hom}(V, W) \cong V^* \otimes W$  is an isomorphism of vector spaces, where  $V^*$  denotes the dual vector space of  $V$ .

**Question 2.** Let  $G$  be a finite group,  $X$  be a finite set on which  $G$  acts,  $\rho$  denote the corresponding permutation representation and  $\chi$  the corresponding character. For every  $g \in G$  show that  $\chi(g)$  is the number of elements of  $X$  fixed by  $g$ .

**Question 3.** Let  $V$  be a finite dimensional vector space with a representation  $\rho : G \rightarrow \text{GL}(V)$  and let  $V^*$  be the vector space dual of  $V$ . Let  $\langle v, \phi \rangle = \phi(v)$  be the natural pairing between the dual space  $V^*$  and  $V$ .

1. Show that there is a unique representation  $\rho^* : G \rightarrow GL(V^*)$  satisfying:

$$\langle \rho(v), \rho^*(s)(\phi) \rangle = \langle v, \phi \rangle;$$

2. By fixing a basis for  $V$  and using the dual basis for  $V^*$ , show that  $\rho^*(g) = \rho(g^{-1})^T$ .

**Question 4.** Let  $p$  be a prime number. Show that any  $p$ -group  $G$  has a faithful irreducible representation if and only if the centre  $Z(G)$  is cyclic.

**Question 5.** Let  $G$  be a finite group.

a) Show that for any irreducible representation  $\rho$  of  $G$  with character  $\chi$ , the set

$$\ker(\chi) := \{g \in G \mid \chi(g) = \chi(1)\}$$

is a normal subgroup of  $G$ ;

b) Show that each normal subgroup  $N \triangleleft G$  is the intersection of subgroups of the form  $\ker(\chi)$ , where  $\chi$  is the character of an irreducible representation.

**Question 6.** Let  $\rho_1 : G \rightarrow GL(V_1)$  and  $\rho_2 : G \rightarrow GL(V_2)$  be two representations with characters  $\chi_1$  and  $\chi_2$  respectively, and let  $W := \text{Hom}(V_1, V_2)$ . For every  $g \in G$  and  $f \in W$ , denote by  $\rho(g)f$  the element of  $W$  given by:

$$\rho(g)f = \rho_2(g) \circ f \circ \rho_1^{-1}(g).$$

a) Show that  $\rho : G \rightarrow GL(W)$  is a representation of  $G$ ;

b) Show that the character of  $\rho$  is  $\chi_1^* \chi_2$ , where  $\chi_1^*$  denotes the character of the dual representation of  $\rho_1$  defined in Question 3;

c) Show that  $\rho$  is isomorphic as a representation to the representation  $\rho_1^* \otimes \rho_2$ .

(Note that here, we are considering the tensor product representation of  $G$ , not  $G \times G$ . This is obtained by composing with the map:  $G \rightarrow G \times G$ , and using the tensor product of the representations  $\rho_i$ ; this has formula:  $\rho_1^* \otimes \rho_2(g) = \rho_1^*(g) \otimes \rho_2(g)$  in this case).