

CHOOSING ROOTS OF POLYNOMIALS SMOOTHLY

DMITRI ALEKSEEVSKY
 ANDREAS KRIEGL
 MARK LOSIK
 PETER W. MICHOR

Erwin Schrödinger International Institute
 of Mathematical Physics, Wien, Austria

ABSTRACT. We clarify the question whether for a smooth curve of polynomials one can choose the roots smoothly and related questions. Applications to perturbation theory of operators are given.

TABLE OF CONTENTS

1. Introduction		1
2. Choosing differentiable square and cubic roots		2
3. Choosing local roots of real polynomials smoothly		7
4. Choosing global roots of polynomials differentiably		11
5. The real analytic case		14
6. Choosing roots of complex polynomials		16
7. Choosing eigenvalues and eigenvectors of matrices smoothly		18

1. INTRODUCTION

We consider the following problem. Let

$$(1) \quad P(t) = x^n - a_1(t)x^{n-1} + \dots + (-1)^n a_n(t)$$

be a polynomial with all roots real, smoothly parameterized by t near 0 in \mathbb{R} . Can we find n smooth functions $x_1(t), \dots, x_n(t)$ of the parameter t defined near 0, which are the roots of $P(t)$ for each t ? We can reduce the problem to $a_1 = 0$, replacing the variable x by with the variable $y = x - a_1(t)/n$. We will say that the curve (1) is smoothly solvable near $t = 0$ if such smooth roots $x_i(t)$ exist.

We describe an algorithm which in the smooth and in the holomorphic case sometimes allows to solve this problem. The main results are: If all roots are real then they can always be chosen differentiable, but in general not C^1 for degree

1991 *Mathematics Subject Classification.* 26C10.

Key words and phrases. smooth roots of polynomials.

Supported by 'Fonds zur Förderung der wissenschaftlichen Forschung, Projekt P 10037 PHY'.

$n \geq 3$; and in degree 2 they can be chosen twice differentiable but in general not C^2 . If they are arranged in increasing order, they depend continuously on the coefficients of the polynomial, and if moreover no two of them meet of infinite order in the parameter, then they can be chosen smoothly. We also apply these results to obtain a smooth 1-parameter perturbation theorem for selfadjoint operators with compact resolvent under the condition, that no pair of eigenvalues meets of infinite order.

We thank C. Fefferman who found a mistake in a first version of 2.4, to M. and Th. Hoffmann-Ostenhof for their interest and hints, and to Jerry Kazdan for arranging [15].

2. CHOOSING DIFFERENTIABLE SQUARE AND CUBIC ROOTS

2.1. Proposition. The case $n = 2$. *Let $P(t)(x) = x^2 - f(t)$ for a function f defined on an open interval, such that $f(t) \geq 0$ for all t .*

If f is smooth and is nowhere flat of infinite order, then smooth solutions x exist.

If f is C^2 then C^1 -solutions exist.

If f is C^4 then twice differentiable solutions exist.

Proof. Suppose that f is smooth. If $f(t_0) > 0$ then we have obvious local smooth solutions $\pm\sqrt{f(t)}$. If $f(t_0) = 0$ we have to find a smooth function x such that $f = x^2$, a smooth square root of f . If f is not flat at t_0 then the first nonzero derivative at t_0 has even order $2m$ and is positive, and $f(t) = (t - t_0)^{2m} f_{2m}(t)$, where $f_{2m}(t) := \int_0^1 \frac{(1-r)^{2m-1}}{(2m-1)!} f^{(2m)}(t_0 + r(t - t_0)) dr$ gives a smooth function and $f_{2m}(t_0) = \frac{1}{(2m)!} f^{(2m)}(t_0) > 0$. Then $x(t) := (t - t_0)^m \sqrt{f_{2m}(t)}$ is a local smooth solution. One can piece together these local solutions, changing sign at all points where the first non-vanishing derivative of f is of order $2m$ with m odd. These points are discrete.

Let us consider now a function $f \geq 0$ of class C^2 . We claim that then $x^2 = f(t)$ admits a C^1 -root $x(t)$, globally in t . We consider a fixed t_0 . If $f(t_0) > 0$ then there is locally even a C^2 -solution $x_{\pm}(t) = \pm\sqrt{f(t)}$. If $f(t_0) = 0$ then $f(t) = (t - t_0)^2 h(t)$ where $h \geq 0$ is continuous and C^2 off t_0 with $h(t_0) = \frac{1}{2} f''(t_0)$. If $h(t_0) > 0$ then $x_{\pm}(t) = \pm(t - t_0)\sqrt{h(t)}$ is C^2 off t_0 , and

$$x'_{\pm}(t_0) = \lim_{t \rightarrow t_0} \frac{x_{\pm}(t) - x_{\pm}(t_0)}{t - t_0} = \lim_{t \rightarrow t_0} \pm\sqrt{h(t)} = \pm\sqrt{h(t_0)} = \pm\sqrt{\frac{1}{2}f''(t_0)}.$$

If $h(t_0) = 0$ then we choose $x_{\pm}(t_0) = 0$, and any choice of the roots is then differentiable at t_0 with derivative 0, by the same calculation.

One can piece together these local solutions: At zeros t of f where $f''(t) > 0$ we have to pass through 0, but where $f''(t) = 0$ the choice of the root does not matter. The set $\{t : f(t) = f''(t) = 0\}$ is closed, so its complement is a union of open intervals. Choose a point in each of these intervals where $f(t) > 0$ and start there with the positive root, changing signs at points where $f(t) = 0 \neq f''(t)$: these points do not accumulate in the intervals. Then we get a differentiable choice of a root $x(t)$ on each of this open intervals which extends to a global differentiable root which is 0 on $\{t : f(t) = f''(t) = 0\}$.

Note that we have

$$x'(t) = \begin{cases} \frac{f'(t)}{2x(t)} & \text{if } f(t) > 0 \\ \pm\sqrt{f''(t)/2} & \text{if } f(t) = 0 \end{cases}$$

In points t_0 with $f(t_0) > 0$ the solution x is C^2 ; locally around points t_0 with $f(t_0) = 0$ and $f''(t_0) > 0$ the root x is C^1 since for $t \neq t_0$ near t_0 we have $f(t) > 0$ and $f'(t) \neq 0$, so by l'Hospital we get

$$\lim_{t \rightarrow t_0} x'(t)^2 = \lim_{t \rightarrow t_0} \frac{f'(t)^2}{4f(t)} = \lim_{t \rightarrow t_0} \frac{2f'(t)f''(t)}{4f'(t)} = \frac{f''(t_0)}{2} = x'(t_0)^2,$$

and since the choice of signs was coherent, x' is continuous at t_0 ; if $f''(t_0) = 0$ then $x'(t_0) = 0$ and $x'(t) \rightarrow 0$ for $t \rightarrow t_0$ for both expressions, by lemma 2.2 below. Thus x is C^1 .

If moreover $f \geq 0$ is C^4 , then the solution x from above may be modified to be twice differentiable. Near points t_0 with $f(t_0) > 0$ any continuous root $t \mapsto x_{\pm}(t) = \pm\sqrt{f(t)}$ is even C^4 . Near points t_0 with $f(t_0) = f'(t_0) = 0$ we have $f(t) = (t - t_0)^2 h(t)$ where $h \geq 0$ is C^2 . We may choose a C^1 -root z with $z^2 = h$ by the arguments above, and then $x(t) := (t - t_0)z(t)$ is twice differentiable at t_0 since we have

$$\begin{aligned} \frac{x'(t) - x'(t_0)}{t - t_0} &= \frac{z(t) + (t - t_0)z'(t) - z(t_0)}{t - t_0} \\ &= z'(t) + \frac{z(t) - z(t_0)}{t - t_0} \rightarrow 2z'(t_0) = \pm\sqrt{\frac{f^{(4)}(t_0)}{4!}}. \end{aligned}$$

If $f(t_0) = f''(t_0) = f^{(4)}(t_0) = 0$ then any C^1 choice of the roots is twice differentiable at t_0 , in particular $x(t) = |t - t_0|z(t)$.

Now we can piece together this solutions similarly as above. Let y be a global C^1 -root of f , chosen as above changing sign only at points t with $f(t) = 0 < f''(0)$. We put $x(t) = \varepsilon(t)y(t)$, where $\varepsilon(t) \in \{\pm 1\}$ will be chosen later. The set $\{t : f(t) = f''(t) = f^{(4)}(t) = 0\}$ has a countable union of open intervals as complements. In each of these intervals choose a point t_0 with $f(t_0) > 0$, near which y is C^4 . Now let $\varepsilon(t_0) = 1$, and let ε change sign exactly at points with $f(t) = f''(t) = 0$ but $f^{(4)}(t) > 0$. These points do not accumulate inside the interval. Then x is twice differentiable. \square

2.2. Lemma. *Let $f \geq 0$ be a C^2 -function with $f(t_0) = 0$, then for all $t \in \mathbb{R}$ we have*

$$(1) \quad f'(t)^2 \leq 2f(t) \max\{f''(t_0 + r(t - t_0)) : 0 \leq r \leq 2\}.$$

Proof. If $f(t) = 0$ then $f'(t) = 0$ so (1) holds. We use the Taylor formula

$$(2) \quad f(t + s) = f(t) + f'(t)s + \int_0^1 (1 - r)f''(t + rs) dr s^2$$

In particular we get (replacing t by t_0 and then $t_0 + s$ by t)

$$(3) \quad \begin{aligned} f(t) &= 0 + 0 + \int_0^1 (1-r)f''(t_0 + r(t-t_0)) dr (t-t_0)^2 \\ &\leq \frac{(t-t_0)^2}{2} \max\{f''(t_0 + r(t-t_0)) : 0 \leq r \leq 1\} \end{aligned}$$

Now in (2) we replace s by $-\varepsilon s$ (where $\varepsilon = \text{sign}(f'(t))$) to obtain

$$(4) \quad 0 \leq f(t - \varepsilon s) = f(t) - |f'(t)|s + \int_0^1 (1-r)f''(t - \varepsilon rs) dr s^2$$

Let us assume $t \geq t_0$ and then choose (using (3))

$$s(t) := \sqrt{\frac{2f(t)}{\max\{f''(t_0 + r(t-t_0)) : 0 \leq r \leq 1\}}} \leq t - t_0.$$

Note that we may assume $f(t) > 0$, then $s(t)$ is well defined and $s(t) > 0$. This choice of s in (4) gives

$$\begin{aligned} |f'(t)| &\leq \frac{1}{s(t)} \left(f(t) + \frac{s(t)^2}{2} \max\{f''(t - \varepsilon rs(t)) : 0 \leq r \leq 1\} \right) \\ &\leq \frac{1}{s(t)} \left(f(t) + \frac{s(t)^2}{2} \max\{f''(t - r(t-t_0)) : -1 \leq r \leq 1\} \right) \\ &= \frac{2f(t)}{s(t)} = \sqrt{2f(t) \max\{f''(t_0 + r(t-t_0)) : 0 \leq r \leq 1\}} \end{aligned}$$

which proves (1) for $t \geq t_0$. Since the assertion is symmetric it then holds for all t . \square

2.3. Examples. If $f \geq 0$ is only C^1 , then there may not exist a differentiable root of $x^2 = f(t)$, as the following example shows: $x^2 = f(t) := t^2 \sin^2(\log t)$ is C^1 , but $\pm t \sin(\log t)$ is not differentiable at 0.

If $f \geq 0$ is twice differentiable there may not exist a C^1 -root: $x^2 = f(t) = t^4 \sin^2(\frac{1}{t})$ is twice differentiable, but $\pm t^2 \sin(\frac{1}{t})$ is differentiable, but not C^1 .

If $f \geq 0$ is only C^3 , then there may not exist a twice differentiable root of $x^2 = f(t)$, as the following example shows: $x^2 = f(t) := t^4 \sin^2(\log t)$ is C^3 , but $\pm t^2 \sin(\log t)$ is only C^1 and not twice differentiable.

2.4. Example. If $f(t) \geq 0$ is smooth but flat at 0, in general our problem has no C^2 -root as the following example shows, which is an application of the general curve lemma 4.2.15 in [5]: Let $h : \mathbb{R} \rightarrow [0, 1]$ be smooth with $h(t) = 1$ for $t \geq 0$ and $h(t) = 0$ for $t \leq -1$. Then the function

$$\begin{aligned} f(t) &:= \sum_{n=1}^{\infty} h_n(t-t_n) \cdot \left(\frac{2n}{2^n} (t-t_n)^2 + \frac{1}{4^n} \right), \quad \text{where} \\ h_n(t) &:= h \left(n^2 \left(\frac{1}{n \cdot 2^{n+1}} + t \right) \right) \cdot h \left(n^2 \left(\frac{1}{n \cdot 2^{n+1}} - t \right) \right) \quad \text{and} \\ t_n &:= \sum_{k=1}^{n-1} \left(\frac{2}{k^2} + \frac{2}{k \cdot 2^{k+1}} \right) + \frac{1}{n^2} + \frac{1}{n \cdot 2^{n+1}}, \end{aligned}$$

is ≥ 0 and is smooth: the sum consists of at most one summand for each t , and the derivatives of the summands converge uniformly to 0: Note that $h_n(t) = 1$ for $|t| \leq \frac{1}{n \cdot 2^{n+1}}$ and $h_n(t) = 0$ for $|t| \geq \frac{1}{n \cdot 2^{n+1}} + \frac{1}{n^2}$ hence $h_n(t - t_n) \neq 0$ only for $r_n < t < r_{n+1}$, where $r_n := \sum_{k=1}^{n-1} \left(\frac{2}{k^2} + \frac{2}{k \cdot 2^{k+1}} \right)$. Let $c_n(s) := \frac{2n}{2^n} s^2 + \frac{1}{4^n} \geq 0$ and $H_i := \sup\{|h^{(i)}(t)| : t \in \mathbb{R}\}$. Then

$$\begin{aligned} n^2 \sup\{|(h_n \cdot c_n)^{(k)}(t)| : t \in \mathbb{R}\} &= n^2 \sup\{|(h_n \cdot c_n)^{(k)}(t)| : |t| \leq \frac{1}{n \cdot 2^{n+1}} + \frac{1}{n^2}\} \\ &\leq n^2 \sum_{i=0}^k \binom{k}{i} n^{2i} H_i \sup\{|c_n^{(k-i)}(t)| : |t| \leq \frac{1}{n \cdot 2^{n+1}} + \frac{1}{n^2}\} \\ &\leq \left(\sum_{i=0}^k \binom{k}{i} n^{2i+2} H_i \right) \sup\{|c_n^{(j)}(t)| : |t| \leq 2, j \leq k\} \end{aligned}$$

and since c_n is rapidly decreasing in $C^\infty(\mathbb{R}, \mathbb{R})$ (i.e. $\{p(n) c_n : n \in \mathbb{N}\}$ is bounded in $C^\infty(\mathbb{R}, \mathbb{R})$ for each polynomial p) the right side of the inequality above is bounded with respect to $n \in \mathbb{N}$ and hence the series $\sum_n h_n(\cdot - t_n) c_n(\cdot - t_n)$ converges uniformly in each derivative, and thus represents an element of $f \in C^\infty(\mathbb{R}, \mathbb{R})$. Moreover we have

$$f(t_n) = \frac{1}{4^n}, \quad f'(t_n) = 0, \quad f''(t_n) = \frac{2n}{2^{n-1}}.$$

Let us assume that $f(t) = g(t)^2$ for t near $\sup_n t_n < \infty$, where g is twice differentiable. Then

$$\begin{aligned} f' &= 2gg' \\ f'' &= 2gg'' + 2(g')^2 \\ 2ff'' &= 4g^3g'' + (f')^2 \\ 2f(t_n)f''(t_n) &= 4g(t_n)^3g''(t_n) + f'(t_n)^2 \end{aligned}$$

thus $g''(t_n) = \pm 2n$, so g cannot be C^2 , and g' cannot satisfy a local Lipschitz condition near $\lim t_n$. Another similar example can be found in 7.4 below.

According to [15], some results of this section are contained in Frank Warners dissertation (around 1963, unpublished): Non-negative smooth functions have C^1 square roots whose second derivatives exist everywhere. If all zeros are of finite order there are smooth square roots. However, there are examples not possessing a C^2 square root. Here is one:

$$f(t) = \sin^2(1/t)e^{-1/t} + e^{-2/t} \text{ for } t > 0, \quad f(t) = 0 \text{ for } t \leq 0.$$

This is a sum of two non-negative C^∞ functions each of which has a C^∞ square root. But the second derivative of the square root of f is not continuous at the origin.

In [6] Glaeser proved that a non-negative C^2 -function on an open subset of \mathbb{R}^n which vanishes of second order has a C^1 positive square root. A smooth function $f \geq 0$ is constructed which is flat at 0 such that the positive square root is not C^2 . In [4] Dieudonné gave shorter proofs of Glaeser's results.

2.5. Example. The case $n \geq 3$. We will construct a polynomial

$$P(t) = x^3 + a_2(t)x - a_3(t)$$

with smooth coefficients a_2, a_3 with all roots real which does not admit C^1 -roots. Multiplying with other polynomials one then gets polynomials of all orders $n \geq 3$ which do not admit C^1 -roots.

Suppose that P admits C^1 -roots x_1, x_2, x_3 . Then we have

$$\begin{aligned} 0 &= x_1 + x_2 + x_3 \\ a_2 &= x_1x_2 + x_2x_3 + x_3x_1 \\ a_3 &= x_1x_2x_3 \\ 0 &= \dot{x}_1 + \dot{x}_2 + \dot{x}_3 \\ \dot{a}_2 &= \dot{x}_1x_2 + x_1\dot{x}_2 + \dot{x}_2x_3 + x_2\dot{x}_3 + \dot{x}_3x_1 + x_3\dot{x}_1 \\ \dot{a}_3 &= \dot{x}_1x_2x_3 + x_1\dot{x}_2x_3 + x_1x_2\dot{x}_3 \end{aligned}$$

We solve the linear system formed by the last three equations and get

$$\begin{aligned} \dot{x}_1 &= \frac{\dot{a}_3 - \dot{a}_2x_1}{(x_2 - x_1)(x_3 - x_1)} \\ \dot{x}_2 &= \frac{\dot{a}_3 - \dot{a}_2x_2}{(x_3 - x_2)(x_1 - x_2)} \\ \dot{x}_3 &= \frac{\dot{a}_3 - \dot{a}_2x_3}{(x_1 - x_3)(x_2 - x_3)}. \end{aligned}$$

We consider the continuous function

$$b_3 := \dot{x}_1\dot{x}_2\dot{x}_3 = \frac{\dot{a}_3^3 + \dot{a}_2^2\dot{a}_3a_2 - \dot{a}_2^3a_3}{4a_2^3 + 27a_3^2}.$$

For smooth functions f and ε with $\varepsilon^2 \leq 1$ we let

$$\begin{aligned} u &:= -12a_2 := f^2 \\ v &:= 108a_3 := \varepsilon f^3. \end{aligned}$$

then all roots of P are real since $a_2 \leq 0$ and $432(4a_2^3 + 27a_3^2) = v^2 - u^3 = f^6(\varepsilon^2 - 1) \leq 0$. We get then

$$\begin{aligned} 11664b_3 &= \frac{4\dot{v}^3 - 27u\dot{u}^2\dot{v} + 27\dot{u}^3v}{v^2 - u^3} \\ &= 4 \frac{f^3\dot{\varepsilon}^3 + 9f^2\dot{f}\dot{\varepsilon}^2\varepsilon + 27f\dot{f}^2\dot{\varepsilon}(\varepsilon^2 - 1) + 27\dot{f}^3\varepsilon(\varepsilon^2 - 3)}{\varepsilon^2 - 1}. \end{aligned}$$

Now we choose

$$\begin{aligned} f(t) &:= \sum_{n=1}^{\infty} h_n(t - t_n) \cdot \left(\frac{n}{2^n} (t - t_n) \right), \\ \varepsilon(t) &:= 1 - \sum_{n=1}^{\infty} h_n(t - t_n) \cdot \frac{1}{8^n}, \end{aligned}$$

where h_n and t_n are as in the beginning of 2.4. Then $f(t_n) = 0$, $\dot{f}(t_n) = \frac{n}{2^n}$, and $\varepsilon(t_n) = \frac{1}{8^n}$, hence

$$108b_3(t_n) = \frac{\dot{f}(t_n)^3 \varepsilon(t_n) (\varepsilon(t_n)^2 - 3)}{\varepsilon(t_n)^2 - 1} \sim n^3 \rightarrow \infty$$

is unbounded on the convergent sequence t_n . So the roots cannot be chosen locally Lipschitz, thus not C^1 .

3. CHOOSING LOCAL ROOTS OF REAL POLYNOMIALS SMOOTHLY

3.1. Preliminaries. We recall some known facts on polynomials with real coefficients. Let

$$P(x) = x^n - a_1 x^{n-1} + \cdots + (-1)^n a_n$$

be a polynomial with real coefficients a_1, \dots, a_n and roots $x_1, \dots, x_n \in \mathbb{C}$. It is known that $a_i = \sigma_i(x_1, \dots, x_n)$, where σ_i ($i = 1, \dots, n$) are the elementary symmetric functions in n variables:

$$\sigma_i(x_1, \dots, x_n) = \sum_{1 \leq j_1 < \cdots < j_i \leq n} x_{j_1} \cdots x_{j_i}$$

Denote by s_i the Newton polynomials $\sum_{j=1}^n x_j^i$ which are related to the elementary symmetric function by

$$(1) \quad s_k - s_{k-1} \sigma_1 + s_{k-2} \sigma_2 + \cdots + (-1)^{k-1} s_1 \sigma_{k-1} + (-1)^k k \sigma_k = 0 \quad (k \leq n)$$

The corresponding mappings are related by a polynomial diffeomorphism ψ^n , given by (1):

$$\begin{aligned} \sigma^n &:= (\sigma_1, \dots, \sigma_n) : \mathbb{R}^n \rightarrow \mathbb{R}^n \\ s^n &:= (s_1, \dots, s_n) : \mathbb{R}^n \rightarrow \mathbb{R}^n \\ s^n &:= \psi^n \circ \sigma^n \end{aligned}$$

Note that the Jacobian (the determinant of the derivative) of s^n is $n!$ times the Vandermonde determinant: $\det(ds^n(x)) = n! \prod_{i>j} (x_i - x_j) =: n! \text{Van}(x)$, and even the derivative itself $d(s^n)(x)$ equals the Vandermonde matrix up to factors i in the i -th row. We also have $\det(d(\psi^n)(x)) = (-1)^{n(n+3)/2} n! = (-1)^{n(n-1)/2} n!$, and consequently $\det(d\sigma^n(x)) = \prod_{i>j} (x_j - x_i)$. We consider the so-called Bezoutiant

$$B := \begin{pmatrix} s_0 & s_1 & \cdots & s_{n-1} \\ s_1 & s_2 & \cdots & s_n \\ \vdots & \vdots & & \vdots \\ s_{n-1} & s_n & \cdots & s_{2n-2} \end{pmatrix}.$$

Let B_k be the minor formed by the first k rows and columns of B . From

$$B_k(x) = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ x_1 & x_2 & \cdots & x_n \\ \vdots & \vdots & & \vdots \\ x_1^{k-1} & x_2^{k-1} & \cdots & x_n^{k-1} \end{pmatrix} \cdot \begin{pmatrix} 1 & x_1 & \cdots & x_1^{k-1} \\ 1 & x_2 & \cdots & x_2^{k-1} \\ \vdots & \vdots & & \vdots \\ 1 & x_n & \cdots & x_n^{k-1} \end{pmatrix}$$

it follows that

$$(2) \quad \Delta_k(x) := \det(B_k(x)) = \sum_{i_1 < i_2 < \dots < i_k} (x_{i_1} - x_{i_2})^2 \dots (x_{i_1} - x_{i_k})^2 \dots (x_{i_{k-1}} - x_{i_k})^2,$$

since for $n \times k$ -matrices A one has $\det(AA^\top) = \sum_{i_1 < \dots < i_k} \det(A_{i_1, \dots, i_k})^2$, where A_{i_1, \dots, i_k} is the minor of A with indicated rows. Since the Δ_k are symmetric we have $\Delta_k = \tilde{\Delta}_k \circ \sigma^n$ for unique polynomials $\tilde{\Delta}_k$ and similarly we shall use \tilde{B} .

3.2. Theorem. (*Sylvester's version of Sturm's theorem, see [14], [10]*) *The roots of P are all real if and only if the symmetric $(n \times n)$ matrix $\tilde{B}(P)$ is positive semidefinite; then $\tilde{\Delta}_k(P) := \tilde{\Delta}_k(a_1, \dots, a_n) \geq 0$ for $1 \leq k \leq n$. The rank of B equals the number of distinct roots of P and its signature equals the number of distinct real roots.*

3.3. Proposition. *Let now P be a smooth curve of polynomials*

$$P(t)(x) = x^n - a_1(t)x^{n-1} + \dots + (-1)^n a_n(t)$$

with all roots real, and distinct for $t = 0$. Then P is smoothly solvable near 0.

This is also true in the real analytic case and for higher dimensional parameters, and in the holomorphic case for complex roots.

Proof. The derivative $\frac{d}{dx}P(0)(x)$ does not vanish at any root, since they are distinct. Thus by the implicit function theorem we have local smooth solutions $x(t)$ of $P(t, x) = P(t)(x) = 0$. \square

3.4. Splitting Lemma. *Let P_0 be a polynomial*

$$P_0(x) = x^n - a_1x^{n-1} + \dots + (-1)^n a_n.$$

If $P_0 = P_1 \cdot P_2$, where P_1 and P_2 are polynomials with no common root. Then for P near P_0 we have $P = P_1(P)\tilde{P}_2(P)$ for real analytic mappings of monic polynomials $P \mapsto P_1(P)$ and $P \mapsto P_2(P)$, defined for P near P_0 , with the given initial values.

Proof. Let the polynomial P_0 be represented as the product

$$P_0 = P_1 \cdot P_2 = (x^p - b_1x^{p-1} + \dots + (-1)^p b_p)(x^q - c_1x^{q-1} + \dots + (-1)^q c_q),$$

let x_i for $i = 1, \dots, n$ be the roots of P_0 , ordered in such a way that for $i = 1, \dots, p$ we get the roots of P_1 , and for $i = p+1, \dots, p+q = n$ we get those of P_2 . Then $(a_1, \dots, a_n) = \phi^{p,q}(b_1, \dots, b_p, c_1, \dots, c_q)$ for a polynomial mapping $\phi^{p,q}$ and we get

$$\begin{aligned} \sigma^n &= \phi^{p,q} \circ (\sigma^p \times \sigma^q), \\ \det(d\sigma^n) &= \det(d\phi^{p,q}(b, c)) \det(d\sigma^p) \det(d\sigma^q). \end{aligned}$$

From 3.1 we conclude

$$\prod_{1 \leq i < j \leq n} (x_i - x_j) = \det(d\phi^{p,q}(b, c)) \prod_{1 \leq i < j \leq p} (x_i - x_j) \prod_{p+1 \leq i < j \leq n} (x_i - x_j)$$

which in turn implies

$$\det(d\phi^{p,q}(b, c)) = \prod_{1 \leq i \leq p < j \leq n} (x_i - x_j) \neq 0$$

so that $\phi^{p,q}$ is a real analytic diffeomorphism near (b, c) . \square

3.5. For a continuous function f defined near 0 in \mathbb{R} let the *multiplicity* or *order of flatness* $m(f)$ at 0 be the supremum of all integer p such that $f(t) = t^p g(t)$ near 0 for a continuous function g . If f is C^n and $m(f) < n$ then $f(t) = t^{m(f)} g(t)$ where now g is $C^{n-m(f)}$ and $g(0) \neq 0$. If f is a continuous function on the space of polynomials, then for a fixed continuous curve P of polynomials we will denote by $m(f)$ the multiplicity at 0 of $t \mapsto f(P(t))$.

The splitting lemma 3.4 shows that for the problem of smooth solvability it is enough to assume that all roots of $P(0)$ are equal.

Proposition. *Suppose that the smooth curve of polynomials*

$$P(t)(x) = x^n + a_2(t)x^{n-2} - \cdots + (-1)^n a_n(t)$$

is smoothly solvable with smooth roots $t \mapsto x_i(t)$, and that all roots of $P(0)$ are equal. Then for $(k = 2, \dots, n)$

$$\begin{aligned} m(\tilde{\Delta}_k) &\geq k(k-1) \min_{1 \leq i \leq n} m(x_i) \\ m(a_k) &\geq k \min_{1 \leq i \leq n} m(x_i) \end{aligned}$$

This result also holds in the real analytic case and in the smooth case.

Proof. This follows from 3.1.(2) for Δ_k , and from $a_k(t) = \sigma_k(x_1(t), \dots, x_n(t))$. \square

3.6. Lemma. *Let P be a polynomial of degree n with all roots real. If $a_1 = a_2 = 0$ then all roots of P are equal to zero.*

Proof. From 3.1.(1) we have $\sum x_i^2 = s_2(x) = \sigma_1^2(x) - 2\sigma_2(x) = a_1^2 - 2a_2 = 0$. \square

3.7. Multiplicity lemma. *Consider a smooth curve of polynomials*

$$P(t)(x) = x^n + a_2(t)x^{n-2} - \cdots + (-1)^n a_n(t)$$

with all roots real. Then, for integers r , the following conditions are equivalent:

- (1) $m(a_k) \geq kr$ for all $2 \leq k \leq n$.
- (2) $m(\tilde{\Delta}_k) \geq k(k-1)r$ for all $2 \leq k \leq n$.
- (3) $m(a_2) \geq 2r$.

Proof. We only have to treat $r > 0$.

(1) implies (2): From 3.1.(1) we have $m(\tilde{s}_k) \geq rk$, and from the definition of $\tilde{\Delta}_k = \det(\tilde{B}_k)$ we get (2).

(2) implies (3) since $\tilde{\Delta}_2 = -2na_2$.

(3) implies (1): From $a_2(0) = 0$ and lemma 3.6 it follows that all roots of the polynomial $P(0)$ are equal to zero and, then, $a_3(0) = \cdots = a_n(0) = 0$. Therefore, $m(a_3), \dots, m(a_n) \geq 1$. Under these conditions, we have $a_2(t) = t^{2r} a_{2,2r}(t)$ and $a_k(t) = t^{m_k} a_{k,m_k}(t)$ for $k = 3, \dots, n$, where the m_k are positive integers and $a_{2,2r}, a_{3,m_3}, \dots, a_{n,m_n}$ are smooth functions, and where we may assume that either $m_k = m(a_k) < \infty$ or $m_k \geq kr$.

Suppose now indirectly that for some $k > 2$ we have $m_k = m(a_k) < kr$. Then we put

$$m := \min\left(r, \frac{m_3}{3}, \dots, \frac{m_n}{n}\right) < r.$$

We consider the following continuous curve of polynomials for $t \geq 0$:

$$\begin{aligned} \bar{P}_m(t)(x) := & x^n + a_{2,2r}(t)t^{2r-2m}x^{n-2} \\ & - a_{3,m_3}(t)t^{m_3-3m}x^{n-3} + \dots + (-1)^n a_{n,m_n}(t)t^{m_n-nm}. \end{aligned}$$

If x_1, \dots, x_n are the real roots of $P(t)$ then $t^{-m}x_1, \dots, t^{-m}x_n$ are the roots of $\bar{P}_m(t)$, for $t > 0$. So for $t > 0$, $\bar{P}_m(t)$ is a family of polynomials with all roots real. Since by theorem 3.2 the set of polynomials with all roots real is closed, $\bar{P}_m(0)$ is also a polynomial with all roots real.

By lemma 3.6 all roots of the polynomial $\bar{P}_m(0)$ are equal to zero, and for those k with $m_k = km$ we have $a_{k,m_k}(0) = 0$ and, therefore, $m(a_k) > m_k$, a contradiction. \square

3.8. Algorithm. Consider a smooth curve of polynomials

$$P(t)(x) = x^n - a_1(t)x^{n-1} + a_2(t)x^{n-2} - \dots + (-1)^n a_n(t)$$

with all roots real. The algorithm has the following steps:

- (1) If all roots of $P(0)$ are pairwise different, P is smoothly solvable for t near 0 by 3.3.
- (2) If there are distinct roots at $t = 0$ we put them into two subsets which splits $P(t) = P_1(t).P_2(t)$ by the splitting lemma 3.4. We then feed $P_i(t)$ (which have lower degree) into the algorithm.
- (3) All roots of $P(0)$ are equal. We first reduce $P(t)$ to the case $a_1(t) = 0$ by replacing the variable x by $y = x - a_1(t)/n$. Then all roots are equal to 0 so $m(a_2) > 0$.
- (3a) If $m(a_2)$ is finite then it is even since $\tilde{\Delta}_2 = -2na_2 \geq 0$, $m(a_2) = 2r$ and by the multiplicity lemma 3.7 $a_i(t) = a_{i,ir}(t)t^{ir}$ ($i = 2, \dots, n$) for smooth $a_{i,ir}$. Consider the following smooth curve of polynomials

$$P_r(t)(x) = x^n + a_{2,2r}(t)x^{n-2} - a_{3,3r}(t)x^{n-3} \dots + (-1)^n a_{n,nr}(t).$$

If $P_r(t)$ is smoothly solvable and $x_k(t)$ are its smooth roots, then $x_k(t)t^r$ are the roots of $P(t)$ and the original curve P is smoothly solvable too. Since $a_{2,2m}(0) \neq 0$, not all roots of $P_r(0)$ are equal and we may feed P_r into step 2 of the algorithm.

- (3b) If $m(a_2)$ is infinite and $a_2 = 0$, then all roots are 0 by 3.6 and thus the polynomial is solvable.
- (3c) But if $m(a_2)$ is infinite and $a_2 \neq 0$, then by the multiplicity lemma 3.7 all $m(a_i)$ for $2 \leq i \leq n$ are infinite. In this case we keep $P(t)$ as factor of the original curve of polynomials with all coefficients infinitely flat at $t = 0$ after forcing $a_1 = 0$. This means that all roots of $P(t)$ meet of infinite order of flatness (see 3.5) at $t = 0$ for any choice of the roots. This can be seen as follows: If $x(t)$ is any root of $P(t)$ then $y(t) := x(t)/t^r$ is a root of $P_r(t)$, hence by 4.1 bounded, so $x(t) = t^{r-1}.ty(t)$ and $t \mapsto ty(t)$ is continuous at $t = 0$.

This algorithm produces a splitting of the original polynomial

$$P(t) = P^{(\infty)}(t)P^{(s)}(t)$$

where $P^{(\infty)}$ has the property that each root meets another one of infinite order at $t = 0$, and where $P^{(s)}(t)$ is smoothly solvable, and no two roots meet of infinite order at $t = 0$, if they are not equal. Any two choices of smooth roots of $P^{(s)}$ differ by a permutation.

The factor $P^{(\infty)}$ may or may not be smoothly solvable. For a flat function $f \geq 0$ consider:

$$x^4 - (f(t) + t^2)x^2 + t^2 f(t) = (x^2 - f(t))(x - t)(x + t).$$

Here the algorithm produces this factorization. For $f(t) = g(t)^2$ the polynomial is smoothly solvable. For the smooth function f from 2.4 it is not smoothly solvable.

4. CHOOSING GLOBAL ROOTS OF POLYNOMIALS DIFFERENTIABLY

4.1. Lemma. *For a polynomial*

$$P(x) = x^n - a_1(P)x^{n-1} + \cdots + (-1)^n a_n(P)$$

with all roots real, i.e. $\tilde{\Delta}_k(P) = \tilde{\Delta}_k(a_1, \dots, a_n) \geq 0$ for $1 \leq k \leq n$, let

$$x_1(P) \leq x_2(P) \leq \cdots \leq x_n(P)$$

be the roots, increasingly ordered.

Then all $x_i : \sigma^n(\mathbb{R}^n) \rightarrow \mathbb{R}$ are continuous.

Proof. We show first that x_1 is continuous. Let $P_0 \in \sigma^n(\mathbb{R}^n)$ be arbitrary. We have to show that for every $\varepsilon > 0$ there exists some $\delta > 0$ such that for all $|P - P_0| < \delta$ there is a root $x(P)$ of P with $x(P) < x_1(P_0) + \varepsilon$ and for all roots $x(P)$ of P we have $x(P) > x_1(P_0) - \varepsilon$. Without loss of generality we may assume that $x_1(P_0) = 0$.

We use induction on the degree n of P . By the splitting lemma 3.4 for the C^0 -case we may factorize P as $P_1(P) \cdot P_2(P)$, where $P_1(P_0)$ has all roots equal to $x_1 = 0$ and $P_2(P_0)$ has all roots greater than 0 and both polynomials have coefficients which depend real analytically on P . The degree of $P_2(P)$ is now smaller than n , so by induction the roots of $P_2(P)$ are continuous and thus larger than $x_1(P_0) - \varepsilon$ for P near P_0 .

Since 0 was the smallest root of P_0 we have to show that for all $\varepsilon > 0$ there exists a $\delta > 0$ such that for $|P - P_0| < \delta$ any root x of $P_1(P)$ satisfies $|x| < \varepsilon$. Suppose there is a root x with $|x| \geq \varepsilon$. Then we get as follows a contradiction, where n_1 is the degree of P_1 . From

$$-x^{n_1} = \sum_{k=1}^{n_1} (-1)^k a_k(P_1) x^{n_1-k}$$

we have

$$\varepsilon \leq |x| = \left| \sum_{k=1}^{n_1} (-1)^k a_k(P_1) x^{1-k} \right| \leq \sum_{k=1}^{n_1} |a_k(P_1)| |x|^{1-k} < \sum_{k=1}^{n_1} \frac{\varepsilon^k}{n_1} \varepsilon^{1-k} = \varepsilon,$$

provided that $n_1|a_k(P_1)| < \varepsilon^k$, which is true for P_1 near P_0 , since $a_k(P_0) = 0$. Thus x_1 is continuous.

Now we factorize $P = (x - x_1(P)) \cdot P_2(P)$, where $P_2(P)$ has roots $x_2(P) \leq \dots \leq x_n(P)$. By Horner's algorithm ($a_n = b_{n-1}x_1$, $a_{n-1} = b_{n-1} + b_{n-2}x_1$, \dots , $a_2 = b_2 + b_1x_1$, $a_1 = b_1 + x_1$) the coefficients b_k of $P_2(P)$ are again continuous and so we may proceed by induction on the degree of P . Thus the claim is proved. \square

4.2. Theorem. *Consider a smooth curve of polynomials*

$$P(t)(x) = x^n + a_2(t)x^{n-2} - \dots + (-1)^n a_n(t)$$

with all roots real, for $t \in \mathbb{R}$. Let one of the two following equivalent conditions be satisfied:

- (1) *If two of the increasingly ordered continuous roots meet of infinite order somewhere then they are equal everywhere.*
- (2) *Let k be maximal with the property that $\tilde{\Delta}_k(P)$ does not vanish identically for all t . Then $\tilde{\Delta}_k(P)$ vanishes nowhere of infinite order.*

Then the roots of P can be chosen smoothly, and any two choices differ by a permutation of the roots.

Proof. The local situation. We claim that for any t_0 , without loss $t_0 = 0$, the following conditions are equivalent:

- (1) *If two of the increasingly ordered continuous roots meet of infinite order at $t = 0$ then their germs at $t = 0$ are equal.*
- (2) *Let k be maximal with the property that the germ at $t = 0$ of $\tilde{\Delta}_k(P)$ is not 0. Then $\tilde{\Delta}_k(P)$ is not infinitely flat at $t = 0$.*
- (3) *The algorithm 3.8 never leads to step (3c).*

(3) \implies (1). Suppose indirectly that two nonequal of the increasingly ordered continuous roots meet of infinite order at $t = 0$. Then in each application of step (2) these two roots stay with the same factor. After any application of step (3a) these two roots lead to nonequal roots of the modified polynomial which still meet of infinite order at $t = 0$. They never end up in a factor leading to step (3b) or step (1). So they end up in a factor leading to step (3c).

(1) \implies (2). Let $x_1(t) \leq \dots \leq x_n(t)$ be the continuous roots of $P(t)$. From 3.1, (2) we have

$$(4) \quad \tilde{\Delta}_k(P(t)) = \sum_{i_1 < i_2 < \dots < i_k} (x_{i_1} - x_{i_2})^2 \dots (x_{i_1} - x_{i_n})^2 \dots (x_{i_{k-1}} - x_{i_k})^2.$$

The germ of $\tilde{\Delta}_k(P)$ is not 0, so the germ of one summand is not 0. If $\tilde{\Delta}_k(P)$ were infinitely flat at $t = 0$, then each summand is infinitely flat, so there are two roots among the x_i which meet of infinite order, thus by assumption their germs are equal, so this summand vanishes.

(2) \implies (3). Since the leading $\tilde{\Delta}_k(P)$ vanishes only of finite order at zero, P has exactly k different roots off 0. Suppose indirectly that the algorithm 3.8 leads to step (3c), then $P = P^{(\infty)}P^{(s)}$ for a nontrivial polynomial $P^{(\infty)}$. Let $x_1(t) \leq \dots \leq x_p(t)$ be the roots of $P^{(\infty)}(t)$ and $x_{p+1}(t) \leq \dots \leq x_n(t)$ those of $P^{(s)}$. We know

that each x_i meets some x_j of infinite order and does not meet any x_l of infinite order, for $i, j \leq p < l$. Let $k^{(\infty)} > 2$ and $k^{(s)}$ be the number of generically different roots of $P^{(\infty)}$ and $P^{(s)}$, respectively. Then $k = k^{(\infty)} + k^{(s)}$, and an inspection of the formula for $\tilde{\Delta}_k(P)$ above leads to the fact that it must vanish of infinite order at 0, since the only non-vanishing summands involve exactly $k^{(\infty)}$ many generically different roots from $P^{(\infty)}$.

The global situation. From the first part of the proof we see that the algorithm 3.8 allows to choose the roots smoothly in a neighborhood of each point $t \in \mathbb{R}$, and that any two choices differ by a (constant) permutation of the roots. Thus we may glue the local solutions to a global solution. \square

4.3. Theorem. *Consider a curve of polynomials*

$$P(t)(x) = x^n - a_1(t)x^{n-1} + \cdots + (-1)^n a_n(t), \quad t \in \mathbb{R},$$

with all roots real, where all a_i are of class C^n . Then there is a differentiable curve $x = (x_1, \dots, x_n) : \mathbb{R} \rightarrow \mathbb{R}^n$ whose coefficients parameterize the roots.

That this result cannot be improved to C^2 -roots is shown already in 2.4, and not to C^1 for $n \geq 3$ is shown in 2.5.

Proof. First we note that the multiplicity lemma 3.7 remains true in the C^n -case for $r = 1$ in the following sense, with the same proof:

If $a_1 = 0$ then the following two conditions are equivalent

- (1) $a_k(t) = t^k a_{k,k}(t)$ for a continuous function $a_{k,k}$, for $2 \leq k \leq n$.
- (2) $a_2(t) = t^2 a_{2,2}(t)$ for a continuous function $a_{2,2}$.

In order to prove the theorem itself we follow one step of the algorithm. First we replace x by $x + \frac{1}{n}a_1(t)$, or assume without loss that $a_1 = 0$. Then we choose a fixed t , say $t = 0$.

If $a_2(0) = 0$ then it vanishes of second order at 0: if it vanishes only of first order then $\tilde{\Delta}_2(P(t)) = -2na_2(t)$ would change sign at $t = 0$, contrary to the assumption that all roots of $P(t)$ are real, by 3.2. Thus $a_2(t) = t^2 a_{2,2}(t)$, so by the variant of the multiplicity lemma 3.7 described above we have $a_k(t) = t^k a_{k,k}(t)$ for continuous functions $a_{k,k}$, for $2 \leq k \leq n$. We consider the following continuous curve of polynomials

$$P_1(t)(x) = x^n + a_{2,2}(t)x^{n-2} - a_{3,3}(t)x^{n-3} \cdots + (-1)^n a_{n,n}(t).$$

with continuous roots $z_1(t) \leq \cdots \leq z_n(t)$, by 4.1. Then $x_k(t) = z_k(t)t$ are differentiable at 0, and are all roots of P , but note that $x_k(t) = y_k(t)$ for $t \geq 0$, but $x_k(t) = y_{n-k}(t)$ for $t \leq 0$, where $y_1(t) \leq \cdots \leq y_n(t)$ are the ordered roots of $P(t)$. This gives us one choice of differentiable roots near $t = 0$. Any choice is then given by this choice and applying afterwards any permutation of the set $\{1, \dots, n\}$ keeping invariant the function $k \mapsto z_k(0)$.

If $a_2(0) \neq 0$ then by the splitting lemma 3.4 for the C^n -case we may factor $P(t) = P_1(t) \cdots P_k(t)$ where the $P_i(t)$ have again C^n -coefficients and where each $P_i(0)$ has all roots equal to c_i , and where the c_i are distinct. By the arguments

above the roots of each P_i can be arranged differentiably, thus P has differentiable roots $y_k(t)$.

But note that we have to apply a permutation on one side of 0 to the original roots, in the following case: Two roots x_k and x_l meet at zero with $x_k(t) - x_l(t) = tc_{kl}(t)$ with $c_{kl}(0) \neq 0$ (we say that they meet slowly). We may apply to this choice an arbitrary permutation of any two roots which meet with $c_{kl}(0) = 0$ (i.e. at least of second order), and we get thus any differentiable choice near $t = 0$.

Now we show that we choose the roots differentiably on the whole domain \mathbb{R} . We start with with the ordered continuous roots $y_1(t) \leq \dots \leq y_n(t)$. Then we put

$$x_k(t) = y_{\sigma(t)(k)}(t)$$

where the permutation $\sigma(t)$ is given by

$$\sigma(t) = (1, 2)^{\varepsilon_{1,2}(t)} \dots (1, n)^{\varepsilon_{1,n}(t)} (2, 3)^{\varepsilon_{2,3}(t)} \dots (n-1, n)^{\varepsilon_{n-1,n}(t)}$$

and where $\varepsilon_{i,j}(t) \in \{0, 1\}$ will be specified as follows: On the closed set $S_{i,j}$ of all t where $y_i(t)$ and $y_j(t)$ meet of order at least 2 any choice is good. The complement of $S_{i,j}$ is an at most countable union of open intervals, and in each interval we choose a point where we put $\varepsilon_{i,j} = 0$. Going right (and left) from this point we change $\varepsilon_{i,j}$ in each point where y_i and y_j meet slowly. These points accumulate only in $S_{i,j}$. \square

5. THE REAL ANALYTIC CASE

5.1. Theorem. *Let P be a real analytic curve of polynomials*

$$P(t)(x) = x^n - a_1(t)x^{n-1} + \dots + (-1)^n a_n(t), \quad t \in \mathbb{R},$$

with all roots real.

Then P is real analytically solvable, globally on \mathbb{R} . All solutions differ by permutations.

By a real analytic curve of polynomials we mean that all $a_i(t)$ are real analytic in t (but see also [8]), and real analytically solvable means that we may find $x_i(t)$ for $i = 1, \dots, n$ which are real analytic in t and are roots of $P(t)$ for all t . The local existence part of this theorem is due to Rellich [11], Hilfssatz 2, his proof uses Puiseux-expansions. Our proof is different and more elementary.

Proof. We first show that P is locally real analytically solvable, near each point $t_0 \in \mathbb{R}$. It suffices to consider $t_0 = 0$. Using the transformation in the introduction we first assume that $a_1(t) = 0$ for all t . We use induction on the degree n . If $n = 1$ the theorem holds. For $n > 1$ we consider several cases:

The case $a_2(0) \neq 0$. Here not all roots of $P(0)$ are equal and zero, so by the splitting lemma 3.4 we may factor $P(t) = P_1(t) \cdot P_2(t)$ for real analytic curves of polynomials of positive degree, which have both all roots real, and we have reduced the problem to lower degree.

The case $a_2(0) = 0$. If $a_2(t) = 0$ for all t , then by 3.6 all roots of $P(t)$ are 0, and we are done. Otherwise $1 \leq m(a_2) < \infty$ for the multiplicity of a_2 at 0, and

by 3.6 all roots of $P(0)$ are 0. If $m(a_2) > 0$ is odd, then $\tilde{\Delta}_2(P)(t) = -2na_2(t)$ changes sign at $t = 0$, so by 3.2 not all roots of $P(t)$ are real for t on one side of 0. This contradicts the assumption, so $m(a_2) = 2r$ is even. Then by the multiplicity lemma 3.7 we have $a_i(t) = a_{i,ir}(t)t^{ir}$ ($i = 2, \dots, n$) for real analytic $a_{i,ir}$, and we may consider the following real analytic curve of polynomials

$$P_r(t)(x) = x^n + a_{2,2r}(t)x^{n-2} - a_{3,3r}(t)x^{n-3} \cdots + (-1)^n a_{n,nr}(t),$$

with all roots real. If $P_r(t)$ is real analytically solvable and $x_k(t)$ are its real analytic roots, then $x_k(t)t^r$ are the roots of $P(t)$ and the original curve P is real analytically solvable too. Now $a_{2,2r}(0) \neq 0$ and we are done by the case above.

Claim. Let $x = (x_1, \dots, x_n) : I \rightarrow \mathbb{R}^n$ be a real analytic curve of roots of P on an open interval $I \subset \mathbb{R}$. Then any real analytic curve of roots of P on I is of the form $\alpha \circ x$ for some permutation α .

Let $y : I \rightarrow \mathbb{R}^n$ be another real analytic curve of roots of P . Let $t_k \rightarrow t_0$ be a convergent sequence of distinct points in I . Then $y(t_k) = \alpha_k(x(t_k)) = (x_{\alpha_k 1}, \dots, x_{\alpha_k n})$ for permutations α_k . By choosing a subsequence we may assume that all α_k are the same permutation α . But then the real analytic curves y and $\alpha \circ x$ coincide on a converging sequence, so they coincide on I and the claim follows.

Now from the local smooth solvability above and the uniqueness of smooth solutions up to permutations we can glue a global smooth solution on the whole of \mathbb{R} . \square

5.2. Remarks and examples. The uniqueness statement of theorem 5.1 is wrong in the smooth case, as is shown by the following example: $x^2 = f(t)^2$ where f is smooth. In each point t where f is infinitely flat one can change sign in the solution $x(t) = \pm f(t)$. No sign change can be absorbed in a permutation (constant in t). If there are infinitely many points of flatness for f we get uncountably many smooth solutions.

Theorem 5.1 reminds of the curve lifting property of covering mappings. But unfortunately one cannot lift real analytic homotopies, as the following example shows. This example also shows that polynomials which are real analytically parameterized by higher dimensional variables are not real analytically solvable.

Consider the 2-parameter family $x^2 = t_1^2 + t_2^2$. The two continuous solutions are $x(t) = \pm|t|$, but for none of them $t_1 \mapsto x(t_1, 0)$ is differentiable at 0.

There remains the question whether for a real analytic submanifold of the space of polynomials with all roots real one can choose the roots real analytically along this manifold. The following example shows that this is not the case:

Consider

$$P(t_1, t_2)(x) = (x^2 - (t_1^2 + t_2^2))(x - (t_1 - a_1))(x - (t_2 - a_2)),$$

which is not real analytically solvable, see above. For $a_1 \neq a_2$ the coefficients describe a real analytic embedding for (t_1, t_2) near 0.

6. CHOOSING ROOTS OF COMPLEX POLYNOMIALS

6.1. In this section we consider the problem of finding smooth curves of complex roots for smooth curves $t \mapsto P(t)$ of polynomials

$$P(t)(z) = z^n - a_1(t)z^{n-1} + \cdots + (-1)^n a_n(t)$$

with complex valued coefficients $a_1(t), \dots, a_n(t)$. We shall also discuss the real analytic and holomorphic cases. The definition of the Bezoutiant B , its principal minors Δ_k , and formula 3.1.(2) are still valid. Note that now there are no restrictions on the coefficients. In this section the parameter may be real and $P(t)$ may be smooth or real analytic in t , or the parameter t may be complex and $P(t)$ may be holomorphic in t .

6.2. The case $n = 2$. As in the real case the problem reduces to the following one: Let f be a smooth complex valued function, defined near 0 in \mathbb{R} , such that $f(0) = 0$. We look for a smooth function $g : (\mathbb{R}, 0) \rightarrow \mathbb{C}$ such that $f = g^2$. If $m(f)$ is finite and even, we have $f(t) = t^{m(f)}h(t)$ with $h(0) \neq 0$, and $g(t) := t^{m(f)/2}\sqrt{h(t)}$ is a local solution. If $m(f)$ is finite and odd there is no solution g , also not in the real analytic and holomorphic cases. If $f(t)$ is flat at $t = 0$, then one has no definite answer, and the example 2.4 is still not smoothly solvable.

6.3. The general case. Proposition 3.3 and the splitting lemma 3.4 are true in the complex case. Proposition 3.5 is true also because it follows from 3.1.(2). Of course lemma 3.6 is not true now and the multiplicity lemma 3.7 only partially holds:

6.4. Multiplicity Lemma. *Consider the smooth (real analytic, holomorphic) curve of complex polynomials*

$$P(t)(z) = z^n + a_2(t)z^{n-2} - \dots + (-1)^n a_n(t).$$

Then, for integers r , the following conditions are equivalent:

- (1) $m(a_k) \geq kr$ for all $2 \leq k \leq n$.
- (2) $m(\tilde{\Delta}_k) \geq k(k-1)r$ for all $2 \leq k \leq n$.

Proof. (2) implies (1): Since $\tilde{\Delta}_2 = na_2$ we have $s_2(0) = -2a_2(0) = 0$. From $\tilde{\Delta}_3(0) = -s_3(0)^2$ we then get $s_3(0) = 0$, and so on we obtain $s_4(0) = \dots = s_n(0) = 0$. Then by 3.1.(1) $a_i(0) = 0$ for $i = 3, \dots, n$. The rest of the proof coincides with the one of the multiplicity lemma 3.7. \square

6.5. Algorithm. Consider a smooth (real analytic, holomorphic) curve of polynomials

$$P(t)(z) = z^n - a_1(t)z^{n-1} + a_2(t)z^{n-2} - \dots + (-1)^n a_n(t)$$

with complex coefficients. The algorithm has the following steps:

- (1) If all roots of $P(0)$ are pairwise different, P is smoothly (real analytically, holomorphically) solvable for t near 0 by 3.3.
- (2) If there are distinct roots at $t = 0$ we put them into two subsets which factors $P(t) = P_1(t).P_2(t)$ by the splitting lemma 3.4. We then feed $P_i(t)$ (which have lower degree) into the algorithm.
- (3) All roots of $P(0)$ are equal. We first reduce $P(t)$ to the case $a_1(t) = 0$ by replacing the variable x by $y = x - a_1(t)/n$. Then all roots are equal to 0 so $a_i(0) = 0$ for all i .

If there does not exist an integer $r > 0$ with $m(a_i) \geq ir$ for $2 \leq i \leq n$, then by 3.5 the polynomial is *not* smoothly (real analytically, holomorphically)

solvable, by proposition 3.5: We store the polynomial as an output of the procedure, as a factor of $P^{(n)}$ below.

If there exists an integer $r > 0$ with $m(a_i) \geq ir$ for $2 \leq i \leq n$, let $a_i(t) = a_{i,ir}(t)t^{ir}$ ($i = 2, \dots, n$) for smooth (real analytic, holomorphic) $a_{i,ir}$. Consider the following smooth (real analytic, holomorphic) curve of polynomials

$$P_r(t)(x) = x^n + a_{2,2r}(t)x^{n-2} - a_{3,3r}(t)x^{n-3} \cdots + (-1)^n a_{n,nr}(t).$$

If $P_r(t)$ is smoothly (real analytically, holomorphically) solvable and $x_k(t)$ are its smooth (real analytic, holomorphic) roots, then $x_k(t)t^r$ are the roots of $P(t)$ and the original curve P is smoothly (real analytically, holomorphically) solvable too.

If for one coefficient we have $m(a_i) = ir$ then $P_r(0)$ has a coefficient which does not vanish, so not all roots of $P_r(0)$ are equal, and we may feed P_r into step (2).

If all coefficients of $P_r(0)$ are zero, we feed P_r again into step (3).

In the smooth case all $m(a_i)$ may be infinite; In this case we store the polynomial as a factor of $P^{(\infty)}$ below.

In the holomorphic and real analytic cases the algorithm provides a splitting of the polynomial $P(t) = P^{(n)}(t)P^{(s)}(t)$ into holomorphic and real analytic curves, where $P^{(s)}(t)$ is solvable, and where $P^{(n)}(t)$ is not solvable. But it may contain solvable roots, as is seen by simple examples.

In the smooth case the algorithm provides a splitting near $t = 0$

$$P(t) = P^{(\infty)}(t)P^{(n)}(t)P^{(s)}(t)$$

into smooth curves of polynomials, where: $P^{(\infty)}$ has the property that each root meets another one of infinite order at $t = 0$; and where $P^{(s)}(t)$ is smoothly solvable, and no two roots meet of infinite order at $t = 0$; $P^{(n)}$ is not smoothly solvable, with the same property as above.

6.6. Remarks. If $P(t)$ is a polynomials whose coefficients are meromorphic functions of a complex variable t , there is a well developed theory of the roots of $P(t)(x) = 0$ as multi-valued meromorphic functions, given by *Puiseux* or *Laurent-Puiseux* series. But it is difficult to extract holomorphic information out of it, and the algorithm above complements this theory. See for example Theorem 3 on page 370 (Anhang, §5) of [1]. The question of choosing roots continuously has been treated in [2]: one finds sufficient conditions for it.

7. CHOOSING EIGENVALUES AND EIGENVECTORS OF MATRICES SMOOTHLY

In this section we consider the following situation: Let $A(t) = (A_{ij}(t))$ be a smooth (real analytic, holomorphic) curve of real (complex) $(n \times n)$ -matrices or operators, depending on a real (complex) parameter t near 0. What can we say about the eigenvalues and eigenfunctions of $A(t)$?

Let us first recall some known results. These have some difficulty with the interpretation of the eigenprojections and the eigennilpotents at branch points of the eigenvalues, see [7], II, 1.11.

7.1. Result. ([7], II, 1.8) *Let $\mathbb{C} \ni t \mapsto A(t)$ be a holomorphic curve. Then all eigenvalues, all eigenprojections and all eigennilpotents are holomorphic with at most algebraic singularities at discrete points.*

7.2. Result. ([11], Satz 1) *Let $t \mapsto A(t)$ be a real analytic curve of hermitian complex matrices. Let λ be a k -fold eigenvalue of $A(0)$ with k orthonormal eigenvectors v_i , and suppose that there is no other eigenvalue of A near λ . Then there are k real analytic eigenvalues $\lambda_i(t)$ through λ , and k orthonormal real analytic eigenvectors through the v_i , for t near 0.*

The condition that $A(t)$ is hermitian cannot be omitted. Consider the following example of real semisimple (not normal) matrices

$$A(t) := \begin{pmatrix} 2t + t^3 & t \\ -t & 0 \end{pmatrix},$$

$$\lambda_{\pm}(t) = t + \frac{t^2}{2} \pm t^2 \sqrt{1 + \frac{t^2}{4}}, \quad x_{\pm}(t) = \begin{pmatrix} 1 + \frac{t}{2} \pm t \sqrt{1 + \frac{t^2}{4}} \\ -1 \end{pmatrix},$$

where at $t = 0$ we do not get a base of eigenvectors.

7.3. Result. (Rellich [13], see also Kato [7], II, 6.8) *Let $A(t)$ be a C^1 -curve of symmetric matrices. Then the eigenvalues can be chosen C^1 in t , on the whole parameter interval.*

For an extension of this result to Hilbert space, under stronger assumptions, see 7.8, whose proof will need 7.3. This result is best possible for the degree of continuous differentiability, as is shown by the following example.

7.4 Example. Consider the symmetric matrix

$$A(t) = \begin{pmatrix} a(t) & b(t) \\ b(t) & -a(t) \end{pmatrix}$$

The characteristic polynomial of $A(t)$ is $\lambda^2 - (a(t)^2 + b(t)^2)$. We shall specify the entries a and b as smooth functions in such a way, that $a(t)^2 + b(t)^2$ does not admit a C^2 -square root.

Assume that $a(t)^2 + b(t)^2 = c(t)^2$ for a C^2 -function c . Then we may compute as follows:

$$\begin{aligned} c^2 &= a^2 + b^2 \\ cc' &= aa' + bb' \\ (c')^2 + cc'' &= (a')^2 + aa'' + (b')^2 + bb'' \\ c'' &= \frac{1}{c} \left((a')^2 + aa'' + (b')^2 + bb'' - (c')^2 \right) \\ &= \frac{1}{c} \left((a')^2 + aa'' + (b')^2 + bb'' - \frac{(aa' + bb')^2}{c^2} \right) \\ &= \frac{(ab' - ba')^2 + a^3 a'' + b^3 b'' + ab^2 a'' + a^2 bb''}{c^3} \end{aligned}$$

By $c^2 = a^2 + b^2$ we have

$$\left| \frac{a^3}{c^3} \right| \leq 1, \quad \left| \frac{b^3}{c^3} \right| \leq 1, \quad \left| \frac{ab^2}{c^3} \right| \leq \frac{1}{\sqrt{3}}, \quad \left| \frac{a^2b}{c^3} \right| \leq \frac{1}{\sqrt{3}}.$$

So for C^2 -functions a , b , and continuous c all these terms are bounded. We will now construct smooth a and b such that

$$\left(\frac{(ab' - ba')^2}{c^3} \right)^2 = \frac{(ab' - ba')^4}{(a^2 + b^2)^3}$$

is unbounded near $t = 0$. This contradicts that c is C^2 . For this we choose a and b similar to the function f in example 2.4 with the same t_n and h_n :

$$a(t) := \sum_{n=1}^{\infty} h_n(t - t_n) \cdot \left(\frac{2n}{2^n}(t - t_n) + \frac{1}{4^n} \right),$$

$$b(t) := \sum_{n=1}^{\infty} h_n(t - t_n) \cdot \left(\frac{2n}{2^n}(t - t_n) \right).$$

Then $a(t_n) = \frac{1}{4^n}$, $b(t_n) = 0$, $|c(t_n)| = \frac{1}{4^n}$, and $b'(t_n) = \frac{2n}{2^n}$.

7.5. Result. (Rellich [12], see also Kato [7], VII, 3.9) *Let $A(t)$ be a real analytic curve of unbounded self-adjoint operators in a Hilbert space with common domain of definition and with compact resolvent.*

Then the eigenvalues and the eigenvectors can be chosen real analytically in t , on the whole parameter domain.

7.6. Theorem. *Let $A(t) = (A_{ij}(t))$ be a smooth curve of complex hermitian $(n \times n)$ -matrices, depending on a real parameter $t \in \mathbb{R}$, acting on a hermitian space $V = \mathbb{C}^n$, such that no two of the continuous eigenvalues meet of infinite order at any $t \in \mathbb{R}$ if they are not equal for all t .*

Then all the eigenvalues and all the eigenvectors can be chosen smoothly in t , on the whole parameter domain \mathbb{R} .

The last condition permits that some eigenvalues agree for all t — we speak of higher ‘generic multiplicity’ in this situation.

Proof. The proof will use an algorithm.

Note first that by 4.2 the characteristic polynomial

$$(1) \quad \begin{aligned} P(A(t))(\lambda) &= \det(A(t) - \lambda \mathbb{I}) \\ &= \lambda^n - a_1(t)\lambda^{n-1} + a_2(t)\lambda^{n-2} - \dots + (-1)^n a_n(t) \\ &= \sum_{i=0}^n \text{Trace}(\Lambda^i A(t)) \lambda^{n-i} \end{aligned}$$

is smoothly solvable, with smooth roots $\lambda_1(t), \dots, \lambda_n(t)$, on the whole parameter interval.

Case 1: distinct eigenvalues. If $A(0)$ has some eigenvalues distinct, then one can reorder them in such a way that for $i_0 = 0 < 1 \leq i_1 < i_2 < \dots < i_k < n = i_{k+1}$ we have

$$\lambda_1(0) = \dots = \lambda_{i_1}(0) < \lambda_{i_1+1}(0) = \dots = \lambda_{i_2}(0) < \dots < \lambda_{i_k+1}(0) = \dots = \lambda_n(0)$$

For t near 0 we still have

$$\lambda_1(t), \dots, \lambda_{i_1}(t) < \lambda_{i_1+1}(t), \dots, \lambda_{i_2}(t) < \dots < \lambda_{i_k+1}(t), \dots, \lambda_n(t)$$

Now for $j = 1, \dots, k+1$ consider the subspaces

$$V_t^{(j)} = \bigoplus_{i=i_{j-1}+1}^{i_j} \{v \in V : (A(t) - \lambda_i(t))v = 0\}$$

Then each $V_t^{(j)}$ runs through a smooth vector subbundle of the trivial bundle $(-\varepsilon, \varepsilon) \times V \rightarrow (-\varepsilon, \varepsilon)$, which admits a smooth framing $e_{i_{j-1}+1}(t), \dots, e_{i_j}(t)$. We have $V = \bigoplus_{j=1}^{k+1} V_t^{(j)}$ for each t .

In order to prove this statement note that

$$V_t^{(j)} = \ker \left((A(t) - \lambda_{i_{j-1}+1}(t)) \circ \dots \circ (A(t) - \lambda_{i_j}(t)) \right)$$

so $V_t^{(j)}$ is the kernel of a smooth vector bundle homomorphism $B(t)$ of constant rank (even of constant dimension of the kernel), and thus is a smooth vector subbundle. This together with a smooth frame field can be shown as follows: Choose a basis of V such that $A(0)$ is diagonal. Then by the elimination procedure one can construct a basis for the kernel of $B(0)$. For t near 0, the elimination procedure (with the same choices) gives then a basis of the kernel of $B(t)$; the elements of this basis are then smooth in t , for t near 0.

From the last result it follows that it suffices to find smooth eigenvectors in each subbundle $V^{(j)}$ separately, expanded in the smooth frame field. But in this frame field the vector subbundle looks again like a constant vector space. So feed each of this parts (A restricted to $V^{(j)}$, as matrix with respect to the frame field) into case 2 below.

Case 2: All eigenvalues at 0 are equal. So suppose that $A(t) : V \rightarrow V$ is hermitian with all eigenvalues at $t = 0$ equal to $\frac{a_1(0)}{n}$, see (1).

Eigenvectors of $A(t)$ are also eigenvectors of $A(t) - \frac{a_1(t)}{n}\mathbb{I}$, so we may replace $A(t)$ by $A(t) - \frac{a_1(t)}{n}\mathbb{I}$ and assume that for the characteristic polynomial (1) we have $a_1 = 0$, or assume without loss that $\lambda_i(0) = 0$ for all i and so $A(0) = 0$.

If $A(t) = 0$ for all t we choose the eigenvectors constant.

Otherwise let $A_{ij}(t) = tA_{ij}^{(1)}(t)$. From (1) we see that the characteristic polynomial of the hermitian matrix $A^{(1)}(t)$ is $P_1(t)$ in the notation of 3.8, thus $m(a_i) \geq i$ for $2 \leq i \leq n$ (which follows from 3.5 also).

The eigenvalues of $A^{(1)}(t)$ are the roots of $P_1(t)$, which may be chosen in a smooth way, since they again satisfy the condition of theorem 4.2. Note that eigenvectors

of $A^{(1)}$ are also eigenvectors of A . If the eigenvalues are still all equal, we apply the same procedure again, until they are not all equal: we arrive at this situation by the assumption of the theorem. Then we apply case 1.

This algorithm shows that one may choose the eigenvectors $x_i(t)$ of $A(t)$ in a smooth way, locally in t . It remains to extend this to the whole parameter interval.

If some eigenvalues coincide locally, then on the whole of \mathbb{R} , by the assumption. The corresponding eigenspaces then form a smooth vector bundle over \mathbb{R} , by case 1, since those eigenvalues, which meet in isolated points are different after application of case 2.

So we we get $V = \bigoplus W_t^{(j)}$ where each $W_t^{(j)}$ is a smooth sub vector bundles of $V \times \mathbb{R}$, whose dimension is the generic multiplicity of the corresponding smooth eigenvalue function. It suffices to find global orthonormal smooth frames for each of these vector bundles; this exists since the vector bundle is smoothly trivial, by using parallel transport with respect to a smooth Hermitian connection. \square

7.7. Example. (see [11], §2) That the last result cannot be improved is shown by the following example which rotates a lot:

$$\begin{aligned} x_+(t) &:= \begin{pmatrix} \cos \frac{1}{t} \\ \sin \frac{1}{t} \end{pmatrix}, & x_-(t) &:= \begin{pmatrix} -\sin \frac{1}{t} \\ \cos \frac{1}{t} \end{pmatrix}, & \lambda_{\pm}(t) &= \pm e^{-\frac{1}{t^2}}, \\ A(t) &:= (x_+(t), x_-(t)) \begin{pmatrix} \lambda_+(t) & 0 \\ 0 & \lambda_-(t) \end{pmatrix} (x_+(t), x_-(t))^{-1} \\ &= e^{-\frac{1}{t^2}} \begin{pmatrix} \cos \frac{2}{t} & \sin \frac{2}{t} \\ \sin \frac{2}{t} & -\cos \frac{2}{t} \end{pmatrix}. \end{aligned}$$

Here $t \mapsto A(t)$ and $t \mapsto \lambda_{\pm}(t)$ are smooth, whereas the eigenvectors cannot be chosen continuously.

7.8. Theorem. *Let $t \mapsto A(t)$ be a smooth curve of unbounded self-adjoint operators in a Hilbert space with common domain of definition and with compact resolvent. Then the eigenvalues of $A(t)$ may be arranged in such a way that each eigenvalue is C^1 .*

Suppose moreover that no two of the continuously chosen eigenvalues meet of infinite order at any $t \in \mathbb{R}$ if they are not equal. Then the eigenvalues and the eigenvectors can be chosen smoothly in t , on the whole parameter domain.

Remarks. That $A(t)$ is a smooth curve of unbounded operators means the following: There is a dense subspace V of the Hilbert space H such that V is the domain of definition of each $A(t)$, and such that $A(t)^* = A(t)$ with the same domains V , where the adjoint operator $A(t)^*$ is defined by $\langle A(t)u, v \rangle = \langle u, A(t)^*v \rangle$ for all v for which the left hand side is bounded as function in $u \in H$. Moreover we require that $t \mapsto \langle A(t)u, v \rangle$ is smooth for each $u \in V$ and $v \in H$. This implies that $t \mapsto A(t)u$ is smooth $\mathbb{R} \rightarrow H$ for each $u \in V$ by [9], 2.3 or [5], 2.6.2.

The first part of the proof will show that $t \mapsto A(t)$ smooth implies that the resolvent $(A(t) - z)^{-1}$ is smooth in t and z jointly, and only this is used later in the proof.

It is well known and in the proof we will show that if for some (t, z) the resolvent $(A(t) - z)^{-1}$ is compact then for all $t \in \mathbb{R}$ and z in the resolvent set of $A(t)$.

Proof. For each t consider the norm $\|u\|_t^2 := \|u\|^2 + \|A(t)u\|^2$ on V . Since $A(t) = A(t)^*$ is closed, $(V, \|\cdot\|_t)$ is again a Hilbert space with inner product $\langle u, v \rangle_t := \langle u, v \rangle + \langle A(t)u, A(t)v \rangle$. All these norms are equivalent since $(V, \|\cdot\|_t + \|\cdot\|_s) \rightarrow (V, \|\cdot\|_t)$ is continuous and bijective, so an isomorphism by the open mapping theorem. Then $t \mapsto \langle u, v \rangle_t$ is smooth for fixed $u, v \in V$, and by the multilinear uniform boundedness principle ([9], 5.17 or [5], 3.7.4 + 4.1.19) the mapping $t \mapsto \langle \cdot, \cdot \rangle_t$ is smooth into the space of bounded bilinear forms on $(V, \|\cdot\|_s)$ for each fixed s . By the exponential law ([9], 3.12 or [5], 1.4.3) $(t, u) \mapsto \|u\|_t^2$ is smooth from $\mathbb{R} \times (V, \|\cdot\|_s) \rightarrow \mathbb{R}$ for each fixed s . Thus all Hilbert norms $\|\cdot\|_t$ are locally uniformly equivalent, since $\{\|u\|_t : |t| \leq K, \|u\|_s \leq 1\}$ is bounded by $L_{K,s}$ in \mathbb{R} , so $\|u\|_t \leq L_{K,s}\|u\|_s$ for all $|t| \leq K$. Let us now equip V with one of the equivalent Hilbert norms, say $\|\cdot\|_0$. Then each $A(t)$ is a globally defined operator $V \rightarrow H$ with closed graph and is thus bounded, and by using again the (multi)linear uniform boundedness theorem as above we see that $t \mapsto A(t)$ is smooth $\mathbb{R} \rightarrow L(V, H)$.

If for some $(t, z) \in \mathbb{R} \times \mathbb{C}$ the bounded operator $A(t) - z : V \rightarrow H$ is invertible, then this is true locally and $(t, z) \mapsto (A(t) - z)^{-1} : H \rightarrow V$ is smooth since inversion is smooth on Banach spaces.

Since each $A(t)$ is hermitian the *global resolvent set* $\{(t, z) \in \mathbb{R} \times \mathbb{C} : (A(t) - z) : V \rightarrow H \text{ is invertible}\}$ is open, contains $\mathbb{R} \times (\mathbb{C} \setminus \mathbb{R})$, and hence is connected.

Moreover $(A(t) - z)^{-1} : H \rightarrow H$ is a compact operator for some (equivalently any) (t, z) if and only if the inclusion $i : V \rightarrow H$ is compact, since $i = (A(t) - z)^{-1} \circ (A(t) - z) : V \rightarrow H \rightarrow H$.

Let us fix a parameter s . We choose a simple smooth curve γ in the resolvent set of $A(s)$ for fixed s .

- (1) *Claim* For t near s , there are C^1 -functions $t \mapsto \lambda_i(t) : 1 \leq i \leq N$ which parametrize all eigenvalues (repeated according to their multiplicity) of $A(t)$ in the interior of γ . If no two of the generically different eigenvalues meet of infinite order they can be chosen smoothly.

By replacing $A(s)$ by $A(s) - z_0$ if necessary we may assume that 0 is not an eigenvalue of $A(s)$. Since the global resolvent set is open, no eigenvalue of $A(t)$ lies on γ or equals 0, for t near s . Since

$$t \mapsto -\frac{1}{2\pi i} \int_{\gamma} (A(t) - z)^{-1} dz =: P(t, \gamma)$$

is a smooth curve of projections (on the direct sum of all eigenspaces corresponding to eigenvalues in the interior of γ) with finite dimensional ranges, the ranks (i.e. dimension of the ranges) must be constant: it is easy to see that the (finite) rank cannot fall locally, and it cannot increase, since the distance in $L(H, H)$ of $P(t)$ to the subset of operators of rank $\leq N = \text{rank}(P(s))$ is continuous in t and is either 0 or 1. So for t near s , there are equally many eigenvalues in the interior, and we may call them $\mu_i(t) : 1 \leq i \leq N$ (repeated with multiplicity). Let us denote by $e_i(t) : 1 \leq i \leq N$ a corresponding system of eigenvectors of $A(t)$. Then by the residue theorem we have

$$\sum_{i=1}^N \mu_i(t)^p e_i(t) \langle e_i(t), \cdot \rangle = -\frac{1}{2\pi i} \int_{\gamma} z^p (A(t) - z)^{-1} dz$$

which is smooth in t near s , as a curve of operators in $L(H, H)$ of rank N , since 0 is not an eigenvalue.

- (2) *Claim.* Let $t \mapsto T(t) \in L(H, H)$ be a smooth curve of operators of rank N in Hilbert space such that $T(0)T(0)(H) = T(0)(H)$. Then $t \mapsto \text{Trace}(T(t))$ is smooth near 0 (note that this implies T smooth into the space of nuclear operators, since all bounded linear functionals are of the form $A \mapsto \text{Trace}(AB)$, by [9], 2.3 or 2.14.(4).

Let $F := T(0)(H)$. Then $T(t) = (T_1(t), T_2(t)) : H \rightarrow F \oplus F^\perp$ and the image of $T(t)$ is the space

$$\begin{aligned} T(t)(H) &= \{(T_1(t)(x), T_2(t)(x)) : x \in H\} \\ &= \{(T_1(t)(x), T_2(t)(x)) : x \in F\} \text{ for } t \text{ near } 0 \\ &= \{(y, S(t)(y)) : y \in F\}, \text{ where } S(t) := T_2(t) \circ (T_1(t)|_F)^{-1}. \end{aligned}$$

Note that $S(t) : F \rightarrow F^\perp$ is smooth in t by finite dimensional inversion for $T_1(t)|_F : F \rightarrow F$. Now

$$\begin{aligned} \text{Trace}(T(t)) &= \text{Trace} \left(\begin{pmatrix} 1 & 0 \\ -S(t) & 1 \end{pmatrix} \begin{pmatrix} T_1(t)|_F & T_1(t)|_{F^\perp} \\ T_2(t)|_F & T_2(t)|_{F^\perp} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ S(t) & 1 \end{pmatrix} \right) \\ &= \text{Trace} \left(\begin{pmatrix} T_1(t)|_F & T_1(t)|_{F^\perp} \\ 0 & -S(t)T_1(t)|_{F^\perp} + T_2(t)|_{F^\perp} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ S(t) & 1 \end{pmatrix} \right) \\ &= \text{Trace} \left(\begin{pmatrix} T_1(t)|_F & T_1(t)|_{F^\perp} \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ S(t) & 1 \end{pmatrix} \right), \text{ since rank} = N \\ &= \text{Trace} \begin{pmatrix} T_1(t)|_F + (T_1(t)|_{F^\perp})S(t) & T_1(t)|_{F^\perp} \\ 0 & 0 \end{pmatrix} \\ &= \text{Trace} \left(T_1(t)|_F + (T_1(t)|_{F^\perp})S(t) : F \rightarrow F \right), \end{aligned}$$

which is visibly smooth since F is finite dimensional.

From claim (2) we now may conclude that

$$\sum_{i=1}^N \mu_i(t)^p = -\frac{1}{2\pi i} \text{Trace} \int_{\gamma} z^p (A(t) - z)^{-1} dz$$

is smooth for t near s .

Thus the Newton polynomial mapping $s^N(\mu_1(t), \dots, \mu_N(t))$ is smooth, so also the elementary symmetric polynomial $\sigma^N(\mu_1(t), \dots, \mu_N(t))$ is smooth, and thus $\{\mu_i(t) : 1 \leq i \leq N\}$ is the set of roots of a polynomial with smooth coefficients. By theorem 4.3 there is an arrangement of these roots such that they become differentiable. If no two of the generically different ones meet of infinite order, by theorem 4.2 there is even a smooth arrangement.

To see that in the general case they are even C^1 note that the images of the projections $P(t, \gamma)$ of constant rank for t near s describe the fibers of a smooth vector bundle. The restriction of $A(t)$ to this bundle, viewed in a smooth framing,

becomes a smooth curve of symmetric matrices, for which by Rellich's result 7.3 the eigenvalues can be chosen C^1 . This finishes the proof of claim (1).

(3) *Claim.* Let $t \mapsto \lambda_i(t)$ be a differentiable eigenvalue of $A(t)$, defined on some interval. Then

$$|\lambda_i(t_1) - \lambda_i(t_2)| \leq (1 + |\lambda_i(t_2)|)(e^{a|t_1 - t_2|} - 1)$$

holds for a continuous positive function $a = a(t_1, t_2)$ which is independent of the choice of the eigenvalue.

For fixed t near s take all roots λ_j which meet λ_i at t , order them differentiably near t , and consider the projector $P(t, \gamma)$ onto the joint eigenspaces for only those roots (where γ is a simple smooth curve containing only $\lambda_i(t)$ in its interior, of all the eigenvalues at t). Then the image of $u \mapsto P(u, \gamma)$, for u near t , describes a smooth finite dimensional vector subbundle of $\mathbb{R} \times H$, since its rank is constant. For each u choose an orthonormal system of eigenvectors $v_j(u)$ of $A(u)$ corresponding to these $\lambda_j(u)$. They form a (not necessarily continuous) framing of this bundle. For any sequence $t_k \rightarrow t$ there is a subsequence such that each $v_j(t_k) \rightarrow w_j(t)$ where $w_j(t)$ is again an orthonormal system of eigenvectors of $A(t)$ for the eigenspace of $\lambda_i(t)$. Now consider

$$\frac{A(t) - \lambda_i(t)}{t_k - t} v_i(t_k) + \frac{A(t_k) - A(t)}{t_k - t} v_i(t_k) - \frac{\lambda_i(t_k) - \lambda_i(t)}{t_k - t} v_i(t_k) = 0,$$

take the inner product of this with $w_i(t)$, note that then the first summand vanishes, and let $t_k \rightarrow t$ to obtain

$$\lambda_i'(t) = \langle A'(t)w_i(t), w_i(t) \rangle \text{ for an eigenvector } w_i(t) \text{ of } A(t) \text{ with eigenvalue } \lambda_i(t).$$

This implies, where $V_t = (V, \|\cdot\|_t)$,

$$\begin{aligned} |\lambda_i'(t)| &\leq \|A'(t)\|_{L(V_t, H)} \|w_i(t)\|_{V_t} \|w_i(t)\|_H \\ &= \|A'(t)\|_{L(V_t, H)} \sqrt{\|w_i(t)\|_H^2 + \|A(t)w_i(t)\|_H^2} \\ &= \|A'(t)\|_{L(V_t, H)} \sqrt{1 + \lambda_i(t)^2} \leq a + a|\lambda_i(t)|, \end{aligned}$$

for a constant a which is valid for a compact interval of t 's since $t \mapsto \|\cdot\|_t^2$ is smooth on V . By Gronwall's lemma (see e.g. [3], (10.5.1.3)) this implies claim (3).

By the following arguments we can conclude that all eigenvalues may be numbered as $\lambda_i(t)$ for i in \mathbb{N} or \mathbb{Z} in such a way that they are differentiable (by which we mean C^1 , or C^∞ under the stronger assumption) in $t \in \mathbb{R}$. Note first that by claim (3) no eigenvalue can go off to infinity in finite time since it may increase at most exponentially. Let us first number all eigenvalues of $A(0)$ increasingly.

We claim that for one eigenvalue (say $\lambda_0(0)$) there exists a differentiable extension to all of \mathbb{R} ; namely the set of all $t \in \mathbb{R}$ with a differentiable extension of λ_0 on the segment from 0 to t is open and closed. Open follows from claim (1). If this interval does not reach infinity, from claim (3) it follows that $(t, \lambda_0(t))$ has an accumulation point (s, x) at the the end s . Clearly x is an eigenvalue of $A(s)$, and by claim (1)

the eigenvalues passing through (s, x) can be arranged differentiably, and thus $\lambda_0(t)$ converges to x and can be extended differentiably beyond s .

By the same argument we can extend iteratively all eigenvalues differentiably to all $t \in \mathbb{R}$: if it meets an already chosen one, the proof of 4.3 shows that we may pass through it coherently.

Now we start to choose the eigenvectors smoothly, under the stronger assumption. Let us consider again eigenvalues $\{\lambda_i(t) : 1 \leq i \leq N\}$ contained in the interior of a smooth curve γ for t in an open interval I . Then $V_t := P(t, \gamma)(H)$ is the fiber of a smooth vector bundle of dimension N over I . We choose a smooth framing of this bundle, and use then the proof of theorem 7.6 to choose smooth sub vector bundles whose fibers over t are the eigenspaces of the eigenvalues with their generic multiplicity. By the same arguments as in 7.6 we then get global vector sub bundles with fibers the eigenspaces of the eigenvalues with their generic multiplicity, and finally smooth eigenvectors for all eigenvalues. \square

REFERENCES

1. Baumgärtel, Hellmut, *Endlichdimensionale analytische Störungstheorie*, Akademie-Verlag, Berlin, 1972.
2. Burghilea, Ana, *On the numbering of roots of a family of generic polynomials*, Bul. Inst. Politehn. Bucuresti, Ser. Mec. **39**, no. 3 (1977), 11–15.
3. Dieudonné, J. A., *Foundations of modern analysis, I*, Academic Press, New York – London, 1960.
4. Dieudonné, J., *Sur un theoreme de Glaeser*, J. Anal. Math. **23** (1970), 85–88.
5. Frölicher, Alfred; Kriegl, Andreas, *Linear spaces and differentiation theory*, Pure and Applied Mathematics, J. Wiley, Chichester, 1988.
6. Glaeser, G., *Racine carré d'une fonction différentiable*, Ann. Inst. Fourier (Grenoble) **13,2** (1963), 203–210.
7. Kato, Tosio, *Perturbation theory for linear operators*, Grundlehren 132, Springer-Verlag, Berlin, 1976.
8. Kriegl, Andreas; Michor, Peter W., *A convenient setting for real analytic mappings*, Acta Mathematica **165** (1990), 105–159.
9. Kriegl, A.; Michor, Peter W., *The Convenient Setting of Global Analysis*, to appear in 'Surveys and Monographs', AMS, Providence.
10. Procesi, Claudio, *Positive symmetric functions*, Adv. Math. **29** (1978), 219–225.
11. Rellich, F., *Störungstheorie der Spektralzerlegung, I*, Math. Ann. **113** (1937), 600–619.
12. Rellich, F., *Störungstheorie der Spektralzerlegung, V*, Math. Ann. **118** (1940), 462–484.
13. Rellich, F., *Perturbation theory of eigenvalue problems*, Lecture Notes, New York University, 1953; Gordon and Breach, New York, London, Paris, 1969.
14. Sylvester, J., *On a theory of the syzygetic relations of two rational integral functions, comprising an application to the theory of Sturm's functions, and that of the greatest algebraic common measure*, Philosoph. Trans. Royal Soc. London **CXLIII**, part III (1853), 407–548; *Mathematical papers*, Vol. I, At the University Press, Cambridge, 1904, pp. 511ff.
15. Warner, Frank, *Personal communication*.

D. V. ALEKSEEVSKY: CENTER 'SOPHUS LIE', KRASNOKAZARMENNAYA 6, 111250 MOSCOW, RUSSIA

A. KRIEGL, P. W. MICHOR: INSTITUT FÜR MATHEMATIK, UNIVERSITÄT WIEN, STRUDLHOF-GASSE 4, A-1090 WIEN, AUSTRIA

E-mail address: `Andreas.Kriegl@univie.ac.at`, `Peter.Michor@esi.ac.at`

M. LOSIK: SARATOV STATE UNIVERSITY, UL. ASTRAKHANSKAYA, 83, 410026 SARATOV, RUSSIA

E-mail address: `losik@scnit.saratov.su`