

Δ-matroids and Pfaffian Forms

Walter Wenzel, Bielefeld

In analogy to the fact that representable matroids may be defined in terms of Grassmann-Plücker maps we want to describe representability of Δ-matroids by skew-symmetric matrices in terms of Pfaffian forms. In the sequel we assume $n \in \mathbb{N}$, $S := \{1, \dots, n\}$, and K denotes some commutative field.

Definition 1 (cf. [B1, §6] or [B2, §1]):

Assume $\mathcal{F} \subseteq \mathcal{P}(S)$. The pair (S, \mathcal{F}) is a **Δ-matroid**, if \mathcal{F} satisfies

$$(SEA) \quad \text{For } F_1, F_2 \in \mathcal{F} \text{ and } e \in F_1 \Delta F_2 \text{ there exists some } f \in F_1 \Delta F_2 \text{ with } F_1 \Delta \{e, f\} \in \mathcal{F}.$$

For $\mathcal{F} \subseteq \mathcal{P}(S)$ and $T \subseteq S$ we put $\mathcal{F} \Delta T := \{F \Delta T \mid F \in \mathcal{F}\}$.

Definition 2 (cf. [B2, §4]):

A Δ-matroid (S, \mathcal{F}) is representable over K by a skew-symmetric matrix $A = (a_{ij})_{i,j \in S}$, if for some $T \subseteq S$ we have

$$\mathcal{F} \Delta T = \mathcal{F}(A) := \{S' \subseteq S \mid A' := (a_{ij})_{i,j \in S'} \text{ is nonsingular}\},$$

where $(a_{ij})_{i,j \in \emptyset}$ is considered to be nonsingular. A is then called a **presentation** of (S, \mathcal{F}) .

Theorem 3: A Δ-matroid (S, \mathcal{F}) is representable over K by a skew-symmetric matrix if and only if there exists some map $P : \mathcal{P}(S) \rightarrow K$ and some $T \in \mathcal{F}$ such that

(P0) For $I \subseteq S$ we have $P(I) \neq 0$ if and only if $I \Delta T \in \mathcal{F}$.

(P1) If $I \subseteq S$ and $\#I \equiv 1 \pmod{2}$, then $P(I) = 0$.

(P2) If $I_1, I_2 \subseteq S$ and $I_1 \Delta I_2 = \{i_1, \dots, i_k\}$ with $i_j < i_{j+1}$ for $1 \leq j \leq k-1$, then

$$\sum_{j=1}^k (-1)^j \cdot P(I_1 \Delta \{i_j\}) \cdot P(I_2 \Delta \{i_j\}) = 0.$$

If (S, \mathcal{F}) is representable by $A = (a_{ij})_{i,j \in S}$, then P may be chosen to be the corresponding Pfaffian form, which may be defined by $P(I) := 0$ for $\#I \equiv 1 \pmod{2}$, $P(\emptyset) := 1$, $P(\{i, j\}) := a_{ij}$ for $1 \leq i < j \leq n$, and

$$P(\{i_1, \dots, i_{2m}\}) := \sum_{j=2}^{2m} (-1)^j \cdot P(\{i_1, i_j\}) \cdot P(\{i_2, \dots, i_{2m}\} \setminus \{i_j\})$$

for $2 \leq m \leq \frac{n}{2}$ and $i_j < i_{j+1}$ for $1 \leq j \leq 2m-1$.

Vice versa, if P satisfies (P0), (P1), (P2), then (S, \mathcal{F}) is representable by the matrix $A = (a_{ij})_{i,j \in S}$ given by $a_{ii} := 0$ for $1 \leq i \leq n$, $a_{ij} = -a_{ji} = P(\{i, j\})$ for $1 \leq i < j \leq n$.

Corollary: A Δ -matroid (S, \mathcal{F}) , representable by some skew-symmetric matrix, satisfies the following strong exchange property:

$$\text{For } F_1, F_2 \in \mathcal{F} \text{ and } e \in F_1 \Delta F_2 \text{ there exists some} \\ f \in (F_1 \Delta F_2) \setminus \{e\} \text{ with } F_1 \Delta \{e, f\} \in \mathcal{F} \text{ and } F_2 \Delta \{e, f\} \in \mathcal{F}.$$

Furthermore, Theorem 3 suggests

Definition 4: Assume $F \subseteq \mathcal{P}(S)$.

- i) (S, \mathcal{F}) is an **orientable Δ -matroid**, if there exists some $T \in \mathcal{F}$ and some map $P : \mathcal{P}(S) \rightarrow \{0, 1, -1\}$ satisfying
 - (OP0) For $I \subseteq S$ we have $P(I) \in \{1, -1\}$ if and only if $I \Delta T \in \mathcal{F}$.
 - (OP1) For $I \subseteq S$ with $\#I \equiv 1 \pmod{2}$ we have $P(I) = 0$.
 - (OP2) If $I_1, I_2 \subseteq S$, $I_1 \Delta I_2 = \{i_1, \dots, i_k\}$ with $i_j < i_{j+1}$ for $1 \leq j \leq k-1$, and if for some $w \in \{1, -1\}$ we have $\kappa_j := w \cdot (-1)^j \cdot P(I_1 \Delta \{i_j\}) \cdot P(I_2 \Delta \{i_j\}) \geq 0$ for $1 \leq j \leq k$, then $\kappa_j = 0$ for every j .
- ii) Assume $P : \mathcal{P}(S) \rightarrow \mathbb{R}^+ \cup \{0\}$ is some map. (S, \mathcal{F}, P) is a **valuated Δ -matroid**, if for some $T \in \mathcal{F}$ we have
 - (VP0) For $I \subseteq S$ we have $P(I) > 0$ if and only if $I \Delta T \in \mathcal{F}$.
 - (VP1) For $I \subseteq S$ with $\#I \equiv 1 \pmod{2}$ we have $P(I) = 0$.
 - (VP2) If $I_1, I_2 \subseteq S$ with $\#I_1 \equiv \#I_2 \equiv 0 \pmod{2}$, then for every $i \in I_1 \Delta I_2$ there exists some $j \in (I_1 \Delta I_2) \setminus \{i\}$ with

$$P(I_1) \cdot P(I_2) \leq P(I_1 \Delta \{i, j\}) \cdot P(I_2 \Delta \{i, j\}).$$

References

- [B1] A. Bouchet: Greedy algorithm and symmetric matroids, *Mathematical Programming* 38 (1987), 147-159.
- [B2] A. Bouchet: Representability of Δ -matroids, *Colloquia Mathematica Societatis János Bolyai* 52. Combinatorics, Eger (Hungary), 1987, 167-182.
- [DW] A.W.M. Dress and W. Wenzel: Grassmann-Plücker relations and matroids with coefficients, to appear in *Advances in Mathematics*.
- [K] G. Kowalewski: *Determinantentheorie*, Chelsea Publishing Company, New York, 1948.
- [S] J.R. Stembridge: Nonintersecting Paths, Pfaffians, and Plane Partitions, *Advances in Mathematics* 83 (1990), 96-131.
- [W] W. Wenzel: Pfaffian Forms and Δ -Matroids, in preparation.