

**A CONTINUED FRACTION EXPANSION FOR A q -TANGENT
FUNCTION:
AN ELEMENTARY PROOF**

HELMUT PRODINGER

ABSTRACT. We prove a continued fraction expansion for a certain q -tangent function that was conjectured by the present writer, then proved by Fulmek, now in a completely elementary way.

1. INTRODUCTION

In [3], the present writer defined the following q -trigonometric functions

$$\begin{aligned} \sin_q(z) &= \sum_{n \geq 0} \frac{(-1)^n z^{2n+1}}{[2n+1]_q!} q^{n^2}, \\ \cos_q(z) &= \sum_{n \geq 0} \frac{(-1)^n z^{2n}}{[2n]_q!} q^{n^2}. \end{aligned}$$

Here, we use standard q -notation:

$$[n]_q := \frac{1 - q^n}{1 - q}, \quad [n]_q! := [1]_q [2]_q \cdots [n]_q.$$

These q -functions are variations of Jackson's [2] q -sine and q -cosine functions.

For the q -tangent function $\tan_q(z) = \frac{\sin_q(z)}{\cos_q(z)}$, the following continued fraction expansion was conjectured in [3]:

$$z \tan_q(z) = \frac{z^2}{[1]_q q^0 - \frac{z^2}{[3]_q q^{-2} - \frac{z^2}{[5]_q q^1 - \frac{z^2}{[7]_q q^{-9} - \dots}}}}.$$

Here, the powers of q are of the form $(-1)^{n-1} n(n-1)/2 - n + 1$.

In [1], this statement was proven using heavy machinery from q -analysis.

Happily, after about 8 years, I was now successful to provide a complete *elementary* proof that I will present in the next section.

Key words and phrases. q -tangent, continued fraction.

2. THE PROOF

We write

$$\frac{z \sin_q(z)}{\cos_q(z)} = \frac{z^2}{N_0} = \frac{z^2}{C_1 - \frac{z^2}{N_1}} = \frac{z^2}{C_1 - \frac{z^2}{C_2 - \frac{z^2}{N_2}}} = \dots,$$

and set

$$N_i = \frac{a_i}{b_i}.$$

This means that

$$N_i = C_{i+1} - \frac{z^2}{N_{i+1}}$$

or

$$\frac{z^2}{N_{i+1}} = C_{i+1} - N_i$$

and

$$\frac{b_{i+1}z^2}{a_{i+1}} = C_{i+1} - \frac{a_i}{b_i} = \frac{C_{i+1}b_i - a_i}{b_i}.$$

Therefore we may identify numerators and denominators, and put $a_i = b_{i-1}$ and

$$b_{i+1}z^2 = C_{i+1}b_i - b_{i-1}.$$

The initial conditions are

$$b_{-1} = \cos_q(z) \quad \text{and} \quad b_0 = \sum_{n \geq 0} \frac{(-1)^n q^{n^2} z^{2n}}{[2n+1]_q!}.$$

The constants C_i guarantee that all the b_i are power series, i.e., they make the constant term in $C_{i+1}b_i - b_{i-1}$ disappear. Our goal is to show that $C_i = [2i-1]_q q^{(-1)^{i-1}i(i-1)/2-i+1}$ are the (unique) numbers that do this. We are proving the claim by proving the following *explicit* formula for b_i :

$$b_i = \sum_{n \geq 0} \frac{(-1)^n z^{2n}}{[2n+2i+1]_q!} \left(\prod_{j=1}^i [2n+2j]_q \right) q^{(n+\lfloor \frac{i+1}{2} \rfloor)^2 + [i \text{ odd}] \binom{i+1}{2}}.$$

Note that the C_i are uniquely determined by the imposed condition, and since the b_i are power series, we are done once we prove this formula by induction. The first two instances satisfy this, and we do the induction step now:

$$\begin{aligned} & C_{i+1}b_i - b_{i-1} \\ &= [2i+1]_q q^{(-1)^i \binom{i+1}{2} - i} \sum_{n \geq 0} \frac{(-1)^n z^{2n}}{[2n+2i+1]_q!} \left(\prod_{j=1}^i [2n+2j]_q \right) q^{(n+\lfloor \frac{i+1}{2} \rfloor)^2 + [i \text{ odd}] \binom{i+1}{2}} \\ &\quad - \sum_{n \geq 0} \frac{(-1)^n z^{2n}}{[2n+2i-1]_q!} \left(\prod_{j=1}^{i-1} [2n+2j]_q \right) q^{(n+\lfloor \frac{i}{2} \rfloor)^2 + [i-1 \text{ odd}] \binom{i}{2}} \\ &= \sum_{n \geq 0} \frac{(-1)^n z^{2n}}{[2n+2i+1]_q!} \left([2i+1]_q \left(\prod_{j=1}^i [2n+2j]_q \right) q^{(n+\lfloor \frac{i+1}{2} \rfloor)^2 + [i \text{ odd}] \binom{i+1}{2}} + (-1)^i \binom{i+1}{2} - i \right) \end{aligned}$$

$$\begin{aligned}
& - [2n + 2i + 1]_q \left(\prod_{j=1}^i [2n + 2j]_q \right) q^{(n + \lfloor \frac{i}{2} \rfloor)^2 + [i-1 \text{ odd}] \binom{i}{2}} \\
&= \frac{1}{1-q} \sum_{n \geq 0} \frac{(-1)^n z^{2n}}{[2n + 2i + 1]_q!} \left(\prod_{j=1}^i [2n + 2j]_q \right) \\
& \quad \times \left((1 - q^{2i+1}) q^{(n + \lfloor \frac{i+1}{2} \rfloor)^2 + [i \text{ even}] \binom{i+1}{2} - i} - (1 - q^{2n+2i+1}) q^{(n + \lfloor \frac{i}{2} \rfloor)^2 + [i \text{ even}] \binom{i}{2}} \right).
\end{aligned}$$

The last bracket in this expression can be simplified for i even:

$$-q^{(n + \frac{i}{2})^2 + \binom{i}{2} + 2i+1} (1 - q^{2n})$$

and for i odd:

$$-q^{(n + \frac{i-1}{2})^2} (1 - q^{2n}).$$

Putting everything together, we arrive at

$$C_{i+1} b_i - b_{i-1} = \sum_{n \geq 0} \frac{(-1)^{n-1} z^{2n}}{[2n + 2i + 1]_q!} \left(\prod_{j=0}^i [2n + 2j]_q \right) q^{(n + \lfloor \frac{i}{2} \rfloor)^2 + [i \text{ even}] \binom{i+2}{2}}.$$

Notice that the constant term vanishes, whence

$$\begin{aligned}
b_{i+1} &= \sum_{n \geq 1} \frac{(-1)^{n-1} z^{2n-2}}{[2n + 2i + 1]_q!} \left(\prod_{j=0}^i [2n + 2j]_q \right) q^{(n + \lfloor \frac{i}{2} \rfloor)^2 + [i \text{ even}] \binom{i+2}{2}} \\
&= \sum_{n \geq 0} \frac{(-1)^n z^{2n}}{[2n + 2(i+1) + 1]_q!} \left(\prod_{j=1}^{i+1} [2n + 2j]_q \right) q^{(n + \lfloor \frac{i+2}{2} \rfloor)^2 + [i+1 \text{ odd}] \binom{i+2}{2}},
\end{aligned}$$

which is the announced formula.

REFERENCES

1. M. FULMEK, *A continued fraction expansion for a q -tangent function*, Sém. Loth. Comb., **45**, (2000), [B45b] (5 pages).
2. F.H. JACKSON, *A basic-sine and cosine with symbolic solutions of certain differential equations*, Proc. Edinburgh Math. Soc., **22**, (1904), 28–39.
3. H. PRODINGER, *Combinatorics of geometrically distributed random variables: new q -tangent and q -secant numbers*, Int. J. Math. Math. Sci., **24**, (2000), 825–838.

HELMUT PRODINGER, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF STELLENBOSCH, 7602 STELLENBOSCH, SOUTH AFRICA

E-mail address: hproding@sun.ac.za