

An extension of a theorem by
Bismut–Zhang to Morse–Bott
vector fields

Dan Burghelea and Stefan Haller

Topology seminar
Max Planck Institut for Mathematics
September 17, 2004

Ray–Singer torsion

(M, g) . . . closed Riemannian manifold

(E, μ) . . . flat Euclidean vector bundle over M

↪ deRham complex $(\Omega^*(M; E), d)$

↪ Laplacian $\Delta^q : \Omega^q(M; E) \rightarrow \Omega^q(M; E)$

↪ zeta regularized $\det' \Delta^q$

↪ Ray–Singer torsion

$$\log T_{\text{an}}^{E, g, \mu} := \frac{1}{2} \sum_q (-)^{q+1} \log \det' \Delta^q$$

↪ deRham cohomology $H^q(M; E) = \ker \Delta^q$

↪ scalar product $\| \cdot \|_{H^q(M; E)}$

Analysis of the ODE

$X = -\text{grad}_{\tilde{g}}(f) \dots$ Morse–Bott–Smale

\rightsquigarrow critical manifold $\Sigma := \text{Zero}(X)$

\rightsquigarrow decomposition $TM|_{\Sigma} = T\Sigma \oplus N^+ \oplus N^-$

\rightsquigarrow Morse index $\text{ind} = \text{rank}(N^-) : \Sigma \rightarrow \mathbb{N}_0$

\rightsquigarrow unstable manifold

$$\Sigma \xleftarrow{p} W \xrightarrow{i} M$$

p diffeomorphic to N^- , $i : W \rightarrow M$ smooth

\rightsquigarrow vertical bundle $V_W \rightarrow W$

$$\text{rank}(V_W) = p^* \text{ind}$$

\rightsquigarrow space of unparametrized trajectories

$$\Sigma \xleftarrow{\pi_-} \mathcal{T} \xrightarrow{\pi_+} \Sigma$$

π_- smooth fiber bundle, π_+ smooth

\rightsquigarrow vertical bundle $V_{\mathcal{T}} \rightarrow \mathcal{T}$

$$\text{rank}(V_{\mathcal{T}}) = \pi_+^* \text{ind} - \pi_-^* \text{ind} - 1$$

\rightsquigarrow canonic compactification to manifold with corners

$$\Sigma \xleftarrow{\hat{\pi}_-} \hat{\mathcal{T}} \xrightarrow{\hat{\pi}_+} \Sigma$$

$\hat{\pi}_-$ smooth fiber bundle, $\hat{\pi}_+$ smooth,

$$\partial_1 \hat{\mathcal{T}} = \mathcal{T} \times_{\Sigma} \mathcal{T}$$

\rightsquigarrow canonic compactification to manifold with corners

$$\Sigma \xleftarrow{\hat{p}} \hat{W} \xrightarrow{\hat{i}} M$$

\hat{p} smooth fiber bundle, \hat{i} smooth,

$$\partial_1 \hat{W} = \mathcal{T} \times_{\Sigma} W$$

Geometric complex

X ... Morse–Bott–Smale vector field

E ... flat vector bundle

$$E_S := (E \otimes \mathcal{O}_{N^-})|_S$$

\rightsquigarrow \mathbb{Z} -graded Morse–Bott complex

$$C^q(X; E) := \bigoplus_{S \subseteq \Sigma} \Omega^{q - \text{ind}(S)}(S; E_S)$$

$$\delta := d \pm (\hat{\pi}_-)_* \circ (\hat{\pi}_+)^* \quad \delta^2 = 0$$

\rightsquigarrow cohomology $HC^*(X; E)$

\rightsquigarrow homomorphism

$$\Omega^q(M; E) \xrightarrow{\text{Int}} C^q(X; E)$$

$$\text{Int} := \hat{p}_* \circ \hat{i}^*$$

$$\text{Int} \circ d = \delta \circ \text{Int}$$

\rightsquigarrow isomorphism

$$H^*(M; E) \xrightarrow{\text{Int}} HC^*(X; E)$$

↪ filtration

$$C_{\geq p}^q(X; E) := \bigoplus_{\text{ind}(S) \geq p} \Omega^{q-\text{ind}(S)}(S; E_S)$$

↪ spectral sequence converging to $HC^*(X; E)$

$$E_1 C_p^q(X; E) = \bigoplus_{\text{ind}(S)=p} H^{q-p}(S; E_S)$$

g . . . Riemannian metric on M

μ . . . fiber metric on E

↪ (Σ, g_S) Riemannian

↪ (E_S, μ_S) flat Euclidean vector bundle

↪ scalar product $\|\cdot\|_{E_1}$

↪ inductively $\|\cdot\|_{E_k}$, $1 \leq k \leq \infty$

Combinatorial and metric torsion

↪ finite dimensional Laplacians Δ_k^q on $E_k C^q$

↪ combinatorial torsion

$$\log T_{\text{comb}}^{E,g,\mu,X} := \sum_k \frac{1}{2} \sum_q (-)^{q+1} q \log \det' \Delta_k^q$$

↪ $GHC^*(X; E) = E_\infty C^*(X; E)$

↪ $\det HC^* = \det GHC^* = \det E_\infty C^*$

↪ geometric scalar product $\|\cdot\|_{\det HC^*(X;E)}$

$$\det H^*(M; E) \xrightarrow{\det \text{Int}} \det HC^*(X; E)$$

↪ metric torsion

$$\log T_{\text{met}}^{E,g,\mu,X} := \log \text{vol det Int}$$

Note that $H^*(M; E) = 0 \Rightarrow \log T_{\text{met}} = 0$.

$\theta := -\frac{1}{2} \operatorname{tr}_\mu \nabla \mu \in \Omega^1(M; \mathbb{R}) \dots$ closed 1-form
 $\rightsquigarrow \Theta := [\theta] \in H^1(M; \mathbb{R})$ holonomy of $\det E \otimes \mathcal{O}_E$

$\Psi \in \Omega^{n-1}(TM \setminus M; \mathcal{O}_M) \dots$ Mathai–Quillen form

Theorem A

$$\begin{aligned} \log T_{\text{an}}^M &= \sum_{S \subseteq \Sigma} (-)^{\operatorname{ind}(S)} \log T_{\text{an}}^S \\ &\quad + \log T_{\text{met}} + \log T_{\text{comb}} \\ &\quad + \int_M \theta \wedge (-X)^* \Psi \end{aligned}$$

Remarks

- localization theorem for analytic torsion
- specializes to Bismut–Zhang theorem if X Morse–Smale (Σ discrete, hence $\log T_{\text{an}}^S = 0$)
- implies (a slight generalization of) a result of Lück–Schick–Thielmann (analytic torsion of fiber bundles)

Euler structures (Turaev)

M^n connected with base point x_0

consider equivalence classes $[X, c]$

X vector field with non-degenerate zeros \mathcal{X}

$c \in C_1(M; \mathbb{R})$... Turaev spider (Euler chain)

$$\partial c = \sum_{x \in \mathcal{X}} \text{IND}(x)x - \chi(M)x_0$$

$\text{IND}(x)$... Hopf index

$[X_1, c_1] = [X_2, c_2]$ iff

$$c_2 - c_1 = c(X_1, X_2) \pmod{\partial(C_2(M; \mathbb{R}))}$$

$c(X_1, X_2)$... one chain obtained as the zero set of a generic homotopy from X_1 to X_2

$\rightsquigarrow \mathcal{Eul}_{x_0}(M)$ Euler structures over \mathbb{R}
affine version of $H_1(M; \mathbb{R})$

Euler structures (deRham approach)

M^n connected with base point x_0

consider equivalence classes $[g, \alpha]$ where

$g \dots$ Riemannian metric

$\alpha \in \Omega^{n-1}(M \setminus x_0; \mathcal{O}_M)$ with

$$d\alpha = E_g$$

$E_g \in \Omega^n(M; \mathcal{O}_M) \dots$ Euler form

$[g_1, \alpha_1] = [g_2, \alpha_2]$ iff

$$\alpha_2 - \alpha_1 = \text{cs}(g_1, g_2) \pmod{d(\Omega^{n-1}(M; \mathcal{O}_M))}$$

$\rightsquigarrow \mathfrak{Eul}_{x_0}^*(M) \dots$ (co)Euler structures over \mathbb{R}
affine version of $H^{n-1}(M; \mathcal{O}_M)$

\rightsquigarrow isomorphism $\mathfrak{Eul}_{x_0}^*(M) = \mathfrak{Eul}_{x_0}(M)$
affine over Poincaré duality

$$H^{n-1}(M; \mathcal{O}_M) = H_1(M; \mathbb{R})$$

Analytic torsion

E . . . flat vector bundle over M

$\mathfrak{e} = [g, \alpha] \in \mathfrak{Eul}_{x_0}(M)$. . . Euler structure

consider real line (one dim. vector space)

$$\text{Det}_{x_0}(M; E) := \det H^*(M; E) \otimes (\det E_{x_0})^{-\chi(M)}$$

choose fiber metric μ on E and define

$$\|\cdot\|_{\text{an}} := T_{\text{an}} \cdot e^{-\int_M \theta \wedge \alpha} \cdot \|\cdot\|_{\det H^*(M; E)} \otimes \|\cdot\|_{x_0}$$

anomaly formula for analytic torsion implies that this does only depend on (E, \mathfrak{e}) .

\rightsquigarrow analytic torsion $\pm \tau_{\text{an}}^{E, \mathfrak{e}} \in \text{Det}_{x_0}(M)$

for $\sigma \in H_1(M; \mathbb{R})$ we have

$$\tau_{\text{an}}^{E, \mathfrak{e} + \sigma} = \tau_{\text{an}}^{E, \mathfrak{e}} \cdot e^{-\langle \Theta, \sigma \rangle}$$

Localization of Euler structures

consider equivalence classes $[X, c, \{\mathfrak{e}_S\}]$

X ... Morse–Bott–Smale vector field

$\mathfrak{e}_S \in \mathfrak{Eul}_{x_S}(S)$... Euler structure on $S \subseteq \Sigma$

$c \in C_1(M; \mathbb{R})$ with

$$\partial c = \sum_{S \subseteq \Sigma} \text{IND}(S)x_S - \chi(M)x_0$$

+ more complicated equivalence relation

$\rightsquigarrow \mathfrak{Eul}_{x_0}(M)$... affine version of $H_1(M; \mathbb{R})$

Geometric torsion

X ... Morse–Bott–Smale vector field

$\mathfrak{e} = [-X, c, \{\mathfrak{e}_S\}]$... Euler structure

$$\text{Det}_{x_0}(X; E) := \det HC^*(X; E) \otimes (\det E_{x_0})^{-\chi(M)}$$

spectral sequence provides canonic

$$\det HC^* = \det GHC^* = \det E_\infty C^* = \dots$$

$$= \det E_1 C^* = \bigotimes_{S \subseteq \Sigma} \left(\det H^*(S; E_S) \right)^{(-)^{\text{ind}(S)}}$$

+ parallel transport along c provides

$$\text{Det}_{x_0}(X; E) \stackrel{c}{=} \bigotimes_{S \subseteq \Sigma} \left(\text{Det}_{x_S}(S; E_S) \right)^{(-)^{\text{ind}(S)}}$$

\rightsquigarrow geometric torsion $\tau_{\text{geom}}^{E, \mathfrak{e}, X} \in \text{Det}_{x_0}(X; E)$

for $\sigma \in H_1(M; \mathbb{R})$ we have

$$\tau_{\text{geom}}^{E, \mathfrak{e} + \sigma, X} = \tau_{\text{geom}}^{E, \mathfrak{e}, X} \cdot e^{-\langle \Theta, \sigma \rangle}$$

Reformulation of theorem A

M ... closed manifold

E ... flat vector bundle over M

X ... Morse–Bott–Smale vector field on M

$\mathfrak{e} \in \mathfrak{Eul}_{x_0}(M)$... Euler structure

Int : $H^*(M; E) = HC^*(X; E)$ provides

$$\text{Det}_{x_0}(M; E) = \text{Det}_{x_0}(X; E)$$

Theorem B. $\tau_{\text{an}}^{E, \mathfrak{e}} = \tau_{\text{geom}}^{E, \mathfrak{e}, X}$

Tracing back the definitions we easily find

theorem A \Leftrightarrow theorem B

Homotopies

\mathbb{X} ... MBS homotopy from X_- to X_+

\rightsquigarrow homomorphism

$$C^*(X_-; E) \xrightarrow{\varphi_{\mathbb{X}}} C^*(X_+; E)$$

$$\varphi_{\mathbb{X}} \circ \delta_{X_-} = \delta_{X_+} \circ \varphi_{\mathbb{X}}$$

\rightsquigarrow homotopy

$$C^*(X_-; E) \xrightarrow{H_{\mathbb{X}}} C^{*-1}(X_+; E)$$

$$H_{\mathbb{X}} \circ d + \delta_{X_+} \circ H_{\mathbb{X}} = \varphi_{\mathbb{X}} \circ \text{Int}_{X_-} - \text{Int}_{X_+}$$

\rightsquigarrow commuting isomorphisms

$$\begin{array}{ccc}
 & H^*(M; E) & \\
 \text{Int}_{X_-} \swarrow & & \searrow \text{Int}_{X_+} \\
 HC^*(X_-; E) & \xrightarrow{\varphi_{\mathbb{X}}} & HC^*(X_+; E)
 \end{array}$$

A small perturbation

X ... MBS vector field

Y ... Morse–Smale vector field on Σ

MBS homotopy $\mathbb{X}_t := X + tY$

- $X_- = X$ and X_+ Morse–Smale
- X_+ tangential to Σ and $X_+|_{\Sigma} = Y$
- $\Sigma_{X_+} \subseteq \Sigma_X$, $\text{ind}_{X_+}^M(x) = \text{ind}(S) + \text{ind}_Y^{\Sigma}(x)$
- $x, x' \in S \Rightarrow \mathcal{T}_{Y_S}(x, x') = \mathcal{T}_{X_+}(x, x')$
- $x \in S \neq S' \ni x', \mathcal{T}(S, S') = \emptyset \Rightarrow \mathcal{T}(x, x') = \emptyset$
- \mathbb{X} tangential to Σ
- $x \in S \Rightarrow \mathcal{T}_{\mathbb{X}}(x, S) = W_{Y_S}^-(x)$
- $x \in S \neq S', \mathcal{T}_X(S, S') = \emptyset \Rightarrow \mathcal{T}_{\mathbb{X}}(x, S') = \emptyset$

\rightsquigarrow filtration compatible with δ_{X_+}

$$C(X_+; E)_{\geq p} := \bigoplus_{\text{ind}(S) \geq p} C(X_+|_S; E_S)$$

\rightsquigarrow homomorphism compatible with filtrations

$$\varphi_{\mathbb{X}} : C_{\geq p}^*(X; E) \rightarrow C^*(X_+; E)_{\geq p}$$

\rightsquigarrow E_1 -term identifies to

$$\begin{array}{ccc} E_1 C_p^q(X; E) & \xrightarrow{\varphi_{\mathbb{X}}} & E_1 C^q(X^+; E)_p \\ \parallel & & \parallel \\ H^{q-p}(\Sigma_p; E_{\Sigma_p}) & \xrightarrow{\text{Int}_Y} & HC^{q-p}(Y_{\Sigma_p}; E_{\Sigma_p}) \end{array}$$

Consider

$$\tau_{\text{geom}}^{E, \mathbf{e}, X} \in \text{Det}_{x_0}(X; E)$$

$$\tau_{\text{geom}}^{E, \mathbf{e}, X_+} \in \text{Det}_{x_0}(X_+; E)$$

$$\tau_{\text{geom}}^{E_S, \mathbf{e}_S, Y_S} \in \text{Det}_{x_S}(Y_S; E_S)$$

as linear mappings

$$(\det E_{x_0})\chi(M) \xrightarrow{\tau_{\text{geom}}^{E, \mathbf{e}, X}} \det HC^*(X; E)$$

$$(\det E_{x_0})\chi(M) \xrightarrow{\tau_{\text{geom}}^{E, \mathbf{e}, X_+}} \det HC^*(X_+; E)$$

$$(\det E_{x_S})\chi(S) \xrightarrow{\tau_{\text{geom}}^{E_S, \mathbf{e}_S, Y_S}} \det HC^*(Y_S; E_S)$$

$$\mathbf{e} = [-X, c, \{\mathbf{e}_S\}]$$

$$\mathbf{e}_S = [-Y_S, c_S]$$

$$c_+ := c + \sum (-)^{\text{ind}(S)} c_S$$

$$\Rightarrow \mathbf{e} = [-X_+, c_+]$$

$$\begin{array}{ccccc}
& & \det H^*(M; E) & \xleftarrow{\tau_{\text{an}}^{E, \epsilon}} & (\det E_{x_0})^{\chi(M)} \\
& \swarrow \text{Int}_X & \downarrow \text{Int}_{X^+} & \swarrow \tau_{\text{geom}}^{E, \epsilon, X^+} & \downarrow c^+ \\
\det HC^*(X; E) & \xrightarrow{\varphi_X} & \det HC^*(X^+; E) & \xlongequal{\quad} & \det C^*(X^+; E) \\
\parallel & & \parallel & & \parallel \\
\det E_\infty C^*(X; E) & \xrightarrow{\varphi_X} & \det E_\infty C^*(X^+; E) & & \\
\vdots & & \vdots & & \\
\det E_2 C^*(X; E) & \xrightarrow{\varphi_X} & \det E_2 C^*(X^+; E) & & \\
\parallel & & \parallel & & \parallel \\
\det E_1 C^*(X; E) & \xrightarrow{\varphi_X} & \det E_1 C^*(X^+; E) & \xlongequal{\quad} & \det GC^*(X^+; E) \\
\parallel & & \parallel & & \parallel \\
\bigotimes (\det H^*(S; E_S))^\pm & \xrightarrow{\text{Int}_{Y_S}} & \bigotimes (\det HC^*(Y_S; E_S))^\pm & \xlongequal{\quad} & \bigotimes (\det C^*(Y_S; E_S))^\pm \\
& \swarrow \otimes (\tau_{\text{an}}^{E_S, \epsilon_S})^\pm & \uparrow \otimes (\tau_{\text{geom}}^{E_S, \epsilon_S})^\pm & \swarrow \otimes (c_S)^\pm & \\
& & \bigotimes ((\det(E_S)_{x_S})^{\chi(S)})^\pm & & \\
& & \uparrow c & & \\
& & (\det E_{x_0})^{\chi(M)} & &
\end{array}$$

this commutes!!!

- Bismut–Zhang applied to X^+
- Bismut–Zhang applied to all $S \subseteq \Sigma$
- A theorem of Maumary (spectral sequences)
- what we discussed up to now