# RAAGEDY RIGHT-ANGLED COXETER GROUPS II: IN THE QUASIISOMETRY CLASS OF THE TREE RAAGS

## CHRISTOPHER H. CASHEN

ABSTRACT. We classify two-dimensional right-angled Coxeter groups that are quasiisometric to a right-angled Artin group defined by a tree, and show that when this is true the right-angled Coxeter group actually contains a visible finite index right-angled Artin subgroup.

#### 1. INTRODUCTION

A tree RAAG is a right-angled Artin group whose presentation graph is a tree of diameter at least three. Such a group is the fundamental group of a compact 3-manifold with boundary whose JSJ decomposition, in the 3-manifold sense, contains only Seifert fibered pieces. A corollary of Behrstock and Neumann's [1] quasiisometry classification of graph manifolds is that all tree RAAGs belong to a single quasiisometry class. The diameter condition excludes the well-understood quasiisometry classes  $\mathbb{Z}$ ,  $\mathbb{Z}^2$ , and  $\mathbb{F}_2 \times \mathbb{Z}$ .

Nguyen and Tran [20] classified planar, triangle-free graphs  $\Gamma$  such that the rightangled Coxeter group (RACG)  $W_{\Gamma}$  with presentation graph  $\Gamma$  is in the quasiisometry class of the tree RAAGs. Dani and Levcovitz [11] subsequently showed that when Nguyen and Tran's conditions are satisfied then  $W_{\Gamma}$  actually has a finite index tree RAAG subgroup that is *visible* in  $\Gamma$ , in a sense that will be made precise later.

Nguyen and Tran's proof uses planarity in an essential way twice: once to invoke the Jordan Curve Theorem to construct a certain suspension-decomposition of  $\Gamma$ , and again to say that  $W_{\Gamma}$  is virtually a 3-manifold group. Then their suspensiondecomposition corresponds to the JSJ decomposition of the manifold and they apply Behrstock and Neumann's result. Dani and Levcovitz's argument also explicitly uses a planar embedding of  $\Gamma$ . However, there are easy examples, such as the graph in Figure 1, of nonplanar graphs  $\Gamma$  such that  $W_{\Gamma}$  contains a finite index tree RAAG subgroup (see Example 3.2), so planarity is not a necessary condition.



FIGURE 1. A nonplanar, triangle-free graph  $\Gamma$  such that  $W_{\Gamma}$  is in the quasiisometry class of the tree RAAGs.

This work is licensed under a Creative Commons "Attribution 4.0 International" license.



Date: June 10, 2025.

<sup>2020</sup> Mathematics Subject Classification. 20F65,20F55.

 $Key\ words\ and\ phrases.$  Right-angled Coxeter group, right-angled Artin group, RACG, RAAG, quasiisometry.

This research was funded by the Austrian Science Fund (FWF) 10.55776/P34214.

#### CHRISTOPHER H. CASHEN

We give a complete classification of triangle-free graphs  $\Gamma$  such that  $W_{\Gamma}$  is in the quasiisometry class of the tree RAAGs, and show that in this case  $W_{\Gamma}$  contains a visible finite index tree RAAG subgroup. See Theorem 3.1. The proof uses the group theoretic version of JSJ decompositions (over 2–ended subgroups). Such decompositions of RAAGs and RACGs can be read off from their presentation graphs. One step in the proof is to describe what the JSJ decomposition of a group in the quasiisometry class of the tree RAAGs must look like. The second is to show that, for a 2–dimensional RACG, these conditions are already strong enough to produce a visible finite index RAAG subgroup.

There are infinitely many commensurability classes of tree RAAGs [2, 3], so the main theorem does not say that quasiisometry to a particular tree RAAG implies commensurability to that same tree RAAG. Nevertheless, we find the commensurability aspect of the theorem surprising. Usually when a geometric relation can be promoted to an algebraic relation it is because there is some strong rigidity phenomenon at play, but the class of tree RAAGs is quite flexible. For example, the results of Huang and Kleiner [17] on groups quasiisometric to RAAGs with finite outer automorphism group do not apply to tree RAAGs.

Acknowledgements. This paper is a spin-off of [6], where the question is when  $W_{\Gamma}$  is quasiisometric to any RAAG whatsoever.

Alexandra Edletzberger had already noticed that Nguyen and Tran's decomposition was related to the JSJ decomposition, and that cycles of cuts should be obstructions to being quasiisometric to a RAAG [15, Remark 1.38, Remark 4.29].

# 2. Preliminaries

2.1. Terminology. The right-angled Artin group with presentation graph  $\Delta$ , for a finite simplicial graph  $\Delta$ , is the group  $A_{\Delta}$  presented by a generator for each vertex of  $\Delta$  and a commuting relation between two generators when they are connected by an edge of  $\Delta$ . The right-angled Coxeter group with presentation graph  $\Gamma$ ,  $W_{\Gamma}$ , is defined similarly, with additional relations saying that each generator has order 2. In both cases these groups have a geometric action on a CAT(0) cube complex, which has dimension at most 2 when the presentation graph is triangle-free.

 $\mathbb{F}_n$  is the free group of rank n, and  $\mathbb{F}$  is  $\mathbb{F}_n$  for some unspecified  $n \ge 2$  when the precise rank is not important. A *basis* of  $\mathbb{F}$  means a free generating set.

An *induced* subgraph of a simplicial graph is a subgraph that contains all possible edges between its vertices. If  $\Upsilon'$  is an induced subgraph of  $\Upsilon$  then the inclusion at the graph level extends to inclusions  $A_{\Upsilon'} < A_{\Upsilon}$  and  $W_{\Upsilon'} < W_{\Upsilon}$ . A subgroup of a RAAG or RACG defined in this way is called a *special subgroup*.

If  $\Gamma$  is a presentation graph of a RACG then a *thick join* means a join subgraph A \* B of  $\Gamma$  such that  $W_A$  and  $W_B$  are infinite groups, which happens when A and B are incomplete subgraphs. A *suspension* is a join in which A is a 2-anticlique. If A is a 2-anticlique and |B| > 2 then A is called the *pole* or the *suspension points* of A \* B and B is called the *suspended points*. A vertex is *essential* if it has valence at least 3. A *cone vertex* is everybody's neighbor. If v is a vertex, its *link* lk(v) is its set of neighbors and its *star* is  $st(v) := lk(v) \cup \{v\}$ . A graph is *biconnected* if it is connected with no cut vertex. Thus, a single edge is biconnected.

Two subsets of a metric space are r-coarsely equivalent if their Hausdorff distance is at most r, and coarsely equivalent if their Hausdorff distance is finite. Coarse equivalence gives an equivalence relation that is preserved by quasiisometries. 2.2. Visible RAAG subgroups. If  $\Gamma$  is a graph, its complement graph  $\Gamma^c$  is the graph on the same vertex set that has an edge precisely when  $\Gamma$  does not. Suppose  $\Lambda \subset \Gamma^c$  is a disjoint union of trees  $\Lambda_0$  and  $\Lambda_1$ . For a set of vertices V in  $\Gamma$ , if V is contained in one of the  $\Lambda_i$  then  $\operatorname{hull}_{\Lambda}(V)$  is the set of vertices of  $\Gamma$  contained in the convex hull of V in  $\Lambda_i$ . The commuting graph  $\Delta$  of  $\Lambda$  is the graph that has a vertex for each edge of  $\Lambda$ , with an edge between two of them if their support spans a square in  $\Gamma$ . There is a natural homomorphism  $A_{\Delta} \to W_{\Gamma}$  defined by sending the generator of  $A_{\Delta}$  corresponding to vertex  $\{a, b\}$  of  $\Delta$  to the element  $ab \in W_{\Gamma}$ . it is easy to come up with examples where this homomorphism is not injective, but Dani and Levcovitz [11] give conditions that imply the homomorphism is injective and the image has finite index.

**Theorem 2.1** ([11, Theorem 4.8, Corollary 3.33, Remark 4.3]). Let  $\Gamma$  be an incomplete, triangle-free graph without separating cliques. Suppose  $\Lambda$  is a disjoint union of trees  $\Lambda_0$  and  $\Lambda_1$  in  $\Gamma^c$  such that  $\Lambda$  spans  $\Gamma$  and for each *i*, no two vertices of  $\Lambda_i$ are adjacent in  $\Gamma$ . Let  $\Delta$  be the commuting graph of  $\Lambda$ .

If  $\Lambda$  satisfies the following conditions, then the homomorphism  $A_{\Delta} \to W_{\Gamma}$  is injective and its image has index 4 in  $W_{\Gamma}$ .

- $\mathcal{R}_3$ : If  $\{a, b\} * \{c, d\}$  is a square in  $\Gamma$  then  $\operatorname{hull}_{\Lambda}\{a, b\} * \operatorname{hull}_{\Lambda}\{c, d\} \subset \Gamma$ .
- $\mathcal{R}_4$ : If  $a \mapsto b$  is an edge in a cycle  $\gamma \subset \Gamma$  then there is a square  $\{a, a'\} * \{b, b'\}$ with  $a', b' \in \operatorname{hull}_{\Lambda}(\gamma)$ .

Call a graph  $\Lambda$  satisfying the hypotheses of Theorem 2.1 a finite index Dani-Levcovitz  $\Lambda$  (FIDL- $\Lambda$ ). When we have a FIDL- $\Lambda$ , the commuting graph  $\Delta$  gives a 'visible' finite index RAAG subgroup of  $W_{\Gamma}$ . It is 'visible' in the sense that vertices of  $\Delta$  correspond to diagonals of squares of  $\Gamma$ , and vertices of  $\Delta$  span an edge precisely when their support spans a square in  $\Gamma$ .

2.3. CFS graphs. For  $W_{\Gamma}$  to be quasiisometric to a RAAG it must have at most quadratic divergence, which is characterized by  $\Gamma$  having property *CFS* [12].

**Definition 2.2** (cf.[7, Definition 2.3]). The *diagonal graph*  $\Box(\Gamma)$  of  $\Gamma$  is the graph with an edge for each induced square of  $\Gamma$ , with vertices representing diagonal pairs. The *support* of a vertex of  $\Box(\Gamma)$  is the two vertices of  $\Gamma$  in the corresponding diagonal pair. The support of a subset of  $\Box(\Gamma)$  is the union of supports of its vertices.  $\Gamma$  is *CFS* if  $\Box(\Gamma)$  contains a component whose support is all non-cone vertices of  $\Gamma$ .

We will not invoke the CFS property directly, but  $\square(\Gamma)$  will figure prominently. Consider that  $A_{\Delta}$  is 1-ended when  $\Delta$  is connected and not a single vertex, and  $W_{\Gamma}$  is 1-ended when  $\Gamma$  is incomplete without separating cliques. Thus, in the 1-ended case, for a FIDL- $\Lambda$  as in the previous section, we may identify  $\Delta$  with a subgraph of  $\square(\Gamma)$ , since  $\Delta$  is connected and an edge in  $\Delta$  corresponds to a square in  $\Gamma$ .

2.4. **JSJ** decompositions. We assume familiarity with Bass-Serre theory. In this paper we are interested in JSJ decompositions of finitely presented groups over 2–ended subgroups (equivalently, over virtually  $\mathbb{Z}$  subgroups), which are graph of groups decompositions that encode all possible splittings of the group over 2–ended subgroups. The existence of non-trivial JSJ decompositions, for finitely presented groups not commensurable to a surface group, is a quasiisometry invariant [22], but the precise structure of a particular JSJ decomposition is not. One can make a canonical object by passing to the Bass-Serre tree of any JSJ decomposition and

collapsing cylinders to make the JSJ tree of cylinders [16]. It is a bipartite tree where one part consists of cylinder vertices, which correspond to commensurability classes of universally elliptic 2–ended splitting subgroups, and one part is made up of rigid and hanging vertices that belong to more than one cylinder. Hanging vertices contain the non-(universally elliptic) 2–ended splitting subgroups; that is, they contain collections of 2–ended splitting subgroups that give incompatible group splittings. Rigid vertices correspond to subgroups that are not split by any 2–ended splitting of the ambient group.

It follows from Papasoglu's work [22] (cf. [9]) that a quasiisometry  $\phi: G \to G'$ between two finitely presented groups induces an isomorphism of their JSJ trees of cylinders,  $\phi_*: T \to T'$ , that preserves vertex type: cylinder/rigid/hanging. Furthermore, if X is a tree of spaces for G over T, and similarly X' for G' over T', then restriction of  $\phi$  to each vertex space  $X_v$  is uniformly close to a quasiisometry  $\phi_v: X_v \to X'_{\phi_*(v)}$  that induces a bijection of incident edge spaces by taking  $X_e \subset X_v$  to a set uniformly coarsely equivalent to  $X'_{\phi_*(e)} \subset X'_{\phi_*(v)}$ .

The quotient of the action of a group on its JSJ tree of cylinders gives the graph of cylinders. Since the JSJ tree of cylinders can be recovered by development, this graph of groups contains information on quasiisometry invariants for the group, such as the vertex types that occur, the quasiisometry classes the vertex groups, and the relative quasiisometry types of the vertices relative to the peripheral pattern of their incident edge groups, which will be discussed in Section 2.5.

Work of Clay [10] and Margolis [18], says that the graph of cylinders of a RAAG can be described 'visually' in terms of the presentation graph  $\Delta$  as follows, where in each case the local group is the special subgroup defined by the relevant subgraph:

- Cylinders are stars of cut vertices of  $\Delta$ .
- Rigid vertices are maximal biconnected subgraphs of  $\Delta$  that are not contained in a single cylinder.
- Hanging vertices do not occur.
- There is an edge between a cylinder and rigid vertex when they intersect.

Edletzberger [14], building on work of Mihalik and Tschantz [19] and Dani and Thomas [13], gave a visual description of the graph of cylinders of a 1-ended RACG with a triangle-free presentation graph  $\Gamma$  that is not a cycle graph. The 2-ended subgroups of interest manifest as special subgroups generated by *cuts*  $\{a - b\}$ , meaning *a* and *b* are not adjacent and either:

- $\{a, b\}$  is a cut pair, that is,  $\Gamma \{a, b\}$  is not connected.
- $\{a, b\}$  is not a cut pair, but there is a vertex  $c \in lk(a) \cap lk(b)$  such that the 2-path  $a \rightarrow c \rightarrow b$  disconnects  $\Gamma$ . In this case  $a \rightarrow c \rightarrow b$  is a *cut 2-path*.

The distinction between 2–ended splitting subgroups that are non-(universally elliptic) vs universally elliptic is reflected topologically: A cut gives a non-(universally elliptic) splitting if there is some other cut that *crosses* it, in the sense that each cut contains vertices in different components of the complement of the other. A cut gives a universally elliptic splitting if it is *uncrossed*, ie, not crossed by any cut.

- Cylinders are suspensions  $\{a, b\} * (\operatorname{lk}(a) \cap \operatorname{lk}(b))$  for uncrossed cut  $\{a b\}$ .
- Rigid vertices are sets of essential vertices of Γ of size at least four that cannot be separated by a cut and that are maximal with respect to inclusion among such sets.

- Hanging vertices consist of maximal collections of pairwise crossing cut pairs or pairwise crossing cut 2–paths. (Cut pairs cannot cross cut 2–paths, and vice versa [7, after Definition 2.8].)
- There is an edge between a cylinder  $\{a, b\}*(lk(a) \cap lk(b))$  and a rigid/hanging vertex if the latter contains a cut  $\{a-b\}$ . In this case the edge corresponds to intersection of subgraphs.

2.5. **Peripheral patterns.** A peripheral pattern  $\mathcal{P}(G, \mathcal{H})$  in a group G relative to a collection  $\mathcal{H} := \{H_0, \ldots, H_{n-1}\}$  of subgroups of G is the set of coarse equivalence classes of left cosets of the  $H_i$  in G. When  $\mathcal{H}$  consists of 2-ended subgroups the peripheral pattern it generates may be called a *line pattern*. A map between two groups with peripheral patterns is *pattern preserving* if it takes each coset of one pattern to within uniformly bounded Hausdorff distance of one in the other, and vice versa, inducing a bijection between patterns. The *relative quasiisometry type* of  $\mathcal{P}(G, \mathcal{H})$  is the equivalence class of spaces with peripheral patterns up to pattern preserving quasiisometry.

Line patterns in free groups have been considered before [21, 8, 5]. Associated to a line pattern there is a *decomposition space*, which is the quotient of the boundary of the free group obtained by identifying the two endpoints of each line in the pattern. Pattern preserving quasiisometries induce homeomorphisms of decomposition spaces. Two cases are of particular interest for us. It turns out that each of these types defines a single relative quasiisometry type of line pattern.

**Definition 2.3.** A surface type (line) pattern in  $\mathbb{F}$  is a peripheral pattern  $\mathcal{P}(\mathbb{F}, \mathcal{H})$  with  $\mathcal{H} = \{\langle f_0 \rangle, \ldots, \langle f_{n-1} \rangle\}$  such that, up to passing to roots, conjugates, and inverses, and removing duplicate entries, there is a compact surface with fundamental group  $\mathbb{F}$  such that  $\{f_0, \ldots, f_{n-1}\}$  are the elements of  $\mathbb{F}$  represented by the boundary curves. Equivalently, its decomposition space is a circle.

**Definition 2.4.** A basic (line) pattern in  $\mathbb{F}$  is a peripheral pattern  $\mathcal{P}(\mathbb{F}, \mathcal{H})$  with  $\mathcal{H} = \{\langle f_0 \rangle, \ldots, \langle f_{n-1} \rangle\}$  such that, up to passing to roots, conjugates, and inverses, and removing duplicate entries,  $\{f_0, \ldots, f_{n-1}\}$  is a subset of a basis of  $\mathbb{F}$ . Equivalently, its decomposition space is totally disconnected.

In the present work we are interest not in line patterns in free groups, but in *plane patterns* in  $\mathbb{F} \times \mathbb{Z}$ . Let z be a generator of the center. By a *plane pattern* in  $\mathbb{F} \times \mathbb{Z}$ , we mean a peripheral pattern  $\mathcal{P}(\mathbb{F} \times \mathbb{Z}, \mathcal{H})$  where  $\mathcal{H}$  consists of  $\mathbb{Z}^2$  subgroups, which we might as well assume, up to coarse equivalence, are maximal  $\mathbb{Z}^2$  subgroups. A maximal  $\mathbb{Z}^2$  subgroup of  $\mathbb{F} \times \mathbb{Z}$  is of the form  $\langle f \rangle \times \langle z \rangle$ , where  $f \in \mathbb{F}$  and  $z \in \mathbb{Z}$  are nontrivial elements that are not proper powers. Using the results and terminology of [22], the only separating quasilines of  $\mathbb{F} \times \mathbb{Z}$  are coarsely equivalent to the central  $\mathbb{Z}$ . Thus, a quasiisometry  $\phi \colon \mathbb{F} \times \mathbb{Z} \to \mathbb{F}' \times \mathbb{Z}$  takes each coset of the central  $\mathbb{Z}$  to within bounded Hausdorff distance, depending only on the quasiisometry constants, of a coset of the central  $\mathbb{Z}$ . From this it follows that  $\pi_{\mathbb{F}'} \circ \phi|_{\mathbb{F} \times \{1\}} \colon \mathbb{F} \to \mathbb{F}'$ , where  $\pi_{\mathbb{F}'} \colon \mathbb{F}' \times \mathbb{Z}$  is projection to the first factor given by killing the center, is a quasiisometry. Conversely,  $(\pi_{\mathbb{F}'} \circ \phi|_{\mathbb{F} \times \{1\}}) \times \mathrm{Id}_{\mathbb{Z}} \colon \mathbb{F} \times \mathbb{Z} \to \mathbb{F}' \times \mathbb{Z}$ is a quasiisometry (but not necessarily one that is close to  $\phi$ ). The projections  $\pi_{\mathbb{F}} \colon \mathbb{F} \times \mathbb{Z} \to \mathbb{F}$  and  $\pi_{\mathbb{F}'} \colon \mathbb{F}' \times \mathbb{Z} \to \mathbb{F}'$  send peripheral plane patterns to peripheral line patterns, and the original quasiisometry  $\phi$  preserves plane patterns if and only if  $\pi_{\mathbb{F}'} \circ \phi|_{\mathbb{F}\times\{1\}}$  preserves the projected line patterns. We extend the basic/surface terminology to the plane patterns according to the type of the projection. More

#### CHRISTOPHER H. CASHEN

generally, extend the terminology to patterns of virtually  $\mathbb{Z}^2$  subgroups in virtually  $\mathbb{F} \times \mathbb{Z}$  groups, according to the type of the pattern obtained by pushing forward the pattern by the map induced by the restriction to a finite index  $\mathbb{F} \times \mathbb{Z}$  subgroup.

**Lemma 2.5.** Let  $\Delta$  be a finite tree of diameter at least three, so that  $A_{\Delta}$  is a 1ended RAAG with a nontrivial graph of cylinders. For every cylinder in the graph of cylinders, the peripheral pattern coming from the incident edge groups is basic.

Proof. Since  $\Delta$  is a tree, every non-leaf is a cut vertex, so cylinders are stars of nonleaf vertices v of  $\Delta$ . If v is a non-leaf then  $lk(v) = \{w_0, \ldots, w_{n-1}, \ell_0, \ldots, \ell_{m-1}\}$ , where the  $w_i$  are non-leaves and the  $\ell_j$  are leaves. Since the diameter of  $\Delta$  is greater than 2, n > 0 and  $m + n \ge 2$ . The cylinder group is  $A_{st(v)} = \langle v \rangle \times \langle w_0, \ldots, w_{n-1}, \ell_0, \ldots, \ell_{m-1} \rangle \cong \mathbb{Z} \times \mathbb{F}_{m+n}$ . Since  $\Delta$  is a tree, the maximal biconnected subgraphs are single edges, and those that are not contained in a single cylinder are the non-leaf edges. Thus, the neighbors of the cylinder st(v) in the graph of cylinders are the rigid vertices  $v \mapsto w_i$  for  $0 \le i < n$ , with  $A_{\{v,w_i\}} \cong \mathbb{Z}^2$ . These give a basic plane pattern, since  $\{w_0, \ldots, w_{n-1}\}$  is a subset of the basis  $\{w_0, \ldots, w_{n-1}, \ell_0, \ldots, \ell_{m-1}\}$  of the  $\mathbb{F}_{m+n}$  factor of  $A_{st(v)}$ .

**Example 2.6.** Let  $\Gamma$  be the nonplanar graph in Figure 2a. Let  $W_{\Gamma}$  be the RACG whose presentation graph is  $\Gamma$ , so that the generators of  $W_{\Gamma}$  are  $s_i$  where  $s_i$  corresponds to vertex i of  $\Gamma$ . (For compactness of notation, we omit the 's' in figures and subscripts.) The graph of cylinders of  $W_{\Gamma}$  is shown in Figure 2b, where the labels are the cylinder groups and each unlabelled vertex is a rigid vertex whose group is the intersection of its neighboring cylinder groups, which in each case is isomorphic to the product of two infinite dihedral groups. All edge groups are isomorphic to their neighboring rigid vertex group, and all edge maps are inclusion.



FIGURE 2. Graph  $\Gamma$  and graph of cylinders of  $W_{\Gamma}$  in Example 2.6.

The middle cylinder vertex  $C := W_{\{3,4\}} \times W_{\{0,1,2,8\}}$  has peripheral pattern generated by  $\mathcal{H} := \{W_{\{0,1\}} \times W_{\{3,4\}}, W_{\{1,2\}} \times W_{\{3,4\}}, W_{\{0,2\}} \times W_{\{3,4\}}\}$ . The group C has an index 4 subgroup isomorphic to  $\mathbb{F}_3 \times \mathbb{Z}$ , namely  $\langle w, x, y \rangle \times \langle z \rangle$ , where  $w := s_0 s_8$ ,  $x := s_0 s_1, y := s_1 s_2$ , and  $z := s_3 s_4$ . Restriction to the finite index subgroup induces a pattern preserving quasiisometry between  $\mathcal{P}(C, \mathcal{H})$  and  $\mathbb{F}_3 \times \mathbb{Z}$  with plane pattern generated by  $\{\langle x \rangle \times \langle z \rangle, \langle y \rangle \times \langle z \rangle, \langle xy \rangle \times \langle z \rangle\}$ . Projection to the  $\mathbb{F}_3$  factor gives line pattern generated by  $\{x, y, xy\}$ . Since the generators live in a free factor  $\langle x, y \rangle$  of  $\mathbb{F}_3$ , it follows that the decomposition space for  $\mathcal{P}(\langle x, y \rangle, \{x, y, xy\})$  embeds into the decomposition space for  $\mathcal{P}(\mathbb{F}_3, \{x, y, xy\})$ . The former is a surface type pattern: it matches the standard description of the fundamental group and boundary curves of a 3-holed sphere. Therefore, the decomposition space for  $\mathcal{P}(\mathbb{F}_3, \{x, y, xy\})$  contains circles; in particular, it is not totally disconnected, so the line pattern is not basic. Thus,  $W_{\Gamma}$  is not quasiisometric to a tree RAAG, since the non-basic plane pattern in the cylinder  $W_{\{0,1,2,8\}} \times W_{\{3,4\}}$  of the RACG cannot match the plane patterns in cylinder vertices of tree RAAGs, which by Lemma 2.5 are all basic.  $\diamond$ 

In the planar case, according to Nguyen and Tran's argument,  $\Gamma$  decomposes into a tree of maximal suspensions, with neighbors determined by two suspensions sharing a square. Playing the game as in the example by restricting to a finite index torsion-free subgroup and then projecting to the free factor, one can see that there are only two possibilities for a maximal suspension  $\{a, b\} * \{c_0, \ldots, c_{n-1}\}$ : either there are fewer than *n* neighbors and the line pattern is basic, or there are exactly *n* neighbors and the line pattern is surface type. Nguyen and Tran use 'suspension of *n* vertices has exactly *n* neighboring suspensions' as an obstruction to being quasiisometric to a tree RAAG. Example 2.6 illustrates that in the nonplanar case there are more varied ways for a pattern to fail to be basic; *C* has a non-basic pattern, but it is a suspension of 4 vertices with only 3 neighboring suspensions.

## 3. RACGS QUASIISOMETRIC TO TREE RAAGS

**Theorem 3.1.** Let  $\Gamma$  be an incomplete, triangle-free graph without separating cliques. The following are equivalent:

- (1)  $W_{\Gamma}$  has an index four visible tree RAAG subgroup.
- (2)  $W_{\Gamma}$  is in the quasiisometry class of the tree RAAGs.
- (3)  $W_{\Gamma}$  has a nontrivial graph of cylinders with no hanging vertices and:
  - (a) Cylinders are virtually  $\mathbb{F} \times \mathbb{Z}$ .
  - (b) Rigid vertices are virtually  $\mathbb{Z}^2$ . Each is adjacent to two cylinders, with edge group equal to the rigid vertex group.
  - (c) In each cylinder the incident edge groups form a basic plane pattern.
- (4)  $\Gamma$  has the following structure:
  - (a) Maximal thick joins are non-square suspensions. Every vertex and edge is contained in a maximal suspension. The pole of each maximal suspension gives a cut, in the sense that it is either a cut pair or forms a cut 2-path with one of the suspended vertices. All cuts come from the pole of a maximal suspension in one of these two ways.
  - (b) Every set of at least four essential vertices that is not separated by a cut is a square, each of whose diagonals is the pole of a maximal suspension.
  - (c) There does not exist a maximal suspension  $\{a, b\} * \{c_0, \ldots, c_{n-1}\}$  and  $3 \leq m \leq n$  such that for all  $0 \leq i < m$  the pair  $\{c_i, c_{(i+1) \mod m}\}$  is the pole of a maximal suspension.

*Proof.* Inclusion of a finite index subgroup induces a quasiisometry, so  $(1) \Longrightarrow (2)$ .

The graph of cylinders of a tree RAAG was described in the proof of Lemma 2.5. Quasiisometry invariance of the JSJ tree of cylinders implies that any group quasiisometric to a tree RAAG has JSJ tree of cylinders whose cylinders are quasiisometric to  $\mathbb{F} \times \mathbb{Z}$ , whose rigid vertices are quasiisometric to  $\mathbb{Z}^2$ , has no hanging vertices, has exactly two cylinders adjacent to each rigid vertex, and such that in each cylinder the plane pattern coming from incident edges is a basic plane pattern. 'Quasiisometric to'  $\mathbb{F} \times \mathbb{Z}$  or  $\mathbb{Z}^2$  can be promoted to 'virtually'  $\mathbb{F} \times \mathbb{Z}$  or  $\mathbb{Z}^2$ , respectively, since these groups are quasiisometrically rigid. Finally, if  $\tilde{r}$  is a valence 2 rigid vertex of the JSJ tree of cylinders and r is its image in the graph of cylinders then r either has two adjacent cylinders connected to r via edge groups equal to the vertex group of r, or it has one adjacent cylinder connected to r via an edge group that is an index 2 subgroup of the vertex group of r. The latter is impossible, by visibility of the graph of cylinders in RACGs, since when  $\Gamma$  is triangle-free the only virtually  $\mathbb{Z}^2$  special subgroups are defined by squares of  $\Gamma$ , so one cannot be properly contained in another. Thus,  $(2) \implies (3)$ .

Assume (3). Cylinders are suspensions  $\{a, b\} * (lk(a) \cap lk(b))$  where there is an uncrossed cut  $\{a - b\}$ . By (3a), the suspension is non-square. Rigid vertices are maximal subsets of  $\Gamma$  of at least four essential vertices that cannot be separated by a cut. By (3b) they are virtually  $\mathbb{Z}^2$ , so they correspond to squares in  $\Gamma$  that are not separated by a cut.

A  $K_{3,3}$  subgraph of  $\Gamma$  cannot be separated by a cut. A suspension  $\{a, b\} * C$  with  $|C| \ge 3$  can only be separated by a cut of the form  $\{a - b\}$ , ie  $\{a, b\}$  or a - c - b for  $c \in C$ . Thus the existence of either a maximal thick join that is not a suspension or a non-square suspension whose suspension points do not make a cut in one of these two ways implies there is a non-virtually- $\mathbb{Z}^2$  rigid vertex, contradicting (3b). Thus, maximal thick joins are non-square suspensions whose pole gives a cut. Since there are no hanging vertices, all cuts arise this way. Since the graph of cylinders gives a visual decomposition, every vertex and edge of  $\Gamma$  is contained in a subgraph corresponding to a vertex of the graph of cylinders. Since there are no hanging vertices are assumed, by (3b), to be contained in cylinders, every vertex and edge is contained in a cylinder. This establishes (4a).

We have said rigid vertices correspond to certain squares and cylinders correspond to maximal suspensions. A square is contained in a maximal suspension if and only if one of its diagonals is the pole of the suspension. Since a rigid vertex is assumed to be adjacent to two cylinders, both diagonals of the corresponding square must be poles of maximal suspensions. This establishes (4b).

Item (4c) describes a cylinder vertex of the graph of cylinders whose incident edges form a non-basic plane pattern, contrary to (3c).

It remains to show (4)  $\implies$  (1). A graph  $\Delta$  for which  $A_{\Delta}$  is a finite index tree RAAG subgroup of  $W_{\Gamma}$  can be constructed essentially by taking the graph of cylinders of  $\Gamma$  and adding some leaves. We prove this by constructing a FIDL- $\Lambda$ with  $\Delta$  as its commuting graph, as in Theorem 2.1. This will take some work.

Assume (4). Any graph of groups decomposition of  $W_{\Gamma}$  has underlying graph a tree because  $W_{\Gamma}$  is generated by torsion elements, so it cannot surject onto  $\mathbb{Z}$ . Let T be the tree with one vertex for each cylinder in the graph of cylinders and an edge between cylinders if they share a square. This is just the underlying graph of the graph of cylinders, where we have 'forgotten' the rigid vertices, in the sense that they are all of valence two, so we think of them as the midpoint of an edge instead of as a valence two vertex. In fact, we will think of T as a subgraph of  $\square(\Gamma)$ , whose edges were defined to be the squares of  $\Gamma$ .

We make some claims about the graph structure of  $\Gamma$  deduced from (4). Assuming the claims, we finish the proof of (1). After that we will prove the claims.

Claim 3.1.1. For every square  $\{a, b\} * \{c, d\}$  of  $\Gamma$ , at least one of the diagonals  $\{a, b\}$  or  $\{c, d\}$  is the pole of a non-square suspension that is a maximal thick join.

Claim 3.1.2.  $\square(\Gamma)$  is a tree and T can be identified, via the map sending a maximal thick join  $\{a, b\} * C$  to the vertex  $\{a, b\} \in \square(\Gamma)$ , with the subtree obtained by removing all leaves of  $\square(\Gamma)$ ,

Examples illustrating Claim 3.1.3 are shown in Figure 3. In the claim, when  $\{p_0, q_0\}, \{p_1, q_1\}, \ldots, \{p_n, q_n\}$  is a geodesic in  $T \subset \square(\Gamma)$  then for a vertex  $v \in \Gamma$  let  $I_v := \{i \mid v \in \{p_i, q_i\}\}$ . Also, let  $\mathbb{E}$  denote the even numbers, and  $\mathbb{O}$  the odd.

Claim 3.1.3. If  $\{p_0, q_0\}, \{p_1, q_1\}, \dots, \{p_n, q_n\}$  is a geodesic in T with n > 0 and there exists a vertex  $z \in \Gamma$  with  $\{p_0, q_0, p_n, q_n\} \subset \text{lk}(z)$  then:

- (i)  $n \in \mathbb{E}$
- (ii)  $I_z = [0, n] \cap \mathbb{O}$
- (iii) For all 0 < i < n, there is a cut  $\{p_i q_i\}$  separating  $\{p_j, q_j \mid j < i\} \{p_i q_i\}$ from  $\{p_j, q_j \mid i < j\} - \{p_i - q_i\}$ . If  $i \in \mathbb{E}$  then the cut is  $\{p_i - q_i\} = p_i \rightarrow z \rightarrow q_i$ .



FIGURE 3. Two graphs as in Claim 3.1.3 with their diagonal graphs containing the geodesic  $\{p_0, q_0\}, \{p_1, q_1\}, \ldots, \{p_n, q_n\}$ .

Claim 3.1.4. If  $\{a, b\} \neq \{a, c\}$  are vertices in  $\square(\Gamma)$  then the distance between them is even and every second vertex on the geodesic between them contains a.

Claim 3.1.5. Every vertex of  $\Gamma$  is suspended in some maximal suspension. A vertex that is not a suspension point of any maximal suspension is contained in a unique maximal suspension.

Assuming the claims, we construct a FIDL– $\Lambda$ .

By Claim 3.1.2,  $\square(\Gamma)$  is a tree, so it can be 2-colored 0/1 such that neighboring vertices have different colors. By Claim 3.1.4, if v is in the support of two different vertices of  $\square(\Gamma)$  then they have even distance in  $\square(\Gamma)$ , so they have the same color. Thus, the 2-coloring on  $\square(\Gamma)$  induces a 2-coloring of  $\Gamma$ .

Let  $\{a, b\}$  be a vertex of T, corresponding to a maximal suspension  $\{a, b\} * \{c_0, \ldots, c_{n-1}\}$ . Let  $\Lambda_{a,b} \subset \Gamma^c$  be a graph on vertices  $\{c_0, \ldots, c_{n-1}\}$ , constructed as follows. Add an edge in  $\Lambda_{a,b}$  between  $c_i$  and  $c_j$  if  $\{c_i, c_j\}$  is a vertex of T. Call these 'mandatory edges'. Condition (4c) implies the mandatory edges form a forest. Add

additional 'discretionary' edges to make  $\Lambda_{a,b}$  a tree. Let  $T_i \subset T$  be the vertices with color *i*. Let  $\Lambda_0$  be the union of trees  $\Lambda_{a,b}$  such that  $\{a,b\} \in T_1$ , so that  $\sup(\Lambda_0)$ consists of vertices of  $\Gamma$  colored 0. Define  $\Lambda_1$  analogously for  $T_0$ . By Claim 3.1.5,  $\Lambda := \Lambda_0 \sqcup \Lambda_1$  spans  $\Gamma$ . Vertices of  $\Lambda_i$  are colored *i*, so they are not adjacent in  $\Gamma$ .

We prove  $\Lambda_0$  is a tree by induction. Pick a vertex  $\{a_0, b_0\} \in T_1$ . By construction,  $\Lambda_{a_0,b_0}$  is a tree. Now suppose that  $\bigcup_{i=0}^{m-1} \Lambda_{a_i,b_i}$  is a tree, where the set of vertices  $\{a_i, b_i\} \in T_1$  with i < m is the intersection of a convex subset of T with  $T_1$ . Let  $\{a_m, b_m\} \in T_1$  that is closest to the existing set. Since  $\Box(\Gamma)$  is a tree, the convexity condition implies that there is a unique vertex  $\{c, d\} \in T_0$  separating  $\{a_m, b_m\}$ from all of the other  $\{a_i, b_i\}$ , and there is at least one index j < m such that  $\{c, d\}$  is a neighbor of  $\{a_j, b_j\}$  in T. Thus,  $c \mapsto d$  is a mandatory edge in  $\Lambda_{a_m, b_m}$ and in any such  $\Lambda_{a_j, b_j}$ . We have  $c \mapsto d \subset \Lambda_{a_m, b_m} \cap \bigcup_{i=0}^{m-1} \Lambda_{a_i, b_i}$ . Conversely, suppose, for some j < m, that  $\Lambda_{a_j, b_j} \cap \Lambda_{a_m, b_m} \neq \emptyset$ , so there is a vertex  $z \in \Gamma$ with  $\{a_j, b_j, a_m, b_m\} \subset lk(z)$ . Since  $\{c, d\}$  is the penultimate vertex on the unique geodesic from  $\{a_j, b_j\}$  to  $\{a_m, b_m\}$  in  $\Box(\Gamma)$ , Claim 3.1.3 says  $z \in \{c, d\}$ . Thus,  $\Lambda_{a_m, b_m} \cap \bigcup_{i=0}^{m-1} \Lambda_{a_i, b_i}$  is exactly the edge  $c \mapsto d$ , but the union of two trees with exactly an edge in common is a tree, so  $\Lambda_0$  is a tree by induction, and  $\Lambda_1$  similarly.

Now we check that conditions  $\mathcal{R}_3$  and  $\mathcal{R}_4$  of Theorem 2.1 are satisfied.

Suppose  $\{a, b\} * \{c, d\}$  is a square in  $\Gamma$ . Claim 3.1.1 says either  $\{a, b\}$  or  $\{c, d\}$  is the pole of a maximal suspension; suppose  $\{a, b\} * C$  is a maximal suspension with  $\{c, d\} \subset C$ . Then  $\operatorname{hull}_{\Lambda}\{a, b\} = \{a, b\}$  and  $\operatorname{hull}_{\Lambda}\{c, d\} \subset \Lambda_{a, b} \subset C$ , so  $\operatorname{hull}_{\Lambda}\{a, b\} * \operatorname{hull}_{\Lambda}\{c, d\} \subset \{a, b\} * C \subset \Gamma$ . Condition  $\mathcal{R}_3$  is satisfied.

Suppose condition  $\mathcal{R}_4$  is not satisfied. Then there is a shortest simple cycle  $\gamma$  in  $\Gamma$  containing an edge e that is not in a square with vertices in hull<sub> $\Lambda$ </sub>( $\gamma$ ). Such a  $\gamma$  is induced, because if  $\gamma$  contains vertices v and w that are adjacent in  $\Gamma$  but not in  $\gamma$  then we could surger  $\gamma$  into two simple cycles  $\gamma'$  and  $\gamma''$  that are each an arc of  $\gamma$  plus the edge  $v \mapsto w$ . Both are strictly shorter than  $\gamma$ , and one of them, say  $\gamma'$ , contains e, but hull<sub> $\Lambda$ </sub>( $\gamma'$ )  $\subset$  hull<sub> $\Lambda$ </sub>( $\gamma$ ), so e is not contained in a square with vertices in hull<sub> $\Lambda$ </sub>( $\gamma'$ ). This would contradict minimality of  $\gamma$ .

Let  $u \to v \to w \to x$  be the subsegment of  $\gamma$  such that  $e = v \to w$ . If  $u \to v$  and  $w \to x$  are contained in a common suspension then so is  $u \to x$ , and  $\{u, w\} * \{v, x\}$  is a square, contradicting the definition of e. Thus, we may assume that there is not a suspension containing all three of the edges. The argument will be to produce a cut through v or w separating u from x. Since  $\gamma$  is a loop, it must pass back through the cut again at some other vertex v', which will lead to a contradiction with the assumption that  $\gamma$  is induced. Thus, there is no witness to the failure of  $\mathcal{R}_4$ .

**Case 1:** There is a maximal suspension  $\sigma$  containing  $u \mapsto v \mapsto w$  and a different maximal suspension  $\sigma'$  containing  $v \mapsto w \mapsto x$ . Suppose the distance between  $\sigma$  and  $\sigma'$  in T is minimal among pairs with these properties.

**Case 1a:** Suppose  $\sigma$  and  $\sigma'$  are adjacent in T. Their intersection is the join of their poles. If  $\{u, w\}$  is the pole of  $\sigma$  then all four of u, v, w, and x are contained in  $\sigma'$ , contrary to hypothesis, and similarly for  $\sigma'$ , so there are vertices  $v' \neq w'$  such that  $\sigma \cap \sigma' = \{v, v'\} * \{w, w'\}$ . If no cut separates u from x then  $\{u\} * \{v, v'\} * \{w, w'\} * \{x\}$  is a non-square rigid vertex, which is impossible, so there is at least one cut of the form  $\{v-v'\}$  or  $\{w-w'\}$  separating u from x. These cases are symmetric, so assume it is  $\{v - v'\}$ . Since v is adjacent to u and w, if  $\gamma$  passes through v' then  $\gamma$  fails to be induced. However, the only way for  $\gamma$  to pass back through the cut avoiding

v' is if the cut is a cut 2-path  $v \leftarrow u' \leftarrow v'$  with  $u' \neq w$ . In this case,  $\gamma$  passes back through u', which is adjacent to v, so  $\gamma$  is not induced.

**Case 1b:** Suppose  $\sigma$  and  $\sigma'$  are not adjacent in T. For  $\{a, b\} \in T$ , let  $\sigma_{a,b} := \{a, b\} * (link(a) \cap link(b))$  be the maximal suspension of  $\Gamma$  with pole  $\{a, b\}$ . Let  $\{p_0, q_0\}, \ldots, \{p_n, q_n\}$  be the geodesic in T between  $\sigma = \sigma_{p_0, q_0}$  and  $\sigma' = \sigma_{p_n, q_n}$ . Vertices v and w cannot both be suspended in the same suspension since they are connected by an edge and  $\Gamma$  is triangle free, so, up to symmetry, there are two subcases: either w is a suspended and v is suspension in both  $\sigma$  and  $\sigma'$ , or v is a suspension point of  $\sigma$  and suspended in  $\sigma'$  and the opposite is true for w.

In the first subcase,  $\{p_0, q_0\} = \{v, v'\}$ , for some v', and  $\{p_n, q_n\} = \{v, x\}$ . Claim 3.1.3 says  $n \in \mathbb{E}$ ,  $w \in \{p_1, q_1\}$ , and there is a cut  $\{p_1 - q_1\}$  separating  $\{v, v'\} - \{p_1 - q_1\}$  from  $\{v, x\} - \{p_1 - q_1\}$ . Since  $v \in \{v, v'\} \cap \{v, x\}$ , the cut is the cut 2-path  $p_1 \rightarrow v \rightarrow q_1$ , and it separates v' from x. It also separates u from x, since u is adjacent to v'. So,  $\gamma$  crosses the cut once through v to get from u to x, and must cross back again later, but both of the other vertices of the cut are adjacent to v, contradicting that  $\gamma$  is induced.

In the second subcase we have that w is suspended in  $\sigma_{p_{n-1}q_{n-1}}$ , and, by minimality of  $n, u \notin \sigma_{p_1,q_1}$  and  $x \notin \sigma_{p_{n-1},q_{n-1}}$ . Apply Claim 3.1.3 to  $\{p_0, q_0\}, \ldots, \{p_{n-1}, q_{n-1}\}$  with z = w, which gives that  $w \in \{p_1, q_1\}$  and there is a cut  $\{p_1 - q_1\}$  in  $\Gamma$  separating  $\{p_0, q_0\} - \{p_1 - q_1\}$  from  $\{p_{n-1}, q_{n-1}\} - \{p_1 - q_1\}$ . Since  $u \notin \sigma_{p_1,q_1}$  and  $x \notin \sigma_{p_{n-1},q_{n-1}}, \{p_1 - q_1\}$  separates u from x. Since v is adjacent to u and v and x are both suspended in  $\sigma'$  with two common neighbors  $\{p_n, q_n\} \neq \{p_1, q_1\}$ , the cut  $\{p_1 - q_1\}$  is a cut 2–path  $p_1 \rightarrow v \rightarrow q_1$ . Thus,  $\gamma$  crosses  $\{p_1 - q_1\}$  once through v and then back again through a vertex adjacent to v, contradicting that it is induced.

**Case 2:** The path  $u \to v \to w$  is not contained in a maximal suspension. Let  $\{p_0, q_0\}, \ldots, \{p_n, q_n\}$  be a geodesic in T such that  $u, v \in \sigma_{p_0, q_0}$  and  $v, w \in \sigma_{p_n, q_n}$ , and such that n > 0 is minimal among such geodesics. Edges sharing a common vertex cannot be contained in adjacent maximal suspensions without both being contained in at least one of the two, so  $n \ge 2$ . Minimality implies  $|\{u, v\} \cap \{p_0, q_0\}| = 1$  and  $\{u, v\} \cap \{p_1, q_1\} = \emptyset$  and  $|\{v, w\} \cap \{p_n, q_n\}| = 1$  and  $\{v, w\} \cap \{p_{n-1}, q_{n-1}\} = \emptyset$ .

If v is suspended in both  $\sigma_{p_0,q_0}$  and  $\sigma_{p_n,q_n}$  then Claim 3.1.3 says  $v \in \{p_1,q_1\}$ , but this contradicts minimality of n.

If v is suspended in  $\sigma_{p_0,q_0}$  but not in  $\sigma_{p_n,q_n}$  then  $v \in \{p_n,q_n\}$  is adjacent to both  $p_{n-1}$  and  $q_{n-1}$ . Apply Claim 3.1.3 to the geodesic subsegment from  $\{p_0,q_0\}$  to  $\{p_{n-1},q_{n-1}\}$ . Again, this implies  $v \in \{p_1,q_1\}$ , contradicting minimality of n. The symmetric argument works if v is suspended in  $\sigma_{p_n,q_n}$  but not in  $\sigma_{p_0,q_0}$ .

Finally, suppose v is a suspension point in both  $\sigma_{p_0,q_0}$  and  $\sigma_{p_n,q_n}$ . If n = 2 then apply Claim 3.1.3 with  $z = p_1$ . If n > 2 then apply Claim 3.1.3 to the geodesic subsegment  $\{p_1, q_1\}, \ldots, \{p_{n-1}, q_{n-1}\}$  with z = v. In both cases we get that  $p_{n-1} \rightarrow v \rightarrow q_{n-1}$  is a cut 2-path, and the loop  $\gamma$  contains v and has vertices u and w on different sides of the cut, so it must cross the cut a second time in either  $p_{n-1}$  or  $q_{n-1}$ , both of which are adjacent to v, contradicting that  $\gamma$  is induced.

Up to symmetry, this accounts for all possibilities, so we have constructed a FIDL- $\Lambda$ . By Theorem 2.1, the commuting graph  $\Delta$  of  $\Lambda$  defines an index 4 visible RAAG subgroup of  $W_{\Gamma}$ . It follows from Behrstock and Neumann's [1] work that  $\Delta$  must be a tree, because tree RAAGs are not quasiisometric to nontree RAAGs, but we can say explicitly what  $\Delta$  is from the construction: it is a tree built by starting

from T and at each vertex  $\{a, b\} \in T$  adding a new leaf for each discretionary edge of  $\Lambda_{a,b}$ . This completes the proof of (4)  $\implies$  (1), modulo the claims.

Proof of Claim 3.1.1: Any square  $\{a, b\} * \{c, d\}$  in  $\Gamma$  is a thick join, so it is contained in some maximal thick join  $\sigma$ , which, by (4a) is a non-square suspension. Since  $\Gamma$ is triangle-free either  $\{a, b\}$  or  $\{c, d\}$  is the pole of  $\sigma$ .  $\Diamond$ 

Proof of Claim 3.1.2: If  $\{a, b\} * \{c, d\}$  in  $\Gamma$  is a square and  $\{a, b\} * \{x, y\}$  is another square then  $|\{c, d, x, y\}| \ge 3$ . The thick join  $\{a, b\} * \{c, d, x, y\}$  is contained in a maximal thick join, which is a suspension, by (4a), so  $\{a, b\}$  is the pole of a maximal thick join that is a non-square suspension.

Every non-leaf of  $\square(\Gamma)$  is a pair of vertices of  $\Gamma$  that are the diagonal of more than one square, so, as above, is the pole of a non-square suspension. These correspond to vertices of T, so all of the non-leaf vertices of  $\square(\Gamma)$  are identified with vertices of T. Furthermore, adjacent vertices of  $\square(\Gamma)$  span a square, so T-vertices of  $\square(\Gamma)$  are adjacent in  $\square(\Gamma)$  if and only if they are adjacent in T. Thus, we identify T with a subgraph of  $\square(\Gamma)$  that includes all non-leaves of  $\square(\Gamma)$ . By Claim 3.1.1,  $\square(\Gamma)$  has no isolated edges, so every leaf is attached to one of the T-vertices of  $\square(\Gamma)$ . Thus,  $\square(\Gamma)$  is a tree and T is a subtree containing all non-leaves.

For  $\{a, b\} \in T$  there is a maximal thick join  $\{a, b\} * \{c_0, c_1, c_2, ...\}$  in  $\Gamma$ . Since  $\Gamma$  is triangle free, there are induced squares  $\{a, b\} * \{c_i, c_j\}$  for all  $i \neq j$ . Thus,  $\{a, b\}$  has at least three neighbors in  $\Box(\Gamma)$ . In particular, it is not a leaf.

Proof of Claim 3.1.3: The proof is by induction on n. Since  $\Gamma$  is triangle-free, n > 1. Suppose n = 2. Since  $\{p_0, q_0\}$  and  $\{p_2, q_2\}$  are distinct vertices in T,  $|\{p_0, q_0, p_2, q_2\}| \ge 3$ , so  $\{p_0, q_0, p_2, q_2, z\}$  is a set of at least 4 essential vertices that is not a square, since z is adjacent to at least three of the others. By (4b), the set must be separated by a cut, since otherwise there would be a non-(virtually  $\mathbb{Z}^2$ ) rigid vertex in the graph of cylinders. By (4a), for all i, vertices  $p_i$  and  $q_i$ have at least three common neighbors, so they cannot be separated by a cut. Now,  $\{z, p_1, q_1\} * \{p_0, q_0, p_2, q_2\} \subset \Gamma$ , but thick joins in  $\Gamma$  are suspensions, by (4a), so  $|\{z, p_1, q_1\}| = 2$ . Thus,  $z \in \{p_1, q_1\}$ , and there is a cut  $\{p_1 - q_1\}$  separating  $\{p_0, q_0\} - \{p_1 - q_1\}$  from  $\{p_2, q_2\} - \{p_1 - q_1\}$ . This proves the claim when n = 2.

Now consider a geodesic of length n > 2 as in the statement of the claim, and suppose the claim is true for all geodesics in T of length strictly less than n. As in the base case, for all i there is no cut separating  $p_i$  from  $q_i$ , but there must be a cut separating  $\{p_0, q_0, p_n, q_n, z\}$ , and the cut must contain z. Thus, there is some cut containing z and separating a subset of  $\{p_0, q_0\}$  from a subset of  $\{p_n, q_n\}$ . The geodesic  $\{p_0, q_0\}, \ldots, \{p_n, q_n\}$  in T gives a chain of squares in  $\Gamma$  with adjacent squares sharing a diagonal. Such a set is 2-connected, and the only potential cut pairs are  $\{p_i, q_i\}$  for 0 < i < n. Thus:

(†) There exists  $0 < i_0 < n$  such that there is a cut  $\{p_{i_0} - q_{i_0}\} \supset \{p_{i_0}, q_{i_0}, z\}$  separating the set  $\{p_0, q_0, p_n, q_n, z\}$ .

The goal will be to split the T-geodesic into two subsegments at index  $i_0$  and apply the induction hypothesis to both sides. We will also use the following fact:

- (†) No cut of the form  $\{p_i q_i\}$  separates the set  $\{p_j, q_j \mid j < i\} \{p_i q_i\}$  or
- (\*) separates the set  $\{p_j, q_j \mid j > i\} \{p_i q_i\}.$

To see (‡), first note that for all j < i,  $\{p_j, q_j\} \notin \{z, p_i, q_i\}$ . This is true because  $\{p_j, q_j\} \neq \{p_i, q_i\}$ , and  $p_j$  and  $q_j$  are not adjacent, but z is adjacent to both of  $p_i$  and

 $q_i$ . Thus, for all j < i at least one of  $p_j$  or  $q_j$  survives in  $\{p_j, q_j \mid j < i\} - \{p_i - q_i\}$  and any vertices with consecutive indices are adjacent.

The combination of (†) and (‡) implies there exists  $0 < i_0 < n$  such that there is a cut  $\{p_{i_0} - q_{i_0}\} \supset \{p_{i_0}, q_{i_0}, z\}$  separating  $\{p_i, q_i \mid i < i_0\} - \{p_{i_0} - q_{i_0}\}$  from  $\{p_i, q_i \mid i > i_0\} - \{p_{i_0} - q_{i_0}\}$ . Consider two cases based on whether z is in  $\{p_{i_0}, q_{i_0}\}$ . **Case**  $z \notin \{p_{i_0}, q_{i_0}\}$ : Then the cut  $\{p_{i_0} - q_{i_0}\}$  is a cut 2-path  $p_{i_0} \rightarrow z \rightarrow q_{i_0}$ . Consider the geodesic subsegments  $\{p_0, q_0\}, \ldots, \{p_{i_0}, q_{i_0}\}$  and  $\{p_{i_0}, q_{i_0}\}, \ldots, \{p_n, q_n\}$ . Each is strictly shorter than the one we started with and has all vertices in the supports of its two endpoints contained in lk(z). Apply the induction hypothesis. Then  $i_0 \in \mathbb{E}$  and  $n - i_0 \in \mathbb{E}$ , so  $n \in \mathbb{E}$ . We have the even number  $i_0$  is not in  $I_z$ , and by induction z occurs in the support of each odd index of each geodesic subsegment, so  $I_z = [0, n] \cap \mathbb{O}$ . Finally, the induction hypothesis for the first subsegment says that for every  $i < i_0$  there is a cut  $\{p_i - q_i\}$  separating  $\{p_j, q_j \mid 0 \leq j < i\} - \{p_i - q_i\}$ from  $\{p_j, q_j \mid i < j \leq i_0\} - \{p_i - q_i\}$ , but (‡) says  $\{p_i - q_i\}$  does not separate

from  $\{p_j, q_j \mid i < j \leq i\} = \{p_i - q_i\}$ , but (4) says  $\{p_i - q_i\}$  does not separate  $\{p_j, q_j \mid i < j \leq n\} - \{p_i - q_i\}$ , so it separates  $\{p_j, q_j \mid 0 \leq j < i\} - \{p_i - q_i\}$  from  $\{p_j, q_j \mid i < j \leq n\} - \{p_i - q_i\}$ . A symmetric argument on the second geodesic subsegment finishes the proof of (iii).

**Case**  $z \in \{p_{i_0}, q_{i_0}\}$ : Without loss of generality, assume  $z = p_{i_0}$ . First suppose  $1 < i_0 < n-1$ , and consider the geodesic subsegments  $\{p_0, q_0\}, \ldots, \{p_{i_0-1}, q_{i_0-1}\}$  and  $\{p_{i_0+1}, q_{i_0+1}\}, \ldots, \{p_n, q_n\}$ . Again, each is strictly shorter than the one we started with and has all vertices in the support of its two endpoints contained in lk(z), so induct. We get  $i_0 - 1 \in \mathbb{E}$  and  $n - (i_0 - 1) \in \mathbb{E}$ , so  $n \in \mathbb{E}$  and  $i_0 \in \mathbb{O}$ . We have that  $I_z$  contains the odd index  $i_0$ , since  $z = p_{i_0}$ , and the induction hypothesis for the two subsegments gives that  $I_z$  contains every other odd index as well. The proof of (iii) is the same as in the previous case.

If  $i_0 = 1$  or  $i_0 = n - 1$  the proof is the same, except one of the geodesic subsegments is a single vertex, so the induction is only necessary on the other.

Proof of Claim 3.1.4: By Claim 3.1.2,  $\Box(\Gamma)$  is a tree, so there is a unique geodesic  $\{a, b\} = \{p_0, q_0\}, \ldots, \{p_n, q_n\} = \{a, c\}$ . The support of a single edge has 4 vertices, so n > 1. If n = 2 the claim is true. We cannot have n = 3, because then  $\{p_1, q_1\} * \{p_2, q_2\}$  is a square in  $\Gamma$  contained in  $\mathbb{Ik}(a)$ , contradicting that  $\Gamma$  is triangle-free. If n > 3, apply Claim 3.1.3 to  $\{p_1, q_1\}, \ldots, \{p_{n-1}, q_{n-1}\}$  with z = a.

Proof of Claim 3.1.5: Since  $\Gamma$  is triangle-free, the intersection of suspensions corresponding to adjacent vertices in T is exactly the square formed by the join of the two poles. By (4a), every vertex of  $\Gamma$  is contained in some maximal suspension. If  $v \in \Gamma$  is a suspension point of a maximal suspension  $\sigma$ , then it is suspended in each neighbor of  $\sigma$  in T, so every vertex of  $\Gamma$  is suspended in some suspension.

If  $z \in \sigma \cap \sigma'$  is contained in two maximal suspensions, but is not in the pole of either, then  $\sigma$  and  $\sigma'$  do not give adjacent vertices in T. Claim 3.1.3 says zoccurs in the support of each odd index vertex on the T-geodesic from  $\sigma$  to  $\sigma'$ . Thus, every vertex that is contained in distinct maximal suspensions is in the pole of some maximal suspension.  $\diamond$ 

**Example 3.2.** The graph of Figure 1 is shown again with its graph of cylinders in Figure 4. The hypotheses of Theorem 3.1 are satisfied, because the central cylinder vertex has basic plane pattern:  $W_{\{3,4\}*\{0,1,2,8\}}$  has an index 4 subgroup  $\langle w, x, y \rangle \times \langle z \rangle \cong \mathbb{F}_3 \times \mathbb{Z}$  given by  $w := s_0 s_8$ ,  $x := s_1 s_8$ ,  $y := s_2 s_8$ , and  $z := s_3 s_4$ ,

such that restriction to the subgroup induces a pattern preserving quasiisometry to  $\mathbb{F}_3 \times \mathbb{Z}$  with plane pattern generated by  $\{\langle w \rangle \times \langle z \rangle, \langle x \rangle \times \langle z \rangle, \langle y \rangle \times \langle z \rangle\}$ , so the projected line pattern is generated by  $\{w, x, y\}$  in  $\langle w, x, y \rangle$ , which is basic.  $\diamond$ 



FIGURE 4. Graph and graph of cylinders for Example 3.2.

# 4. Further remarks

The implication  $(3) \implies (2)$  of Theorem 3.1 can be proven directly as in [4]. The description of the JSJ decomposition means these groups, like the tree RAAGs, are 'two-line tubular groups with bounded height change', which are all quasiisometric [4, Example 5.1]. The implication  $(4) \implies (3)$  is also easy. It should be possible to generalize Edletzberger's [14] visual description of the graph of cylinders in the case that  $\Gamma$  has triangles. In this case one could hope to replace the 'no-triangles' hypothesis on  $\Gamma$  by 'no-icosahedra' and retain the equivalence between (2), (3), and (4). Graphs with no icosahedra are the most general family for which such a classification would be interesting, because if  $\Gamma$  contains an icosahedron then  $\mathbb{Z}^3 \leq W_{\Gamma}$ , so  $W_{\Gamma}$  is not quasiisometric to a tree RAAG.

The equivalence of (1) to the others if the 'no-triangles' hypothesis is relaxed is another matter. Theorem 2.1 is an application of conditions of Dani and Levcovitz [11] specialized to the triangle-free case. Their work partially handles more general situations, but when there are triangles there are more conditions and they are only necessary, not sufficient, for  $\Lambda$  to spawn a visible RAAG subgroup. The 'sufficient' direction would be needed to extend Theorem 3.1 to handle triangles.

# References

- J. A. Behrstock and W. D. Neumann, Quasi-isometric classification of graph manifold groups, Duke Math. J. 141 (2008), no. 2, 217–240.
- [2] M. Casals-Ruiz, I. Kazachkov, and A. Zakharov, On commensurability of right-angled Artin groups I: RAAGs defined by trees of diameter 4, Rev. Mat. Iberoam. 35 (2019), no. 2, 521– 560.
- [3] M. Casals-Ruiz, I. Kazachokov, and A. Zakharov, On commensurability of right-angled Artin groups II: RAAGs defined by paths, Math. Proc. Cambridge Philos. Soc. 170 (2021), no. 3, 559–608.
- [4] C. H. Cashen, Quasi-isometries between tubular groups, Groups Geom. Dyn. 4 (2010), no. 3, 473–516.
- [5] C. H. Cashen, Splitting line patterns in free groups, Algebr. Geom. Topol. 16 (2016), no. 2, 621–673.
- [6] C. H. Cashen, P. Dani, A. Edletzberger, and A. Karrer, *RAAGedy right-angled Coxeter groups*, in preparation.
- [7] C. H. Cashen and A. Edletzberger, Visual right-angled Artin subgroups of two-dimensional right-angled Coxeter groups, J. Group Theory (in press).
- [8] C. H. Cashen and N. Macura, *Line patterns in free groups*, Geom. Topol. 15 (2011), no. 3, 1419–1475.

- C. H. Cashen and A. Martin, Quasi-isometries between groups with two-ended splittings, Math. Proc. Cambridge Philos. Soc. 162 (2017), no. 2, 249–291.
- [10] M. Clay, When does a right-angled Artin group split over Z?, Internat. J. Algebra Comput. 24 (2014), no. 6, 815–825.
- [11] P. Dani and I. Levcovitz, Right-angled Artin subgroups of right-angled Coxeter and Artin groups, Algebr. Geom. Topol. 24 (2024), 755–802.
- [12] P. Dani and A. Thomas, Divergence in right-angled Coxeter groups, Trans. Amer. Math. Soc. 367 (2015), no. 5, 3549–3577.
- [13] P. Dani and A. Thomas, Bowditch's JSJ tree and the quasi-isometry classification of certain Coxeter groups, J. Topol. 10 (2017), no. 4, 1066–1106.
- [14] A. Edletzberger, Quasi-isometries for certain right-angled Coxeter groups, Groups Geom. Dyn. 18 (2024), no. 3, 1037–1098.
- [15] A. Edletzberger, Quasi-isometries for two-dimensional right-angled Coxeter groups, Ph.D. thesis, University of Vienna, 2024.
- [16] V. Guirardel and G. Levitt, JSJ decompositions of groups, Astérisque (2017), no. 395.
- [17] J. Huang and B. Kleiner, Groups quasi-isometric to right-angled Artin groups, Duke Math. J. 167 (2018), no. 3, 537–602.
- [18] A. Margolis, Quasi-isometry classification of RAAGs that split over cyclic subgroups, Groups Geom. Dyn. 14 (2020), 1351–1417.
- [19] M. Mihalik and S. Tschantz, Visual decompositions of Coxeter groups, Groups Geom. Dyn. 3 (2009), no. 1, 173–198.
- [20] H. T. Nguyen and H. C. Tran, On the coarse geometry of certain right-angled Coxeter groups, Algebr. Geom. Topol. 19 (2019), no. 6, 3075–3118.
- [21] J.-P. Otal, Certaines relations d'equivalence sur l'ensemble des bouts d'un groupe libre, J. London Math. Soc. s2-46 (1992), no. 1, 123–139.
- [22] P. Papasoglu, Quasi-isometry invariance of group splittings, Ann. of Math. (2) 161 (2005), no. 2, 759–830.

Faculty of Mathematics, University of Vienna, Oskar-Morgenstern-Platz 1, 1090 Vienna, Austria, 100000002-6340-469X

Email address: christopher.cashen@univie.ac.at