

# Oscillators and Resonance


- 1 -

harmonic oscillator:

$$\ddot{x} + \omega^2 x = 0$$

$$\begin{cases} \dot{x} = y \\ \dot{y} = -\omega^2 x \end{cases}$$

with solution  $\begin{cases} x(t) = A \cos \omega t + B \sin \omega t \\ y(t) = -A \omega \sin \omega t + B \omega \cos \omega t. \end{cases}$



All solutions are periodic except a stationary point of center type at the origin.

Pendulum

$$\ddot{x} + \omega^2 \sin x = 0$$

Newton's equation becomes

$$F = m \frac{d^2x}{dt^2}$$


force      acceleration  
mass

$$-F_g \sin x = m l \ddot{x}$$

Insert  $F_g = mg$   
and divide by  $ml$ :

$$\ddot{x} + \frac{g}{l} \sin x = 0$$

$= \omega^2$  = frequency squared.



$$\begin{aligned} F_g &= \text{gravitational force} \\ &= mg \end{aligned}$$

gravitational constant

$\approx 9.8 \frac{m}{sec^2}$  on Earth

Different derivation via preservation of energy:

$$E_{\text{kin}} = \frac{1}{2}mv^2 = \frac{1}{2}m(\ell\dot{x})^2$$



kinetic energy

$$E_{\text{pot}} = \text{Const} + mgh = \underbrace{\text{Const} - mg\ell \cos x}_{\text{potential energy}}$$

$$\begin{aligned}\ddot{\theta} &= \ddot{E} = \frac{d}{dt}(E_{\text{kin}} + E_{\text{pot}}) \\ &= m\ell^2 \ddot{x} \dot{x}^2 + mg\ell \sin x \dot{x}\end{aligned}$$

Divide by  $m\ell^2 \dot{x}$

$$\ddot{\theta} = \ddot{x} + \frac{g}{\ell} \sin x$$



no explicit formulas  
for the solutions known

phase portrait  
of pendulum

## damped pendulum

$$\ddot{x} + r \dot{x} + \omega^2 \sin x = 0$$

friction term,  $r > 0$

For the damped pendulum, energy dissipates:

$$\begin{aligned}
 \dot{E} &= \frac{d}{dt}(E_{kin} + E_{pot}) = \frac{d}{dt}\left(\frac{1}{2}m(\ell\dot{x})^2 + (m\ell - mg\ell \cos x)\right) \\
 &= m\ell^2 \ddot{x} \dot{x} + mgl \sin x \dot{x} \\
 &= m\ell^2 \dot{x} \left( \ddot{x} + \left(\frac{g}{\ell} \sin x\right) \right) \\
 &= m\ell^2 \dot{x} (-r\dot{x}) = -m\ell^2 g r \dot{x}^2 \leq 0
 \end{aligned}$$

NB The damped harmonic oscillator can be solved explicitly.

$$\ddot{x} + r\dot{x} + \omega^2 x = 0$$

## Ansatz:

$$x(t) = e^{\lambda t}$$

$$\dot{x}(t) = \lambda e^{\lambda t} \quad x_r$$

$$\ddot{x}(t) = \lambda^2 e^{\lambda t}$$

$$(\omega^2 + r\lambda + \lambda^2) e^{\lambda t} = 0$$

$$\text{Hence } \lambda = -\frac{r}{2} \pm \sqrt{\left(\frac{r}{2}\right)^2 - \omega^2}, \quad \text{with solution}$$

$$x(t) = A e^{-\frac{r}{2} + i \sqrt{\omega^2 - \left(\frac{r}{2}\right)^2} t} + B e^{-\frac{r}{2} - i \sqrt{\omega^2 - \left(\frac{r}{2}\right)^2} t}$$

For small friction:  $\frac{r}{2} < |\omega|$

$$= e^{-\frac{rt}{2}} \left( A' \cos \sqrt{\omega^2 - \left(\frac{r}{2}\right)^2} t + B' \sin \sqrt{\omega^2 - \left(\frac{r}{2}\right)^2} t \right)$$

## Driven oscillators

$$\ddot{x} + r\dot{x} + \omega^2 x = A \cos t$$

Friction term
driving force  
(also oscillating)

describes the motion of a child on a swing pushed by its parent, but also the behaviour of a radio antenna.

For simplicity, we give it a proper linear term (i.e. not  $\omega^2 \sin x$ ), so this is a 2<sup>nd</sup> order inhomogeneous linear ODE.

The homogeneous solution (with RHS = 0) was just given. To solve the entire ODE we have to find a particular solution and add it to the homogeneous solution.

$$\text{Ansatz } x_{\text{par}}(t) = p \cos t + q \sin t \quad \times \omega^2$$

$$\dot{x}_{\text{par}}(t) = -p \sin t + q \cos t \quad \times r$$

$$\ddot{x}_{\text{par}}(t) = -p \cos t - q \sin t \quad \times 1$$

$$A \cos t = (\omega^2 p + rq - p) \cos t + (\omega^2 q - rp - q) \sin t$$

$$\Rightarrow \begin{cases} q = \frac{rp}{\omega^2 - 1} \\ p = \frac{A - rq}{\omega^2 - 1} \end{cases}$$

Driven oscillator continued

$$\ddot{x} + r\dot{x} + \omega^2 x = A \cos t$$

$$\left\{ \begin{array}{l} q = \frac{rp}{\omega^2 - 1} \\ p = \frac{A - rq}{\omega^2 - 1} \end{array} \right. \Rightarrow p = \frac{A}{\omega^2 - 1 + r^2}$$

Hence the amplitude of the particular solution  $x_{\text{par}}$  is very large if both friction  $r \approx 0$  and  $\omega^2 \approx 1$  resonance.

In the extreme case of a frictionless driven oscillator:  $\ddot{x} + \omega^2 x = A \cos t$  the same method with adjusted Ansatz gives:

$$\text{Ansatz: } x_{\text{par}}(t) = p \sin t + qt \sin t \quad \times \omega?$$

$$\dot{x}_{\text{par}}(t) = p \cos t + qs \in t + qt \cos t \quad \times 0$$

$$\ddot{x}_{\text{par}}(t) = -ps \in t + 2q \cos t - qt \sin t \quad \times 1$$

$$A \cos t = 2q \cos t$$

$$\text{so } x_{\text{par}}(t) = \frac{At}{2} \cos t$$

but this solution is unbounded over time!

## Coupled oscillators

$$\ddot{x}_1 + \omega_1^2 x_1 = \varepsilon f_1(x_2)$$

$$\ddot{x}_2 + \omega_2^2 x_2 = \varepsilon f_2(x_1)$$

Naturally there can be more than two oscillators and friction terms are left out for simplicity

If there is no coupling i.e.  $\varepsilon = 0$  then the solutions are

$$\begin{cases} x_1(t) = A_1 \cos(t + t_1) \\ x_2(t) = A_2 \cos(t + t_2) \end{cases}$$

Consider a Poincaré section  $\Sigma = \{x_1 = A_1\}$ ,

then  $(x_1(t), x_2(t)) \in \Sigma$  if  $t = \frac{2\pi k}{\omega_1} - t_1$ ,  $k \in \mathbb{Z}$

and then  $x_2(t) = A_2 \cos(\underbrace{2\pi k \frac{\omega_2}{\omega_1} - \omega_2 t_1 + t_2}_{u_k})$

i.e.  $x_2(t) = A_2 \cos(u_k)$  for  $u_k = R_\alpha^k(u_0)$   $u_0 = t_2 - \omega_2 t_1$

and  $R_\alpha$  is the rotation with angle  $\alpha = \frac{\omega_2}{\omega_1}$  (usually  $\notin \mathbb{Q}$ )

Also assume  $\omega_2 \approx \omega_1$ , so

$$\frac{\omega_2}{\omega_1} \bmod 1 \equiv \delta \approx 0$$

If there is coupling, i.e.  $\varepsilon > 0$ ,


then  $R_\alpha$  becomes  $R_{\alpha, \varepsilon}$ ,

some nonlinear perturbation

of  $R_\alpha$ . If  $\delta$  is small

compared to  $\varepsilon$ , then  $R_{\alpha, \varepsilon}$

is likely to have a fixed point



fixed point:  
phase locking

time dependent,  
usually periodic

-7-

## Oscillators with parametric resonance

$$\ddot{x} + \omega^2(t) \sin x = 0$$

This models the behaviour of a person on a swing keeping it swinging by shifting his body back and forth.

This can make the stationary point  $x(t) \equiv 0$  unstable and the stationary point  $x(t) \equiv \pi$  stable.

