

# Exactness and Maximal Automorphic Factors of Unimodal Interval Maps

Henk Bruin

Mathematics Department 253-37  
California Institute of Technology, Pasadena CA 91125

Jane Hawkins

Mathematics Department CB #3250  
University of North Carolina at Chapel Hill, NC 27599

August 26, 2003

## Abstract

We study exactness and maximal automorphic factors of  $C^3$  unimodal maps of the interval. We show that for a large class of infinitely renormalizable maps, the maximal automorphic factor is an odometer with an ergodic nonsingular measure. We give conditions under which maps with absorbing Cantor sets have an irrational rotation on a circle as a maximal automorphic factor, as well as giving exact examples of this type. We also prove that every  $C^3$   $S$ -unimodal map with no attractor is exact with respect to Lebesgue measure. Additional results about measurable attractors in locally compact metric spaces are given.

## 1 Introduction

The notion of exactness of a noninvertible map was first introduced by Rohlin who proved, in the measure preserving case, that exact endomorphisms have no nontrivial measurable factors with zero entropy [21]. He showed that some piecewise monotone and continuous interval maps described by Renyi in [20] were exact. In this paper we address the question of exactness for smooth maps of an interval and study the structure of many nonexact interval maps as well as give sufficient conditions for  $S$ -unimodal maps to be Lebesgue exact.

It is well-known that every exact measure preserving endomorphism has positive measure theoretic entropy. The notion of exactness of an endomorphism extends easily to non-measure-preserving, nonsingular maps even though a satisfactory definition of nonsingular entropy is still elusive. A nonsingular map  $T$  of a measure space  $(X, \mathcal{B}, \mu)$  is *exact* if the intersection  $\bigcap_{n \geq 0} T^{-n} \mathcal{B}$ , called the *tail field* of  $\mathcal{B}$ , contains only sets of full measure or measure 0. The maximal automorphic factor of  $T$  is its induced action on the tail field; it is the trivial map on a one-point space if and only if  $T$  is exact.

In this paper, the main dynamical systems of interest are smooth unimodal maps of the interval. Throughout, the measure of interest is one-dimensional normalized Lebesgue measure  $\lambda$ . We show that a variety of measurable automorphic behavior is exhibited by these maps; it is well-known that the presence of a measure theoretic attractor of Lebesgue measure zero forces the map to be dissipative with respect to  $\lambda$ , but does not preclude the existence of conservative Lebesgue factors.

After giving a brief review of nonsingular ergodic theory and preliminary definitions, we prove some results about dynamical systems with measurable attractors. We show that every ergodic nonsingular dynamical system of a locally compact metric space with an attractor of measure zero admits an equivalent invariant infinite  $\sigma$ -finite measure.

We then turn to unimodal maps of the interval for examples. We prove that every  $C^3$   $S$ -unimodal map with no attractor is exact with respect to  $\lambda$ . Next, we consider examples with measure theoretic attractors which have been studied and classified by several authors [2, 11, 14]. We exhibit the easily identified maximal automorphic factor in the presence of either a stable periodic orbit or a cycle of intervals (finitely renormalizable). In the case of an infinitely renormalizable map (of bounded type), we show that it is a dissipative map with maximal automorphic factor isomorphic to an odometer. The factor measure on the odometer is nonsingular but not necessarily invariant.

Finally we discuss the case of maps which have absorbing Cantor sets. These maps were shown to exist in [5]. We give conditions under which the maximal automorphic factor is an irrational rotation on a circle as well as an exact example. Irrational rotation factors, as topological factors were shown to exist in [6]. The authors would like to thank Gerhard Keller for useful remarks and corrections on an earlier version of this paper.

## 2 Preliminaries

We assume throughout this paper that  $(X, \mathcal{B}, \mu)$  is a locally compact metric space with metric  $d$ , Borel  $\sigma$ -algebra  $\mathcal{B}$  on  $X$  and  $\mu$  a regular Borel probability measure on  $\mathcal{B}$ . Infinite measures are always assumed to be  $\sigma$ -finite. We assume that  $T$  is *nonsingular*;

i.e.,  $T : X \rightarrow X$  satisfies:  $\mu(A) = 0 \iff \mu(T^{-1}A) = 0$  for every  $A \in \mathcal{B}$ . We also assume that every point in  $X$  has at most countably many preimages under  $T$ . Furthermore in all of our examples we will assume without loss of generality that  $T$  is forward nonsingular as well; i.e., that  $\mu(A) = 0 \iff \mu(TA) = 0$  for all measurable sets  $A$ . For example, any  $C^1$  map of a manifold onto itself whose differential is nonvanishing except at finitely many points is forward and backward nonsingular with respect to the Riemannian volume form (locally equivalent to Lebesgue measure). Let  $\mathcal{B}_+ \subset \mathcal{B}$  denote the collection of measurable sets of positive measure. In order to stress the presence of both a topology and a Borel measurable structure, we will refer to  $(X, \mathcal{B}, \mu, T)$  as a *nonsingular dynamical system*.

**Definition 2.1.** *Let  $(X, \mathcal{B}, \mu, T)$  be a nonsingular dynamical system. The nonsingular dynamical system  $(Y, \mathcal{C}, \nu, S)$  is a (measurable) factor of  $(X, \mathcal{B}, \mu, T)$  if there exists a surjective measurable map  $\pi : X \rightarrow Y$  such that  $S \circ \pi(x) = \pi \circ T(x)$   $\mu$ -a.e., and  $\nu \sim \mu \circ \pi^{-1}$ .*

A sub- $\sigma$ -algebra  $\mathcal{B}_o \subset \mathcal{B}$  is  $T$ -invariant if  $T^{-1}\mathcal{B}_o \subset \mathcal{B}_o$ . It is well-known that every factor map gives rise to a  $T$ -invariant sub- $\sigma$ -algebra,  $\{\pi^{-1}C\}_{C \in \mathcal{C}} \subset \mathcal{B}$ , and conversely. We refer the reader to Rohlin [21] for details.

## 2.1 Nonsingular Measure Theory

We review some basic definitions used in nonsingular measure theory; when the map in question is neither invertible nor measure-preserving, ergodic properties need careful definitions. Some equivalent definitions are no longer equivalent in this setting and others simply do not extend. (For example, it is still an open question as to what the definition of mixing should be for nonsingular noninvertible maps [1]). We assume  $(X, \mathcal{B}, \mu, T)$  is a nonsingular dynamical system though these notions apply in more general measure theoretic settings.

A measurable set  $W$  is (*backward*) *wandering* if the sets  $\{T^{-n}W\}_{n=0}^{\infty}$  are all disjoint. Equivalently, no point in a wandering set  $W$  ever returns to  $W$ . A measurable set  $V$  is *forward wandering* if the sets  $\{T^n V\}_{n=0}^{\infty}$  are all disjoint. Every forward wandering set is also (backward) wandering, but the converse is not true. If  $T$  is invertible the concepts are identical. We will use the usual convention that wandering set always refers to a backward wandering set.

The map  $T$  is *conservative* if there exist no wandering sets of positive measure. There exists a maximal set  $C$  on which  $T$  is conservative, and  $C \subset T^{-1}C$ . A non-conservative map is called *dissipative*; if  $T$  is not conservative on any set of positive measure, then  $T$  is *completely dissipative*, and we can write  $X$  as the (at most countable) union of wandering sets up to a set of measure 0.

The map  $T$  is *ergodic* if  $T$  has a trivial field of invariant sets, or equivalently, if any measurable set  $B$  with the property that  $\mu(B \Delta T^{-1}B) = 0$  has either zero or full measure.

It follows from the definitions that  $T$  is conservative and ergodic if and only if for all sets  $A, B \in \mathcal{B}_+$ , there is a positive integer  $n$  such that  $\mu(B \cap T^{-n}A) > 0$ .

A map is *exact* if it has a trivial tail field  $\bigcap_{n \geq 0} T^{-n}\mathcal{B} \subset \mathcal{B}$ , or equivalently, if any set  $B$  with the property  $\mu(T^{-n} \circ T^n(B) \Delta B) = 0$  for all  $n$  has either zero or full measure. For any set  $A \in \mathcal{B}_+$ , we define a tail set from it by:

$$\text{Tail}(A) := \bigcup_{n \in \mathbb{N}} T^{-n} \circ T^n(A).$$

Denoting the tail sets ( $\mu \bmod 0$ ) by  $\mathcal{T} \subset \mathcal{B}$ , we have  $\bigcap_{n \geq 0} T^{-n}\mathcal{B} = \mathcal{T} (\mu \bmod 0)$ . There is a natural factor mapping onto  $\mathcal{T}$  called the exact decomposition (of  $T$  with respect to  $\mu$ ), and  $T$  acts as an automorphism on the factor space. We denote the factor space by  $(Y, \mathcal{C}, \nu)$ , and the induced automorphism by  $S$ ; note that a point in  $Y$  is an atom of the measurable partition generated by the relation  $x \sim w \iff T^n x = T^n w$  for some  $n \in \mathbb{N}$ . We call this factor the *maximal automorphic factor*; this is because if there is a factor map  $\phi : X \rightarrow Z$  with induced factor automorphism  $R$ , then  $R$  is a factor of  $S$ . We remark that in general  $(Y, \mathcal{C}, S, \nu)$  is a nonsingular endomorphism of a Lebesgue space with no specified topology.

It is well-known that any invertible ergodic nonsingular transformation of a nonatomic measure space is conservative, and virtually all of the examples we consider below are ergodic, so their automorphic factors will be either conservative or atomic, or both. In the next result, we give a condition that rules out the dissipative possibility for an automorphic factor. We recall that every invertible, dissipative, ergodic, nonsingular transformation of a  $\sigma$ -finite space is isomorphic to  $x \mapsto x+1$  on  $\mathbb{Z}$  with an appropriately weighted counting measure.

**Lemma 2.1.** *If  $(X, \mathcal{B}, \mu, T)$  is a nonsingular dynamical system, and if  $\mathcal{B}_+$  contains no forward wandering sets, then every automorphic factor of  $T$  is conservative.*

*Proof.* The trivial factor is conservative, so we assume that  $T$  has a nontrivial automorphic factor. We will denote the projection onto the automorphic factor by  $\pi$ , the factor space by  $Y$ , and the induced automorphism by  $S$ . Then if  $S$  is not conservative, there exists a wandering set  $W$  of positive measure. Since  $S$  is an automorphism, the sets  $\{S^{-n}W\}_{n \in \mathbb{Z}}$  are all disjoint (in  $Y$ ). Then, by definition of the factor, the sets  $\pi^{-1}(S^{-n}W)$  are also disjoint and equal to the (disjoint) collection of sets  $\{T^{-n}(\pi^{-1}W)\}_{n \in \mathbb{Z}}$  (in  $X$ ). This contradicts the hypothesis since  $\pi^{-1}W$  has positive measure and is forward wandering.  $\square$

**Remark:** One can construct examples of ergodic dissipative maps with conservative factors. If we consider the product of a K-automorphism  $S$  with the dissipative ergodic endomorphism  $R(n) = n + 1$  on  $\mathbb{N}$  with counting measure, then the map  $T = S \times R$  is dissipative and ergodic with a conservative automorphic factor (which is  $S$ ). Below we will show that such examples occur within the family of S-unimodal maps.

The following result is well-known but we include it for completeness.

**Lemma 2.2.** *If  $T$  is nonsingular and exact, then  $T$  is totally ergodic; i.e., for each  $n \in \mathbb{N}$ ,  $T^n$  is ergodic.*

We give a necessary and sufficient condition for exactness which will be useful in the context of interval maps.

**Proposition 2.1.** *An ergodic nonsingular endomorphism  $T$  is exact on  $(X, \mathcal{B}, \mu)$  if and only if for every set  $B \in \mathcal{B}_+$ ,  $\mu(T^{-n} \circ T^{n+1}(B) \cap B) > 0$  for some  $n \in \mathbb{N}$ .*

*Proof.* Assume first that  $T$  is exact and  $A \in \mathcal{B}_+$  with  $\mu(T^{-n} \circ T^{n+1}(A) \cap A) = 0$  for every  $n \in \mathbb{N}$ . This means that  $\mu(\text{Tail}(TA) \cap A) = 0$ , but this is impossible since  $\mu(\text{Tail}(TA)) = 1$  by exactness and nonsingularity. To prove the other direction let  $(Y, \mathcal{C}, \nu, S)$  be the maximal automorphic factor of  $(X, \mathcal{B}, \mu, T)$ ,  $\pi : X \rightarrow Y$  being the factor map. Assume by contradiction that  $T$  is not exact, i.e.  $(Y, \mathcal{B}, \nu, S)$  is nontrivial. Then there exists  $C \in \mathcal{C}$  such that  $\nu(C) > 0$  and  $C \cap S(C) = \emptyset$ . Take  $B := \pi^{-1}(C)$ . Since the maximal automorphic factor is isomorphic to the tail field of  $(X, \mathcal{B}, \mu, T)$ ,  $B$  satisfies  $T^{-n} \circ T^n(B) = B$  ( $\mu$  mod 0) for all  $n$ . The same thing is true for  $T(B) = \pi^{-1} \circ S(C)$ . By assumption  $0 < \mu(T^{-n} \circ T^{n+1}(B) \cap B) = \mu(T^{-n} \circ T^n(T(B)) \cap B) = \mu(T(B) \cap B)$ . Hence  $0 < \nu(\pi(T(B) \cap B)) = \nu(S(C) \cap C)$ , contradicting the choice of  $C$ .  $\square$

### 3 Measurable Attractors

An important link between the topological and the measure theoretical dynamics occurs when there are attractors present. The definition of measurable attractor we give here was introduced by Milnor [18]. We assume that  $(X, \mathcal{B}, \mu, T)$  is a nonsingular dynamical system.

For any point  $x \in X$ , the omega limit set  $\omega(x)$  is defined as  $\omega(x) = \overline{\bigcap_n \bigcup_{i>n} T^i(x)}$ . With our standing assumptions on  $X$ , for each  $x \in X$ ,  $\omega(x)$  is a Borel measurable set.

**Definition 3.1.** *For a set  $\mathcal{A}$ , we define*

$$B(\mathcal{A}) := \{x \in X; \omega(x) \neq \emptyset, \omega(x) \subset \mathcal{A}\},$$

and call it the basin of  $\mathcal{A}$ . An attractor is a compact subset  $\mathcal{A} \subset X$  such that  $\mu(B(\mathcal{A})) > 0$ , and there is no proper subset  $\mathcal{A}' \subset \mathcal{A}$ , such that  $\mu(B(\mathcal{A}')) > 0$ .

Obviously, an attractor is invariant:  $T(\mathcal{A}) = \mathcal{A}$ . Milnor defined attractors to be closed, but as he was considering endomorphisms on compact manifolds and our space is only locally compact, we define attractors to be compact.

**Proposition 3.1.** *If  $(X, \mathcal{B}, \mu, T)$  is ergodic, there can be at most one attractor and  $\mu(X \setminus B(\mathcal{A})) = 0$ . Moreover, for any neighborhood  $U$  of  $\mathcal{A}$  and any  $x \in B(\mathcal{A})$ , there exists  $N$  such that  $T^n(x) \in U$  for all  $n \geq N$ .*

*Proof.* The basin of an attractor is clearly completely invariant and has positive measure, so it must have full measure.

It suffices to prove the second statement for small neighborhoods only. By continuity of  $T$ , we can take  $U \supset \mathcal{A}$  so small that  $T(U)$  is contained in a compact set  $K \subset X$ . If  $x \in B(\mathcal{A})$ , then there exists a sequence  $\{n_i\}$  such that  $T^{n_i}(x) \rightarrow \mathcal{A}$ . Suppose that  $T^n(x) \notin U$  infinitely often. Then there exists a sequence  $\{m_i\}$ ,  $m_i \geq n_i$ , such that  $T^{m_i}(x) \in U$  but  $T^{m_i+1}(x) \in K \setminus U$ . Because  $K \setminus U$  is compact, the sequence  $\{T^{m_i+1}(x)\}$  has an accumulation point  $y \in \omega(x) \setminus U$ , contradicting that  $\omega(x) \subset \mathcal{A}$ .  $\square$

Without loss of generality we can assume that  $\mu$  is a probability measure on  $X$  (by replacing  $\mu$  by an equivalent one if necessary). Typically we are interested in the case where  $\mu(\mathcal{A}) = 0$  and  $\mu(B(\mathcal{A})) = 1$ . In this case we can show that the map is completely dissipative.

**Proposition 3.2.** *If  $(X, \mathcal{B}, \mu, T)$  is an ergodic nonsingular dynamical system with an attractor  $\mathcal{A}$  satisfying  $\mu(\mathcal{A}) = 0$ , then  $T$  is completely dissipative.*

*Proof.* Suppose that there exists a set  $C \in \mathcal{B}_+$  on which  $T$  is conservative. By the regularity of  $\mu$ , we can find a compact set  $K \subset C$ ,  $\mu(K) > \mu(C)/2$ , and such that  $K$  does not intersect some neighborhood  $U$  of  $\mathcal{A}$ . Conservativity on  $K$  implies that  $\mu$ -a.e.  $x \in K$  returns to  $K$  infinitely often, but the previous lemma shows that  $K$  is in the basin of  $\mathcal{A}$   $\mu$ -a.e., so  $\mu$ -a.e. point enters  $U$  and stays there. Hence no such  $K$  exists.  $\square$

### 3.1 Maps with Attractors have $\sigma$ -finite Measures

Suppose the nonsingular dynamical system  $(X, \mathcal{B}, \mu, T)$  has an attractor of measure 0. Even though  $T$  is completely dissipative, we show that there exists a  $\sigma$ -finite invariant measure equivalent to  $\mu$ .

Before giving the proof of the result, we review some properties of nonsingular countable-to-one maps. We assume that  $(X, \mathcal{B}, \mu)$  is a Borel probability space and

$T : X \rightarrow X$  is a nonsingular ergodic endomorphism which is surjective and countable-to-one almost everywhere. Since  $T$  is countable-to-one, we apply a well-known result of Rohlin [13] to obtain a measurable partition  $\zeta = \{A_1, A_2, A_3, \dots\}$  of  $X$  into at most countably many sets, called atoms, satisfying:

1.  $\mu(A_i) > 0$  for each  $i$ ;
2. the restriction of  $T$  to each  $A_i$ , which we will write as  $T_i$ , is one-to-one;
3. each  $A_i$  is of maximal measure in  $X \setminus \bigcup_{j < i} A_j$  with respect to property 2;
4.  $T_1$  is one-to-one and onto  $X$  (by numbering the atoms  $A_i$  so that

$$\mu(TA_i) \geq \mu(TA_{i+1})$$

for  $i \in \mathbb{N}$ ).

We call a partition  $\zeta$  of this form a *Rohlin partition*.

The map  $T_1^{-1} \circ T$  gives an obvious factor map from  $X$  onto  $A_1$ . Since  $T$  is countable-to-one, each fiber over  $x \in A_1$  contains at most countably many points (these are the points  $x = w_1, \dots, w_n, \dots$  such that  $T(x) = T(w_j)$ ) and there is an atomic probability measure  $\mu_x$  associated to each point  $x$ ,  $\mu_x^j, j = 1, \dots, n, \dots$ , which is just the factor decomposition of the measure  $\mu$ , viewed as a measure on  $X$  over the fibers of  $A_1$ . The measures  $\mu_x$  vary measurably in  $x$ .

We now turn to the main result of this section.

**Theorem 3.1.** *Let  $(X, \mathcal{B}, \mu, T)$  be an ergodic nonsingular dynamical system, where  $\mu$  is a  $\sigma$ -finite regular Borel measure and  $X$  is a metric space such that  $T$  is countable-to-one and  $T(X) = X$ . Assume that  $X$  has an attractor  $\mathcal{A}$  such that  $\mu(\mathcal{A}) = 0$ . Then there exists a  $\sigma$ -finite invariant measure  $\nu$  which is equivalent to  $\mu$ .*

*Proof.* As mentioned above, we can assume that  $\mu(X) = 1$ . We define the sets  $X_i$ ,  $i \geq 1$ , as follows:  $X_1 := X$ ,  $X_{2i} := \{x \in T(X_{2i-1}); d(T^n(x), \mathcal{A}) \leq 2^{-i} \text{ for all } n \geq 0\}$  and  $X_{2i+1} := T(X_{2i})$ . Obviously  $T(X_i) \supset X_{i+1}$  and because  $\mathcal{A}$  is an attractor,  $\mu(X_i) > 0$  for all  $i$ .

Let  $\hat{X}$  be the disjoint union  $\sqcup_{i \geq 1} X_i$  equipped with the action

$$\hat{T}(x \in X_i) = T(x) \in \begin{cases} X_{i+1} & \text{if } i \text{ is even,} \\ X_{i+1} & \text{if } i \text{ is odd and } T(x) \in X_{i+1}, \\ X_i & \text{if } i \text{ is odd and } T(x) \in X_i \setminus X_{i+1}. \end{cases}$$

If  $\pi_0 : \hat{X} \rightarrow X$  is the standard projection, then  $\pi_0 \circ \hat{T} = T \circ \pi_0$ . We note that each  $X_i$  can be viewed as a subset of  $X$  with the restriction measure  $\mu_i := \mu|_{X_i}$ . Then the nonsingularity and forward and backward measurability of  $T$  gives the corresponding properties for  $\hat{T}$  with respect to the measure

$$\rho(B) := \sum_i \mu_i(B \cap X_i).$$

By construction,  $X_{2i}$  is backward wandering in  $\hat{X}$  for each  $i$ . Moreover, for  $\rho$ -a.e.  $x \in \sqcup_{j \leq 2i} X_j$ , there exists a unique  $n \geq 0$  such that  $\hat{T}^n(x) \in X_{2i}$ .

We define a sequence of measures  $\hat{\mu}_n$  on  $\hat{X}$  starting with  $\hat{\mu}_1$ . Let  $\hat{\mu}_1|_{X_2} = \mu|_{X_2}$ , and inductively extend the measure to  $X_1 \sqcup X_2$  as follows. By our assumptions we can write  $X_1 = \cup_j \hat{T}^{-j}(X_2)$ , where the sets in the union are mutually disjoint and each has positive  $\rho$ -measure. By using the ergodicity of  $T$ , we can consider this disjoint union of  $X_1 := X$  as defining a measurable partition of  $X_1$ ,  $\mathcal{Q} = \{\hat{T}^{-1}(X_2), \hat{T}^{-2}(X_2) \dots\}$ . Therefore it is enough to define  $\hat{\mu}_1|_{\hat{T}^{-j}(X_2)}$ . We proceed inductively on  $j$ .

Consider any  $B \subset \hat{T}^{-1}X_2$  and, we suppose first that  $B = \hat{T}^{-1} \circ \hat{T}(B)$   $\rho$ -a.e.; i.e.,  $B$  is  $\hat{T}^{-1}\mathcal{B}$  measurable set. Then we define

$$\hat{\mu}_1(B) = \hat{\mu}_1(\hat{T}(B)),$$

In order to compensate for the fact that  $T$ , hence  $\hat{T}$  is not one-to-one everywhere (and therefore there are measurable sets not of the above form), we refine  $\mathcal{Q}$  by a Rohlin partition (for  $T$  on  $X$ )  $\zeta = \{A_1, A_2, \dots, A_n, \dots\}$ . We note that  $A_1$  intersects every atom of  $\mathcal{Q}$  in a set of positive measure since  $T$  maps  $A_1$  onto  $X$ . Define the sets

$$Q_k^j = A_k \cap \hat{T}^{-j}X_2,$$

for each  $j, k \in \mathbb{N}$ . Denote for each  $j, k$ , the relative size of each  $A_k$  in  $\hat{T}^{-j}X_2$ , by

$$\alpha_k^j = \frac{\mu(Q_k^j)}{\mu(\hat{T}^{-j}X_2)}.$$

Then for every  $j$ ,  $\sum_{k \geq 1} \alpha_k^j = 1$ . Write  $B_k := B \cap Q_k^1$  and define:

$$\hat{\mu}_1(B_k) = \alpha_k^1 \cdot \hat{\mu}_1 \circ \hat{T}(B_k) \text{ and } \hat{\mu}_1(B) = \sum_{k \geq 1} \hat{\mu}_1(B_k).$$

In particular,

$$\hat{\mu}_1(\hat{T}^{-1}X_2) = \sum_{k \geq 1} \hat{\mu}_1(Q_k^1) \leq \mu(X_2).$$



We give the inductive step. If  $\hat{\mu}_1$  is defined on  $\cup_{i < j} \hat{T}^{-i} X_2$ , we extend it to  $\hat{T}^{-j} X_2$  as follows. For  $B \subset \hat{T}^{-j} X_2$ ,  $B_k := B \cap Q_k^j$ , and  $\hat{T}(B_k) \subset \hat{T}^{-(j-1)} X_2$ , so we define:

$$\hat{\mu}_1(B_k) = \alpha_k^j \cdot \hat{\mu}_1 \circ \hat{T}(B_k).$$

Then, as before,

$$\hat{\mu}_1(B) = \sum_{k \geq 1} \hat{\mu}_1(B_k).$$

We extend the measure  $\hat{\mu}_1$  to all of  $\hat{X}$  by setting  $\hat{\mu}_1(B) = 0$  if  $B \subset \sqcup_{m > 2} X_m$ . Therefore  $\hat{\mu}_1(B) = \hat{\mu}_1(\hat{T}^{-1}(B))$  if  $B \subset X_1 \sqcup X_2$ , and  $\hat{\mu}_1(B) = 0$  if  $B \cap (X_1 \sqcup X_2) = \emptyset$ . Note that by construction  $\hat{\mu}_1(\hat{T}^{-j}(X_2)) = \mu(X_2)$  for every  $j$ , so  $\hat{\mu}_1$  is infinite and  $\sigma$ -finite.

We now continue inductively on  $n$  and define  $\hat{\mu}_{n+1}$  by

$$\hat{\mu}_{n+1} = \hat{\mu}_n \circ \hat{T}^{-1}.$$

Clearly  $\hat{\mu}_{n+1} \geq \hat{\mu}_n$ ; furthermore, if  $B \subset \sqcup_{j \leq n+1} X_j$  for any  $n \geq 1$  then  $\hat{\mu}_{n+1}(B) = \hat{\mu}_n(\hat{T}^{-1}(B)) = \hat{\mu}_n(B)$ , so equality holds.

We claim that  $\hat{X}$  admits a countable partition  $\{X_k^j\}_{j,k \geq 1}$  (up to a set of  $\rho$ -measure 0), where  $X_k^j$  is a measurable subset of  $X_k$  and  $\hat{\mu}_n(X_k^j) \leq \mu(X_2)$  for all  $j, k \geq 1$ . We then define the limit measure  $\hat{\mu}$  by

$$\hat{\mu}(B) = \sum_{j,k \geq 1} \lim_{n \rightarrow \infty} \hat{\mu}_n(B \cap X_k^j).$$

Because  $\hat{\mu}_n$  is invariant on  $\sqcup_{k \leq n} X_k$ , this limit is a  $\hat{T}$  invariant measure. Since  $\hat{\mu}(X_k^j) \leq \mu(X_2) < \infty$ ,  $\hat{\mu}$  is clearly  $\sigma$ -finite. Finally, on  $X$  the desired measure is  $\nu := \hat{\mu} \circ \pi_0^{-1}$ , which is  $T$ -invariant and absolutely continuous with respect to  $\mu$ . Moreover  $\hat{\mu}|_{X_1} \geq \hat{\mu}_1|_{X_1}$ , and due to the nonsingularity of  $T$  and the construction of  $\hat{\mu}_1$  above,  $\hat{\mu}_1|_{X_1}$ , viewed as a measure on  $X$ , is equivalent to  $\mu$ . Hence  $\nu$  is equivalent to  $\mu$ .

Ergodicity and invariance pass to any factor measure; however we must show that  $\nu$  is  $\sigma$ -finite since this property can fail when taking factors. Recall that  $\nu(\mathcal{A}) = 0$  and  $\mathcal{A} = \cap_i \cup_{j \geq i} X_j$ . Therefore, if  $A_\varepsilon$  is an  $\varepsilon$ -neighborhood of  $\mathcal{A}$ ,  $\pi_0^{-1}(X \setminus A_\varepsilon)$  intersects only finitely many levels  $X_j$ . Hence  $\nu|_{X \setminus A_\varepsilon}$  is  $\sigma$ -finite. Since this is true for all  $\varepsilon > 0$  and  $\mu$  is regular Borel,  $\nu$  is  $\sigma$ -finite.

It remains to define the sets  $X_k^j$  and prove the claim. For each  $k = 2i$  and  $j \geq 1$ , let

$$X_k^j = \hat{T}^{-j}(X_{k+2}) \cap X_k \text{ and } X_{k+1}^j = \hat{T}^{-j}(X_{k+2}) \cap X_{k+1}.$$

Because the  $X_k^j$  are preimages of backward wandering sets, they are pairwise disjoint. We claim that

$$\sup_n \hat{\mu}_n(X_{2i}) \leq \mu(X_2) \text{ for all } i.$$

By construction,  $\hat{\mu}_n(X_2) = \mu(X_2)$  for all  $n$ . We continue by induction on  $i$ . Because  $\hat{\mu}_n(\hat{T}^{-1}(B)) \geq \hat{\mu}_n(B)$ , it follows that

$$\hat{\mu}_n(X_{2i+2}) \leq \sum_j \hat{\mu}_n(X_{2i}^j) \leq \hat{\mu}_n(X_{2i}).$$

Furthermore,  $\hat{\mu}_n(X_k^j) \leq \hat{\mu}_n(X_{2i+2}) \leq \mu(X_2)$  for all  $j, i \geq 1$  and  $k = 2i$  or  $2i+1$ . Because  $\hat{\mu}(X_k \setminus \cup_j X_k^j) = 0$  for all  $k$ , this establishes the asserted partition of  $\hat{X}$ .  $\square$

## 4 S-unimodal Maps

We now consider a class of smooth maps of the interval as our main examples of nonsingular dynamical systems. Let  $f : I \rightarrow I$  be a unimodal map. By this we mean that there is a unique point  $c$ , called the *critical point*, such that  $f$  is increasing on the left and decreasing on right of  $c$ . The iterates  $f^i(c)$  will be denoted by  $c_i$ . Assuming that  $c_2 < c < c_1$ , we can scale  $f$  in such a way that  $I = [c_2, c_1]$ . For  $x \in I$ , let  $\tau(x) = \bar{x}$  be the point such that  $f^{-1} \circ f(x) = \{x, \bar{x}\}$ . Note that  $f$  is two-to-one on  $[c_2, \bar{c}_2] \setminus \{c\}$ . Therefore it is precisely on this set that  $x \neq \bar{x}$ , so that  $\tau$  is defined; we have  $\tau \circ \tau = id$  on its domain. We call a set  $A \subset [c_2, \bar{c}_2] \setminus \{c\}$  *symmetric* if  $\tau(A) = A$ .

Throughout the paper we will assume that  $f$  is  $C^3$  and has negative Schwarzian derivative ( $f$  is *S-unimodal*), i.e.  $\frac{f'''}{f'} - \frac{3}{2}(\frac{f''}{f'})^2 \leq 0$  wherever defined. This assumption will enable us to make certain distortion estimates. We call a periodic point of period  $n$  *stable* if  $|(f^n)'(p)| \leq 1$ , thus comprising the hyperbolic attracting and neutral case. Under the assumption of negative Schwarzian derivative, a stable periodic orbit must attract the critical point. Hence there can be only one such orbit.

The *critical order* of  $f$  is  $\ell$  if there exists  $M > 0$  such that  $\frac{\ell|x-c|^{\ell-1}}{M} \leq |f'(x)| \leq M\ell|x-c|^{\ell-1}$  for all  $x \in I$ . For some of our results it is important that  $f$  is *nonflat*, i.e.  $\ell < \infty$ . The critical omega limit set,  $\omega(c)$ , is of particular importance. If  $f$  is nonflat S-unimodal, then either  $\omega(c)$  contains an interval or has Lebesgue measure 0, see [14].

The ergodic properties of Lebesgue measure, which we will denote by  $\lambda$ , for unimodal maps are well-understood. We quote a result by Blokh and Lyubich [2]:

**Proposition 4.1.** *Suppose  $f$  is  $C^3$  S-unimodal and has no stable periodic orbit. Then any forward invariant measurable symmetric set  $A$  (i.e.  $\tau(A) = A$ ) with  $\lambda(A) > 0$  has  $c$  as density point.*

From this, one can easily derive that  $f$  is Lebesgue ergodic if and only if there is no stable periodic orbit. Moreover  $\lambda$ -a.e. point accumulates on the critical point in this case.

**Theorem 4.1.** [2, 11, 14] *If  $f$  is  $C^3$   $S$ -unimodal, then  $f$  is conservative if and only if there is no attractor. There is at most one attractor, which is of one of the following types:*

1. *a stable periodic orbit,*
2. *a cycle of intervals (the finitely renormalizable case),*
3. *a Cantor set (the infinitely renormalizable case),*
4. *an absorbing Cantor set.*

*In the cases 1, 3, and 4,  $\mathcal{A}$  necessarily equals  $\omega(c)$ , and  $f$  is completely dissipative.*

A map is *renormalizable* if there exists a interval  $J$ ,  $c \in J \neq I$ , such that  $f^n(J) \subset J$  and  $J, f(J), \dots, f^{n-1}(J)$  have disjoint interiors. As a rule, we take  $J$  minimal with this property; such a  $J$  is called a *restrictive interval*. The map  $f^n|_J$  is again a unimodal map; it is called the renormalization of  $f$ . If there is a smallest restrictive interval  $J$ , then  $f$  is finitely renormalizable. If there are arbitrarily small restrictive intervals (of arbitrarily large period) then  $f$  is *infinitely renormalizable*. (The best known example is the Feigenbaum map; here the periods of the restrictive intervals are the powers of 2.) The attractor is a Cantor set. It attracts all points except for a nullset of first Baire category. No point in  $I$  has a dense forward orbit. By Lemma 2.2, a renormalizable map cannot be exact.

In case 4,  $f$  is not infinitely renormalizable, but  $\omega(c)$  is a Cantor set such that  $\omega(x) \subset \omega(c)$  for  $x$  in a full measure set of first Baire category. There is a second category set of points whose orbit lies dense in  $I$  (or if  $f$  is finitely renormalizable in a cycle of intervals). It was shown in [5] that so-called Fibonacci unimodal maps with sufficiently large critical orders have absorbing Cantor sets. Theorem 3.1 applied to unimodal maps yields the following result.

**Proposition 4.2.** *If  $f$  is a nonflat  $C^3$   $S$ -unimodal map with an attractor of type 1, 3 or 4, then  $f$  admits a (dissipative)  $\sigma$ -finite invariant measure which is absolutely continuous with respect to  $\lambda$ .*

This result is known when  $\mathcal{A}$  is a stable periodic orbit, or when  $\mathcal{A}$  is an absorbing Cantor set [16]. To our knowledge, the result is new for infinitely renormalizable maps. We conjecture that the result does not hold for Misiurewicz' example of a  $C^\infty$  Feigenbaum map whose attractor  $\omega(c)$  has positive Lebesgue measure [19]. He showed that  $(\omega(c), f)$  admits no  $\sigma$ -finite absolutely continuous invariant measure, and it seems likely that  $\lambda(I \setminus \cup_n f^{-n}(\omega(c))) = 0$ .

## 4.1 Exact S-unimodal Maps

In this section we prove some results concerning exactness and automorphic factors for general S-unimodal maps. Let  $f$  be an S-unimodal map; we show that if  $f$  has no attractor then  $f$  is Lebesgue exact. We also demonstrate the maximal automorphic factor of  $f$  in the cases where either  $f$  has a stable periodic orbit or an attracting cycle of intervals. In later sections we deal with certain specific (dissipative) unimodal maps from the remaining two cases; i.e., where  $f$  has an attractor of type 3 or 4.

The first result is already known in the case where  $f$  admits an absolutely continuous invariant measure  $\mu$  and is nonrenormalizable. Ledrappier [12] showed that  $f$  has a Bernoulli natural extension in this case and is exact with respect to  $\mu \ll \lambda$ .

**Theorem 4.2.** *Let  $f$  be  $C^3$  S-unimodal. If  $f$  has no attractor, then  $f$  is Lebesgue exact.*

Our proof relies on a result by Martens and involves a property which he calls the strong Markov property [15, 14, 17].

**Proposition 4.3 (Strong Markov Property).** *Suppose  $f$  has no attractor of type 1, 3, or 4. Then there exist symmetric neighborhoods  $U$  and  $V$  of  $c$ ,  $c \in \text{cl}U \subset \text{int}V$ , such that for  $\lambda$ -a.e.  $x$  the following holds: There exist integers  $k_1(x) < k_2(x) < \dots$  and nested intervals  $x \in \dots \subset I_2(x) \subset I_1(x)$  such that for all  $i$ ,  $f^{k_i(x)}(x) \in U$  and  $f^{k_i(x)}$  maps  $I_i(x)$  monotonically onto  $V$ .*

**Remark.** Let  $L_1$  and  $L_2$  be the components of  $U \setminus V$ , and say  $\lambda(L_1) \leq \lambda(L_2)$ ; define  $\delta := \frac{\lambda(L_1)}{\lambda(V)}$ . Then the Koebe Principle (see [17, Theorem IV 1.3]) gives the distortion bound  $K(\delta) = (\frac{1+\delta}{\delta})^2$ :

$$\begin{aligned} \text{dist}(f^{k_i(x)}, I_i(x) \cap f^{-k_i(x)}(U)) &:= \\ \sup\left\{\frac{|Df^{k_i(x)}(y)|}{|Df^{k_i(x)}(z)|}; y, z \in I_i(x) \cap f^{-k_i(x)}(U)\right\} &\leq K(\delta). \end{aligned} \tag{1}$$

*Proof of Theorem 4.2:* Let  $U \subset V$  be as in Proposition 4.3 and let  $\delta > 0$  and distortion bound  $K = K(\delta) \geq 1$  be as in (1).

Let  $p$  be the orientation reversing fixed point of  $f$ . Without loss of generality, we can assume that  $V \subset (\bar{p}, p)$ . It is well-known (see e.g. [17, Theorem III 4.6]) that, since  $f$  is not renormalizable,  $\cup_n f^n(U) = I$ . Therefore there exists a minimal  $r > 0$  such that  $p \in \text{int}f^r(U)$ . Let  $H \subset f^r(U)$  be an interval such that  $f(H) \subset f^r(U)$ . Let  $H_1, H_2 \subset U$  be such that  $f^r$  maps  $H_1$  and  $H_2$  diffeomorphically onto  $H$  and  $f(H)$  respectively. Let  $K_0 = K_0(H_1, H_2)$  such that  $\text{dist}(f^{r+1}, H_1), \text{dist}(f^r, H_2) \leq K_0$ . Take  $\varepsilon := \min\{\lambda(H_1), \lambda(H_2)\}$ .

Now let  $A$  be any set of positive measure. Let  $x$  be a Lebesgue density point of  $A$  such that the integers  $k_i$  and intervals  $I_i(x)$  from Proposition 4.3 are well-defined. Since  $x$  is a density point, we can take  $i$  so large that  $\frac{\lambda(I_i(x) \setminus A)}{\lambda(I_i(x))} \leq \frac{\varepsilon}{3K_0}$ . By the Koebe Principle  $\text{dist}(f^{k_i}, I_i(x) \cap f^{-k_i}(U)) \leq K$ . Therefore

$$\frac{\lambda(H_1 \setminus f^{k_i}(A))}{\lambda(H_1)} \leq \frac{\lambda(U \setminus f^{k_i}(A))}{\lambda(U)} \frac{\lambda(U)}{\lambda(H_1)} \leq \frac{\varepsilon}{3K_0} \frac{1}{\varepsilon} \leq \frac{1}{3K_0}.$$

and similarly  $\frac{\lambda(H_2 \setminus f^{k_i}(A))}{\lambda(H_2)} \leq \frac{1}{3K_0}$ . Using the distortion bound  $K_0$  of  $f^{r+1}|_{H_1}$  and  $f^r|_{H_2}$ , we obtain

$$\frac{\lambda(f(H) \setminus f^{k_i+r+1}(A))}{\lambda(f(H))}, \frac{\lambda(f(H) \setminus f^{k_i+r}(A))}{\lambda(f(H))} \leq \frac{1}{3}.$$

It follows that  $\lambda(f^{k_i+r+1}(A) \cap f^{k_i+r}(A)) \geq \frac{1}{3}\lambda(f(H)) > 0$ . By Proposition 2.1,  $f$  is exact.  $\square$

## 4.2 Maps with Stable Periodic Orbits

We have shown that an S-unimodal map without an attractor has a trivial maximal automorphic factor. We next characterize the maximal automorphic factor of  $f$  if  $f$  has an attractor of type 1 or 2.

**Theorem 4.3.** *1. If  $(I, \mathcal{B}, \lambda, f)$  has a stable periodic orbit, then its maximal automorphic factor is isomorphic to  $(\mathbb{R}, \mathcal{B}, \lambda, x \mapsto x + 1)$ .*

*2. If  $(I, \mathcal{B}, \lambda, f)$  has an attractor  $J \cup f(J) \cup \dots \cup f^{n-1}(J)$  of type 2, then its maximal automorphic factor is isomorphic to  $(\mathbb{Z}/n\mathbb{Z}, i \mapsto i + 1 \pmod{n})$  with counting measure.*

*Proof.* Suppose  $p$  is an  $n$ -periodic stable periodic point with immediate basin  $B$ . Then  $\omega(x) = \text{orb}(p)$   $\lambda$ -a.e. If  $p$  reverses orientation, take  $b \in B \setminus \{p\}$  and  $U := (b, f^{2n}(b)]$ . If  $p$  preserves orientation, then take points  $b$  and  $b'$  in either component of  $B \setminus \{p\}$  and  $U := (b, f^n(b)] \cup [f^n(b'), b')$ . (If  $B \setminus \{p\}$  has only one component, namely because  $|(f^n)'(p)| = 1$ , then take  $U := (b, f^n(b)]$ .) In each case  $U$  is a forward wandering and fundamental in the sense that for  $\lambda$ -a.e.  $x \in I$ , there exist  $i, j \geq 0$  such that  $f^i(x) = f^j(y)$  for a unique  $y \in U$ . Define  $\pi(x) = (y, i - j)$ . Then it is easy to see that  $\pi : I \rightarrow U \times \mathbb{Z}$  is a factor map and  $(U \times \mathbb{Z}, (y, n) \mapsto (y, n + 1))$  is the maximal automorphic factor. This system is obviously isomorphic to  $(\mathbb{R}, \mathcal{B}, \lambda, x \mapsto x + 1)$ .

If  $f$  has an  $n$ -period restrictive interval  $J$ , then for  $\lambda$ -a.e.  $x$  there exists  $i$  such that  $f^i(x) \in J$ . The map  $\pi(x) = i \pmod{n}$  is obviously a factor map. If  $J \cup \dots \cup f^{n-1}(J)$  is the attractor, then by Theorem 4.2,  $f^n|_J$  is exact. Hence  $(\mathbb{Z}/n\mathbb{Z}, i \mapsto i + 1 \pmod{n})$  is also the maximal automorphic factor.  $\square$

The next result states that many dissipative unimodal maps have conservative maximal automorphic factors.

**Proposition 4.4.** *Let  $f$  be a  $C^3$  nonflat  $S$ -unimodal map having no stable periodic orbit. Then the maximal automorphic factor of  $(I, \mathcal{B}, \lambda, f)$  is conservative.*

*Proof.* A result by Blokh & Lyubich [2, Section 7] states that  $f$  does not admit wandering sets. (The proof was carried out for critical order  $\ell = 2$ , but can be generalized to any  $\ell \in (1, \infty)$ .) The proposition follows now from Lemma 2.1.  $\square$

### 4.3 Infinitely Renormalizable Maps

As mentioned earlier, a unimodal map is infinitely renormalizable if it has arbitrarily small restrictive intervals. We first review the structure of infinitely renormalizable maps in detail.

Let  $f$  denote an infinitely renormalizable  $S$ -unimodal map. Then  $f$  has a forward invariant Cantor set  $\Omega$ , and the following hold:

- There exists a decreasing chain of closed subsets of  $I$ , denoted by  $\Omega^k$  and satisfying

$$\dots \Omega^{k+1} \subset \Omega^k \dots \subset \Omega^1 \subset \Omega^0 = I;$$

each  $\Omega^k$  contains the critical point and each is mapped onto itself by  $f$ .

- There exists a sequence  $\{p_k\}_{k \in \mathbb{N}}$ , such that  $p_k$  divides  $p_{k+1}$ , such that the following hold. For each  $k \in \mathbb{N}$ , there exist  $p_k$  disjoint closed subintervals  $\Omega_{i_1, \dots, i_k}$ ,  $i_j \in \{0, 1\}$ , which are cyclically permuted by the first  $p_k - 1$  iterates of  $f$ , and such that  $f^{p_k}(\Omega_{i_1, \dots, i_k}) \subset \Omega_{i_1, \dots, i_k}$ ; furthermore  $\Omega^k = \cup \Omega_{i_1, \dots, i_k}$ .
- The critical point  $c$  always lies in the subinterval which is labelled  $\Omega_{0, 0, \dots, 0}$ ; i.e.,  $\Omega_{0, 0, \dots, 0}$  is a restrictive interval, and the rest are labelled so that the action of  $f$  moves the cylinders in the usual  $\{p_k\}$ -odometer order (add 1 to  $i_0$  and carry when necessary).
- The intervals are nested in the obvious way; that is,

$$\Omega_{i_1, \dots, i_{k-1}, i_k} \subset \Omega_{i_1, \dots, i_{k-1}}.$$

As  $f^{p_k}(\Omega_{i_1, \dots, i_k}) \subset \Omega_{i_1, \dots, i_k}$ , we can rescale  $f^{p_k}|_{\Omega_{i_1, \dots, i_k}}$  to a map on the unit interval. For this reason,  $f$  is called renormalizable. The fact that we can do this for every  $k$  makes  $f$  infinitely renormalizable.

- The intersection

$$\Omega := \bigcap_{k \geq 1} \Omega_{i_1, \dots, i_k} = \bigcap_{k \geq 1} \Omega^k$$

is known to be a Cantor set of Lebesgue measure 0 when  $f$  is nonflat S-unimodal (however cf. [19]). Furthermore,  $\Omega = \omega(c)$ , the  $\omega$ -limit set of the critical point, and  $f|_{\Omega}$  is the  $\{p_k\}$ -odometer.

- The orbit of  $\lambda$ -a.e. point  $x$  converges to the Cantor set  $\Omega$  in the sense that for every  $k$ , the orbit of  $x$  eventually lands inside  $\Omega^k$ ; i.e.,  $\Omega$  is an attractor. By  $\Omega^k(x)$  we will denote the specific subinterval  $\Omega_{i_1, i_2, \dots, i_k}$  containing  $x$ . Therefore  $\Omega^k(x)$  is defined for all  $k$  for  $\lambda$ -a.e.  $x$ .

If  $p_k = 2^k$  for an infinitely renormalizable S-unimodal map of the interval, then  $f$  is commonly known as a *Feigenbaum map*. Independently, Feigenbaum [8] and Coulet and Tresser [7] discovered this pattern of renormalization for these maps.

**Lemma 4.1.** *If  $f$  is S-unimodal and infinitely renormalizable, for  $\lambda$ -a.e.  $x \in I$  there exists a unique  $y \in \Omega$  such that  $|f^n(x) - f^n(y)| \rightarrow 0$  as  $n \rightarrow \infty$ . In this case we say that  $x$  copies  $y \in \Omega$ .*

*Proof.* Using the subintervals  $\Omega_{i_1, \dots, i_k}$  as a basis for the topology of the Cantor set, we have that the sequences  $(i_1, \dots, i_k, \dots)$  from the corresponding basis elements give a  $\{p_k\}$ -adic coding for each  $y \in \Omega$ , and, as mentioned above, the labeling is chosen to correspond under the action of  $f$  to the usual odometer action.

We fix some  $x \in I$  such that  $\omega(x) \subset \Omega$ . For each  $k \geq 0$ , there exists a smallest positive integer  $n_k$  such that  $f^{n_k}(x) \in J_k = \Omega_{0, \dots, 0}$  ( $k$  zeroes). As  $f^n(x) \in J_k$  only if  $n = n_k + jp_k$  for some  $j \geq 0$ , it follows that  $p_k$  divides  $n_{k+1} - n_k$ . Let  $i_1, \dots, i_{k-1}$  be such that  $n_k + \sum_{j=1}^{k-1} i_j p_j \equiv 0 \pmod{p_k}$ . Then  $f^{n_k}(x) \in f^{n_k}(\Omega_{i_1, \dots, i_{k-1}})$ . Furthermore,  $\Omega_{i_1, \dots, i_{k-1}} \supset \Omega_{i_1, \dots, i_k}$  for each  $k$ . Let  $y = \bigcap_k \Omega_{i_1, \dots, i_k}$ . We will show that  $x$  copies  $y$  and no other point.

For  $\varepsilon > 0$ , let  $k$  be so large that each component of  $\Omega^k$  has length  $< \varepsilon$ . Then  $|f^n(x) - f^n(y)| \leq \text{diam}(f^{n-n_k}(J_k)) < \varepsilon$  for all  $n \geq n_k$ . Since this holds for any  $\varepsilon > 0$ ,  $|f^n(x) - f^n(y)| \rightarrow 0$ . On the other hand,  $f|_{\Omega}$  is distal: If  $y' \in \Omega$ ,  $y' \neq y$ , then  $|f^n(y) - f^n(y')|$  is bounded away from 0 uniformly in  $n$ . Indeed, for  $k$  sufficiently large,  $y$  and  $y'$  are contained in different components of  $\Omega^k$ , and these components remain disjoint under iteration of  $f$ . Hence  $y$  is the unique point copied by  $x$ .  $\square$

Define the map  $\pi : I \rightarrow \Omega$  by  $\pi(x) = y$ ; i.e.,  $x$  is mapped to the unique point that it copies. This map is well-defined except on a set of  $\lambda$  measure zero. Using  $g$  for the odometer action on  $\Omega$ , we have  $\pi \circ f(x) = g \circ \pi(x)$ . It is easy to see from the dynamics of the map that the usual Borel structure generated by cylinder sets on  $\Omega$  agrees with

the factor measure structure (i.e., a set  $C$  is Borel in  $\Omega \iff \pi^{-1}C$  is Borel in  $I$ .) With respect to the factor measure  $\mu(\cdot) = \lambda \circ \pi^{-1}(\cdot)$ , the factor map is invertible. We take the completion of the Borel sets in  $\Omega$  with respect to  $\mu$  to obtain a factor Lebesgue space. Note that as a factor space, the space  $\Omega$  has  $\mu$  measure 1 (in contrast with the 0 Lebesgue measure it has as a subset of  $I$ ). The aim of the rest of this section is to show that  $(\Omega, \pi(\mathcal{B}), \mu, g)$  is the maximal automorphic factor for a certain class of infinitely renormalizable maps.

We have defined  $J_k = \Omega_{0\dots 0}$  ( $k$  zeroes) to be the  $k$ -th restrictive interval; its period is  $p_k$ . For each  $x \in B(\Omega)$ , let

$$n_k(x) := \min\{n; f^n(x) \in J_k\}$$

and

$$m_k(x) := \min\{m; f^{p_{k-1}m}(y) \in J_k \text{ for } y = f^{n_{k-1}}(x)\}.$$

In this way (taking  $p_0 = 1$ ) we obtain  $n_k = \sum_{i=1}^k m_k p_{k-1}$ . Basically,  $m_k$  plays the same role for  $f^{p_{k-1}}|_{J_{k-1}}$  as  $m_1$  plays for  $f|_I$ .

A unimodal map  $f$  is infinitely renormalizable of *bounded type* if it is infinitely renormalizable and the sequence the quotients  $p_k/p_{k-1}$  is bounded. The geometry of the Cantor set  $\Omega$  has been particularly well-studied for these maps, see [17, Chapter VI]. For our purposes we need the following facts.

**Proposition 4.5.** *1. Let  $f$  be a nonflat S-unimodal infinitely renormalizable map. Then there exists  $K \geq 1$  such that for all restrictive intervals  $J_k$  the following holds: if  $f^n : J \rightarrow J_k$  is a branch of the first return map to  $J_k$ , then the distortion  $\text{dist}(f^n, J) \leq K$ . (In particular  $f^n : J \rightarrow J_k$  is a diffeomorphism; if  $J = J_k$  then we should take  $n = 0$ .)*

*2. If in addition  $f$  is renormalizable of bounded type, then there exist  $\rho_- < \rho_+ < 1$  such that  $\rho_-^{n+1} \lambda(J_k) \leq \lambda(\{x \in J_{k-1}; m_k(x) = n\}) \leq \rho_+^{n+1} \lambda(J_k)$ .*

*Proof.* We only sketch the proof, since most of the details can be found in [17, Section VI.2]. In particular, it is shown that the central gaps of the  $k$ -th level, i.e. the components of  $I \setminus \Omega^k$  adjacent to the restrictive interval  $J_k$ , have size  $\geq \sigma |J_k|$  for some uniform constant  $\sigma > 0$ . (The proof in [17] of this fact deals with infinitely renormalizable maps of bounded type, but it remains valid for arbitrary non-flat S-unimodal infinitely renormalizable maps.) Let  $J$  be any maximal interval such that  $f^n : J \rightarrow J_k$  is a diffeomorphism. Let  $T \supset J$  be the maximal interval such that  $f^n|_T$  is a diffeomorphism. Then there exist  $a < b < n$  such that  $c \in \partial f^a(T), \partial f^b(T)$ , and  $f^n(T) = (c_{n-a}, c_{n-b}) \supset J_k$ . Because  $n - a$  and  $n - b \neq n$ ,  $c_{n-a}$  and  $c_{n-b}$  lie in  $\Omega^k \setminus J_k$ . Hence  $f^n(T)$  contains both gaps adjacent to  $J_k$ . Therefore the components of  $f^n(T) \setminus J_k$



both have size  $\geq \sigma|J_k|$ . This is the space needed to apply the Koebe Principle that yields the distortion bound given in formula (1).

For the second statement we remark that the dynamics of  $f^{p_{k-1}}$  on  $J_{k-1} \setminus J_k$  are hyperbolic (see e.g. [17, Theorem III.5.1]). More precisely, there exist  $\tau > 1$  and  $C > 0$  depending only on  $\lambda(J_k)/\lambda(J_{k-1})$ , such that  $|Df^{np_{k-1}}(x)| \geq C\tau^n$  whenever  $f^{jp_{k-1}}(x) \in J_{k-1} \setminus J_k$  for  $0 \leq j < n$ . If  $f$  is infinitely renormalizable of bounded type,  $\lambda(J_k)/\lambda(J_{k-1})$  is bounded uniformly away from 0. This implies that the assertion holds for a uniform choice of  $\rho_-$  and  $\rho_+$ .  $\square$

We now state the main result of this section.

**Theorem 4.4.** *Every  $S$ -unimodal infinitely renormalizable map of bounded type has as its maximal automorphic factor an ergodic conservative nonsingular adic odometer action with respect to the factor measure induced by  $\lambda$ .*

The idea of the proof is to show that for  $\lambda$ -a.e.  $x$ ,  $\frac{n_k(x)}{p_k}$  is bounded for sufficiently many  $k$ 's in a sense made precise in Proposition 4.6 below. From this it will follow that for  $\lambda \times \lambda$ -a.e. pair  $(x, y)$ ,  $\frac{|n_k(x) - n_k(y)|}{p_k}$  is bounded sufficiently often. Passing to fibers  $I_\omega = \pi^{-1}(\omega)$ , we can show that for  $\mu$ -a.e.  $\omega \in \Omega$  and  $\lambda_\omega \times \lambda_\omega$ -a.e.  $(x, y) \in I_\omega \times I_\omega$ ,  $\frac{|n_k(x) - n_k(y)|}{p_k}$  is bounded infinitely often. (Here  $\lambda_\omega$  denotes the fiber measure that  $\lambda$  induces on  $I_\omega$ .) We apply this to density points of certain tail sets and complete the proof by a distortion argument which shows that the tail sets must intersect eventually under forward iteration.

**Proposition 4.6.** *Under the assumptions of Theorem 4.4, there exists  $N > 0$  and a sequence  $\{k_j\}$  ( $k_j = j^2$  will suffice) such that for  $\lambda$ -a.e.  $x \in B(\Omega)$ ,*

$$\liminf_{i \rightarrow \infty} \frac{1}{i} \#\{j \leq i; n_{k_j}(x) \leq Np_{k_j-1}\} \geq \frac{2}{3}.$$

*Proof.* Recall the constants  $\rho_+ < 1$  and  $K \geq 1$  from Proposition 4.5. Choose  $N_0$  so large that

$$\frac{K}{1 - \rho_+} \rho_+^{N+1} \leq K \left( \frac{\rho_+^{N+1}}{1 - \rho_+} + \frac{\rho_+^{2N/3+1}}{1 - \rho_+^{1/3}} + \frac{\rho_+^{N-N_0/2}}{1 - \rho_+} \right) \leq \rho_+^{N/3} \quad (2)$$

for all  $N \geq N_0$ . We show that for any  $k$  and  $N \geq N_0$ ,

$$P(n_k \geq Np_{k-1}) = \lambda(\{x; n_k(x) \geq Np_{k-1}\}) \leq \rho_+^{N/3}. \quad (3)$$

We will use induction on  $k$ . For  $k = 1$ , Proposition 4.5 and (2) immediately give

$$P(n_1 \geq Np_0) = P(m_1 \geq N) \leq \sum_{i=N}^{\infty} K\rho_+^{i+1} \leq \frac{K}{1 - \rho_+} \rho_+^{N+1} \leq \rho_+^{N/3}$$

for all  $N \geq N_0$ . For the induction step,

$$\begin{aligned} P(n_k \geq Np_{k-1}) &\leq P(m_k \geq N) + \sum_{i=0}^{N-1} P(m_k = i \text{ and } n_{k-1} \geq (N-i)p_{k-1}) \\ &\leq \frac{K}{1-\rho_+} \rho_+^{N+1} + \sum_{i=0}^{N-N_0/2} K \rho_+^{i+1} P(n_{k-1} \geq 2(N-i)p_{k-2}) \\ &\quad + \sum_{i=N-N_0/2}^{N-1} K \rho_+^{i+1}. \end{aligned}$$

Here we used Proposition 4.5 and the fact that  $p_{k-1} \geq 2p_{k-2}$ . By induction,  $P(n_{k-1} \geq 2(N-i)p_{k-2}) \leq \rho_+^{2(N-i)/3}$  for  $i \leq N - N_0/2$ . Together with (2), this gives

$$\begin{aligned} P(n_k \geq Np_{k-1}) &\leq \frac{K}{1-\rho_+} \rho_+^{N+1} + K \sum_{i=0}^{N-N_0/2} \rho_+^{i+1} \rho_+^{2(N-i)/3} + K \sum_{i=N-N_0/2}^{N-1} \rho_+^{i+1} \\ &\leq K \left( \frac{\rho_+^{N+1}}{1-\rho_+} + \frac{\rho_+^{2N/3+1}}{1-\rho_+^{1/3}} + \frac{\rho_+^{N-N_0/2}}{1-\rho_+} \right) \leq \rho_+^{N/3}. \end{aligned}$$

This proves formula (3).

Now take  $k_i = i^2$ . Since  $p_n \geq 2p_{n-1}$  for all  $n \geq 2$ , we have  $p_{k_{i+1}-1} \geq 2^{2i} p_{k_i-1}$ . By (3) we find for  $i$  sufficiently large

$$P(n_{k_i} \geq \frac{1}{i^2} p_{k_{i+1}-1}) = P(n_{k_i} \geq \frac{1}{i^2} \frac{p_{k_{i+1}-1}}{p_{k_i-1}} p_{k_i-1}) \leq \rho_+^{2^{2i}/3i^2},$$

which is summable over  $i$ . The Borel-Cantelli Lemma gives that the set

$$X := \{x; \exists j \forall i \geq j \ n_{k_i}(x) \leq \frac{1}{i^2} p_{k_{i+1}-1}\} \quad (4)$$

has full measure in the basin of  $\Omega$ . Write  $W_i(x) := \sum_{j=k_{i-1}+1}^{k_i} m_j(x) p_{j-1}$ . The random variables  $W_i$  are not independent. Nevertheless, by the arguments that proved (3) we can show that for any sequence  $v_1, \dots, v_{i-1} \in \mathbb{N}$  and any  $N \geq N_0$ ,

$$\begin{aligned} P(W_i \geq Np_{k_{i-1}} | W_j = v_j \text{ for } j < i) = \\ \lambda(\{x; W_i(x) \geq Np_{k_{i-1}}, W_j(x) = v_j \text{ for } j < i\}) \leq \rho_+^{N/3}. \end{aligned}$$

Next take  $N_1 \geq N_0$  so large that  $\rho_+^{N_1/3} \leq 3^{-3}$ . By the binomial formula and Stirling's formula,

$$P\left(\frac{1}{i} \#\{j \leq i; W_j \geq N_1 p_{k_j-1}\} \geq \frac{1}{3}\right) \leq \sum_{j=\lfloor i/3 \rfloor}^i \binom{i}{j/3} [\rho_+^{N_1/3}]^{j/3} \leq i \binom{i}{i/3} 3^{-i} \leq i 2^{-2i/3},$$

which is summable in  $i$ . Therefore the Borel-Cantelli Lemma gives

$$\limsup_{i \rightarrow \infty} \frac{1}{i} \#\{j \leq i; W_j(x) \geq N_1 p_{k_{j-1}}\} \leq \frac{1}{3} \text{ } \lambda\text{-a.e.}$$

Combining this with (4), and noting that  $\sum_i 1/i^2 < 2$ , we obtain

$$\limsup_{i \rightarrow \infty} \frac{1}{i} \#\{j \leq i; n_{k_j} \geq (N_1 + 2)p_{k_{j-1}}\} \leq \frac{1}{3} \text{ for } \lambda\text{-a.e. } x \in X.$$

This proves the proposition using  $N = N_1 + 2$ .  $\square$

Let  $N$  be the integer chosen in Proposition 4.6, and recall that  $\lambda_\omega$  is the fiber measure that Lebesgue measure induces on the fiber  $I_\omega = \pi^{-1}(\omega)$ .

**Corollary 4.1.** *For  $\mu$ -a.e.  $\omega \in \Omega$  and  $\lambda_\omega \times \lambda_\omega$ -a.e.  $(x, y) \in I_\omega \times I_\omega$ , there are infinitely many values of  $i$  such that  $|n_{k_i}(x) - n_{k_i}(y)| \leq N p_{k_{i-1}}$ .*

*Proof.* By a standard argument on fiber measures, Proposition 4.6 implies that for  $\mu$ -a.e.  $\omega \in \Omega$  and  $\lambda_\omega$ -a.e.  $x \in I_\omega$ ,

$$\limsup_{i \rightarrow \infty} \frac{1}{i} \#\{j \leq i; n_{k_j} \geq N p_{k_{j-1}}\} \leq \frac{1}{3}.$$

Hence the lower density of the set of integers  $i$  such that  $n_{k_i} \leq N p_{k_{i-1}}$  is at least  $\frac{2}{3}$  within each fiber. The corollary follows immediately since  $n_{k_i} \geq 0$ .  $\square$

We are now ready to prove Theorem 4.4.

*Proof.* Let  $\Omega$  be the attractor of  $f$  and let  $\pi : I \rightarrow \Omega$  denote the factor map. We know that  $(\Omega, \mathcal{C}, \mu, g)$  with  $\mathcal{C} = \pi(\mathcal{B})$  and  $\mu = \lambda \circ \pi^{-1}$  is a measurable automorphic factor of  $(I, \mathcal{B}, \lambda, f)$  and that it is isomorphic to some odometer. Assume by contradiction that it is not the maximal automorphic factor. Then there exist  $B, B' \in \mathcal{B}_+$  such that  $\pi(B) = \pi(B') \in \mathcal{C}_+$ , but  $f^n(B) \cap f^n(B') = \emptyset$  for all  $n \geq 0$ . Take  $x$  and  $x'$  Lebesgue density points of  $B$  respectively  $B'$ . By Corollary 4.1 we can assume that  $\pi(x) = \pi(x')$ , and that there is an integer  $N$  such that  $|n_k(x) - n_k(x')| \leq N p_{k-1} \leq N p_k$  infinitely often. Let  $I_k \ni x$  and  $I'_k \ni x'$  be the maximal intervals such that  $f^{n_k(x)}(I_k) = f^{n_k(x')}(I'_k) = J_k$ .

Given  $r \geq 0$ , write  $J_{k,r} = \{y \in J_k; m_{k+1}(y) = r\}$ , so  $J_{k,0} = J_{k+1}$ . By Proposition 4.5,  $\lambda(J_{k,r}) \geq \rho_-^{r+1} \lambda(J_k)$ . Write  $\delta := \rho_-^{N+1}$ . Because  $x$  and  $x'$  are density points, we can take  $n_k(x) \leq n_k(x') = n_k(x) + r p_k$  (for some  $r \leq N$ ) so large that

$$\frac{\lambda(I_k \setminus B)}{\lambda(I_k)}, \frac{\lambda(I'_k \setminus B')}{\lambda(I'_k)} \leq \frac{\delta}{3K^2}.$$

Here  $K \geq 1$  is the distortion constant from Proposition 4.5. It follows that  $\lambda(J_{k,0} \setminus f^{n_k(x')}(B')) \leq \frac{1}{3K}\lambda(J_{k,0})$  and  $\lambda(J_{k,r} \setminus f^{n_k(x)}(B)) \leq \frac{1}{3K}\lambda(J_{k,r})$ . Applying another  $rp_k$  iterates to  $f^{n_k}(B) \cap J_{k,r}$  we find (using the same distortion bound  $K$ ) that  $\lambda(J_{k,0} \setminus f^{n_k(x)+rp_k}(B)) \leq \frac{1}{3}\lambda(J_{k,0})$ . Therefore  $f^{n_k(x')}(B') \cap f^{n_k(x)}(B) \neq \emptyset$ . This contradicts the choice of  $B$  and  $B'$ .  $\square$

## 5 The Fibonacci Unimodal Map

The aim of this section is to prove that a Fibonacci map with a Cantor attractor has a circle rotation as maximal automorphic factor. For this we have to recall some facts from [5, 6].

### 5.1 Factors of $(\omega(c), f)$

Fix a unimodal map  $f$ . Let  $D_n$  be the image of the central branch of  $f^n$  (the largest monotone subinterval containing  $c$ ). If  $c \in D_n$ , we say that  $n$  is a *cutting time*. The cutting times are denoted as

$$1 = S_0 < S_1 < S_2 < \dots$$

They are very important as they determine the combinatorial structure of the unimodal map completely. Obviously  $c_n$  is one endpoint of  $D_n$ . It is not hard to show that  $c_{n-S_k}$ , where  $S_k$  is the maximal cutting time less than  $n$ , is the other.

It can be shown that the difference of two subsequent cutting times is again a cutting time. Hence

$$S_k - S_{k-1} = S_{Q(k)},$$

for some integer function  $Q$ , which is called the *kneading map*, [10]. For more details see [3]. We assume for the rest of the paper that

$$Q(k) \rightarrow \infty \text{ as } k \rightarrow \infty. \tag{5}$$

If the cutting times are the Fibonacci numbers  $1, 2, 3, 5, 8, \dots$ , i.e.  $Q(k) = \max\{0, k - 2\}$ , then  $f$  is called a *Fibonacci map*.

**Lemma 5.1.** *If  $Q(k) \rightarrow \infty$ , then  $\lambda(D_n) \rightarrow 0$  and  $\omega(c)$  is a minimal Cantor set.*

*Proof.* See [3].  $\square$

For the Fibonacci map,  $(\omega(c), f)$  is isomorphic to  $(\mathbb{S}^1, R_\gamma)$ , where  $R_\gamma$  is the circle rotation over  $\gamma = \frac{\sqrt{5}-1}{2}$ . This was shown in [13]. In [6], the result was generalized

to many other unimodal maps and group rotations. We will discuss some tools from [6]. First there is the  $S$ -adic transformation (where  $S = \{S_k\}$  refers to the sequence of cutting times). Let

$$E := \{e \in \{0, 1\}^{\mathbb{N}}; e_i = 1 \Rightarrow e_j = 0 \text{ for } Q(i+1) < j < i\},$$

endowed with product topology. On  $E$  we define  $T$  to be the addition of 1 by means of “add and carry”; ones at entries  $i$  and  $Q(i+1)$  carry to a one at entry  $i+1$ . If  $S_k = 2^k$  (i.e.  $f$  is the Feigenbaum map), we recover the usual dyadic odometer. The set

$$E_0 := \{e \in E; \#\{i; e_i = 1\} < \infty\}$$

is the *greedy* representation of  $\mathbb{N} \cup \{0\}$ , see e.g. [9]. Indeed, if  $n \geq 0$ , there is a canonical way of assigning a sequence  $\langle n \rangle \in E_0$  such that  $n = \sum_i \langle n \rangle_i S_i$ . Take  $i := \max\{k; S_k \leq n\}$  and set  $\langle n \rangle_i = 1$ . Repeat this process with  $n - S_i$ , etc. Then the restriction in the definition of  $E$  will automatically be satisfied.

**Lemma 5.2.** *If  $Q(k) \rightarrow \infty$ , then  $T : E \rightarrow E$  is continuous,  $T(\langle n \rangle) = \langle n+1 \rangle$  and  $T$  is invertible, with a possible exception at  $\langle 0 \rangle$ .*

Let  $\pi_1 : E \rightarrow \omega(c)$  be a projection defined on  $E_0$  by  $\pi_1(\langle n \rangle) = f^n(c)$ , and extended to  $E$  by uniform continuity. Equivalently we can define:

$$\pi_1(\langle n \rangle) = c_n \text{ and for } e \notin E_0, \pi_1(e) = \bigcap_k D_{n_k}, \quad (6)$$

where  $n_k = \sum_{i \leq k} e_i S_i$ . We have  $f \circ \pi_1 = \pi_1 \circ T$ . Note that  $\pi_1$  need not be invertible.

For  $x \in \mathbb{R}$ , choose the fractional part  $\text{frac}(x) \in [-\frac{1}{2}, \frac{1}{2})$  such that  $x - \text{frac}(x)$  is an integer. If there exists  $\alpha \in \mathbb{R}$  such that

$$\sum_i |\text{frac}(\alpha S_i)| < \infty, \quad (7)$$

then we can define a second projection  $\pi_0 : E \rightarrow \mathbb{S}^1$  by

$$\pi_0(e) := \sum_i e_i \text{frac}(\alpha S_i) \pmod{1}.$$

One can show that  $\pi_0 \circ T(e) = \pi_0(e) + \alpha$  for all  $e \in E$ .

In particular, the Fibonacci numbers can be written as  $S_k = \frac{5+3\sqrt{5}}{10}(1+\gamma)^k + \frac{5-3\sqrt{5}}{10}(-\gamma)^k$ , where  $\gamma = \frac{\sqrt{5}-1}{2}$ . Hence there exists  $L$  such that

$$|\text{frac}(\gamma S_k)| = |\text{frac}((1+\gamma)S_k)| = |\text{frac}(S_{k+1} + (1+2\gamma)\frac{5-3\sqrt{5}}{10}(-\gamma)^k)| \leq L\gamma^k \quad (8)$$

for all  $k$ . Therefore (7) is satisfied, and the projection  $\pi_0$  is well-defined.

**Proposition 5.1.** *If  $Q(k) \rightarrow \infty$  and  $\sum_k |\text{frac}(\alpha S_k)| < \infty$ , then  $\pi := \pi_0 \circ \pi_1^{-1}$  is a well-defined continuous mapping and the diagram*

$$\begin{array}{ccc} & (E, T) & \\ \pi_1 \swarrow & & \searrow \pi_0 \\ (\omega(c), f) & \xrightarrow{\pi} & (S^1, R_\alpha) \end{array}$$

*commutes.*

*Proof.* The case where  $S_k$  are the Fibonacci numbers was shown already in [13]. Moreover  $\pi$  is one-to-one, except on the backward orbit of  $c$ , where it is two-to-one. The general case was presented in [6].  $\square$

The projection  $\pi$  can be shown to be one-to-one on a set of full Lebesgue measure on  $\mathbb{S}^1$  for many other unimodal maps as well.

## 5.2 Fibonacci Maps with Attractors

The main result of [5] is the following:

**Theorem 5.1.** *If  $f$  is a  $C^2$  unimodal Fibonacci map with a sufficiently degenerate critical point, then  $\omega(c)$  is an absorbing Cantor set.*

The idea of the proof is as follows: Let  $u_1 = \bar{p}$ , where  $p$  is the orientation reversing fixed point, and for  $k \geq 2$ ,

$$u_k := \begin{cases} f^{-S_{k-1}}(u_{k-1}) \cap (u_{k-1}, c) & \text{if } c_{S_{k-1}} < c, \\ f^{-S_{k-1}}(\bar{u}_{k-1}) \cap (u_{k-1}, c) & \text{if } c_{S_{k-1}} > c. \end{cases} \quad (9)$$

It is shown in [5] that this is a valid definition. In fact, the points  $u_k$  are *closest-to- $c$  prefixed* in the sense that if  $(u_{k-1}, \bar{u}_{k-1}) \cap f^{-n}(p) \neq \emptyset$ , then  $n := \min\{i; f^i(u_k) \in \{p, \bar{p}\}\}$ .  $U_1 = I \setminus [u_1, \bar{u}_1]$  and  $U_k = (u_{k-1}, u_k) \cup (\bar{u}_k, \bar{u}_{k-1})$  for  $k \geq 2$ . Define an *induced map*  $F$  by

$$F|_{U_k} := f^{S_{k-1}} \text{ for all } k \geq 1.$$

By construction, for  $k \geq 2$ ,  $F(U_k) = (u_{k-3}, \bar{u}_{k-1})$  if  $c_{S_{k-1}} > c$  and  $F(U_k) = (u_{k-1}, \bar{u}_{k-3})$  if  $c_{S_{k-1}} < c$ . (Here  $u_0 = u_{-1} = p$ .) Hence  $F$  preserves the partition of  $I$  into sets  $U_k$ . Let

$$\chi_n(x) := k \text{ if } F^n(x) \in U_k.$$

As was shown in [5], there exists a constant  $K$  such that the distortion  $\text{dist}(F^n, J) \leq K$  for any  $n \geq 0$  and interval  $J$  on which  $F^n$  is continuous. The behavior of points under iteration of  $F$  is interpreted as a random walk. It is shown that for maps of sufficiently large critical order, the expectation (with respect to Lebesgue measure)

$$\mathbb{E}(\chi_n - k | \chi_{n-1} = k) \geq \eta > 0, \quad (10)$$

where  $\eta$  is independent of  $k$ , of  $n$  and of the precise path used to get to state  $U_k$ . A similar estimate can be made for the variances:

$$\text{Var}(\chi_n - k | \chi_{n-1} = k) \leq V < \infty. \quad (11)$$

These estimates imply that  $\chi_n(x) \rightarrow \infty$   $\lambda$ -a.e., and this implies that  $f^n(x) \rightarrow \omega(c)$   $\lambda$ -a.e.

### 5.3 The Maximal Automorphic Factor of a Fibonacci Map

Let  $f$  be a Fibonacci map with an absorbing Cantor set. To be precise, assume that (10) and (11) hold. For  $x \in I$  define

$$\beta_k(x) := \max\{n; \chi_n(x) \leq k\},$$

and let  $b_k(x)$  be such that  $F^{\beta_k(x)}\{x\} = f^{b_k(x)}\{x\}$ . Because  $\chi_m \rightarrow \infty$   $\lambda$ -a.e., these sequences are defined  $\lambda$ -a.e. Recall that  $\langle n \rangle$  denotes the  $S$ -adic representation of  $n$ . Define

$$\tilde{\pi}(x) := \lim_{k \rightarrow \infty} \pi_0(\langle b_k(x) \rangle),$$

whenever it exists.

**Lemma 5.3.** *If  $\tilde{\pi}(x)$  exists, then for every  $N \geq 0$ ,  $\tilde{\pi}(f^N(x))$  exists and  $\tilde{\pi}(f^N(x)) = \tilde{\pi}(x) - N\gamma$ .*

*Proof.* We will show that for every fixed  $N \geq 0$ , there exists  $i, j \geq 0$  such that

$$F^i(x) = F^j(f^N(x)). \quad (12)$$

From this it follows that  $b_k(f^N(x)) = b_k(x) - N$  for  $k$  sufficiently large, which proves the lemma.

Let  $\{m_i\}$  and  $\{n_j\}$  be such that  $F^{m_i}\{x\} = f^{m_i}$  and  $F^{n_j}\{f^N(x)\} = f^{n_j}$ . Find  $i, j$  maximal such that  $m_i \leq N$  and  $n_j + N \leq m_{i+1}$ . If in one of these cases equality holds, (12) is true. Let  $y = F^i(x)$  and  $y' = F^j(f^N(x))$ . Then  $y' = f^{N'}(y)$  for some  $N' > 0$ . Take also  $s$  and  $s'$  such that  $y \in U_s$  and  $y' \in U_{s'}$ . In this notation,  $N' < S_{s-1} < S_{s'-1} + N'$ .

Recall that the points  $\{u_k\}$  are the closest-to- $c$  prefixed points, none omitted. Therefore

$$f^n(u_k) \notin [u_k, \bar{u}_k] \text{ for all } n > 0 \text{ and } f^n(u_k) \notin [u_{k-1}, \bar{u}_{k-1}] \text{ for all } 0 < n < S_{k-1}. \quad (13)$$

Indeed, if  $f^n(u_k) \in [u_k, \bar{u}_k]$ , then  $u_k$  is not a closest-to- $c$  prefixed point. If  $f^n(u_k) \in (u_{k-1}, u_k)$  or  $(\bar{u}_k, \bar{u}_{k-1})$ , then  $u_k$  is not the first closest-to- $c$  prefixed point after  $u_{k-1}$ . Finally, by equation (9),  $f^{S_{k-1}}(u_k) \in \{u_{k-1}, \bar{u}_{k-1}\}$ . Therefore  $f^n(u_k) \notin \{u_{k-1}, \bar{u}_{k-1}\}$  for  $n < S_{k-1}$ .

Since  $f^{N'}(y) = y'$ ,  $f^{N'}(U_s)$  intersects  $U_{s'}$ . We distinguish three cases:

- $f^{N'}(U_s) \subset U_{s'}$ . Then  $c \in f^{S_{s-1}}(U_s) \subset f^{S_{s-1}-N'}(U_{s'})$ . This contradicts that  $S_{s'-1} + N' > S_{s-1}$ .
- $u_{s'} \in f^{N'}(U_s)$ . By equation (13),  $s' \leq s - 2$ . On the other hand,  $f^{S_{s-1}-N'}(u_{s'}) \in F(U_s) \subset (u_{s-3}, \bar{u}_{s-3})$ . Therefore  $s - 3 < s' - 1$ . This contradicts  $s' \leq s - 2$ .
- $u_{s'-1} \in f^{N'}(U_s)$ . By equation (13),  $s' - 1 \leq s - 2$ . On the other hand,  $f^{S_{s-1}-N'}(u_{s'-1}) \in F(U_s) \subset (u_{s-3}, \bar{u}_{s-3})$ . Therefore  $s - 3 < s' - 2$ , contradicting  $s' - 1 \leq s - 2$ .

These contradictions establish the proof.  $\square$

The main result of this section is that the map  $\tilde{\pi}$  is defined  $\lambda$ -a.e., and that the circle rotation with appropriate measure algebra is the maximal automorphic factor.

**Theorem 5.2.** *Let  $(I, \mathcal{B}, \lambda, f)$  be a Fibonacci map satisfying (10) and (11) (and therefore has an absorbing Cantor set). Then  $\tilde{\pi} : I \rightarrow \mathbb{S}^1$  is defined  $\lambda$ -a.e. If  $\mu := \lambda \circ \tilde{\pi}^{-1}$  and  $\mathcal{C} := \tilde{\pi}(\mathcal{B})$ , then the rotation  $(\mathbb{S}^1, \mathcal{C}, \mu, R_\gamma^{-1})$  is the maximal automorphic factor.*

*Proof.* Let  $V_k = \{x \in U_k; \chi_m(x) > k \text{ for all } m \geq 1\}$ . Equations (10) and (11) show that a definite proportion of the set  $U_k$  never returns to  $\cup_{i \leq k} U_i$ . Hence there exists  $\eta_1 > 0$  such that  $\lambda(V_k) \geq \eta_1 \lambda(U_k)$  for all  $k \geq 1$ . Moreover, the branches on  $F^m$  have a uniform distortion bound  $K$ . Therefore, taking  $\eta_0 = \eta_1/K$ , we find that the probability

$$P(F^m(x) \in V_k | \chi_m(x) = k) \geq \eta_0. \quad (14)$$

These estimates are independent of  $m$  and of the precise path used to get to state  $U_k$ . If  $\beta_{k+1}(x) - \beta_k(x) > k$ , then  $F^i(x) \in \cup_{i > k} U_i \setminus V_i$  for  $b_k(x) < i < \beta_k(x) + k$ . Therefore

$$P(\beta_{k+1}(x) - \beta_k(x) > k) \leq (1 - \eta_0)^k.$$



The Borel-Cantelli Lemma gives that  $P\{\beta_{k+1}(x) - \beta_k(x) > k \text{ infinitely often}\} = 0$ . Write  $\tilde{\pi}_k(x) = \pi_0(\langle b_k(x) \rangle)$ . For  $\lambda$ -a.e.  $x \in I$  there exists  $k_0$  such that

$$b_{k+1}(x) - b_k(x) = \sum_{i \geq 0} a_{k,i}(x) S_{k+i}, \text{ where } \sum_i a_{k,i}(x) \leq k,$$

for all  $k \geq k_0$ . Therefore (8) gives

$$|\tilde{\pi}_{k+1}(x) - \tilde{\pi}_k(x)| \leq \sum_{i \geq 0} a_{k,i} \text{frac}(\gamma S_{k+i}) \leq k |\text{frac}(\gamma S_k)| = Lk\gamma^k.$$

Hence  $\tilde{\pi}_k(x)$  is a Cauchy sequence, converging to  $\tilde{\pi}(x)$ . This shows that  $\tilde{\pi}$  is defined  $\lambda$ -a.e. and that  $\mathcal{C} = \tilde{\pi}(\mathcal{B})$  and  $\mu = \lambda \circ \tilde{\pi}^{-1}$  are well-defined up to measure 0. The relation  $\tilde{\pi} \circ f(x) = R_\gamma^{-1} \circ \tilde{\pi}(x)$  was already established in Lemma 5.3. It follows that  $R_\gamma^{-1}$  is nonsingular with respect to  $\mu$ , and that  $(\mathbb{S}^1, \mathcal{C}, \mu, R_\gamma^{-1})$  is a measurable automorphic factor.

We need the following lemma to show the automorphic factor is maximal.

**Lemma 5.4.** *There exists a decreasing function  $\xi : \mathbb{R}_+ \rightarrow [0, 1]$  converging to 0 as  $x \rightarrow \infty$  such that*

$$P(b_k \geq NS_k) := \lambda(\{x; b_k(x) \geq NS_k\}) \leq \xi(N)$$

for all  $k$ .

*Proof.* Using the  $\eta_0$  obtained in (14), define for any  $N \in \mathbb{R}^+$ ,

$$\xi(N) := 1 - (1 - (1 - \eta_0)^{M/2})(1 - \frac{1}{\eta_0} (1 - \eta_0)^M),$$

with  $M = \frac{\log N - \log 10}{\log \frac{1}{\gamma}}$ . Clearly  $\xi(N) \rightarrow 0$  as  $N \rightarrow \infty$ . We will show that  $\xi(N)$  gives the desired estimate.

Fix  $k, N \in \mathbb{N}$  and let  $x \in B(\omega(c))$  arbitrary (provided  $b_k(x)$  exists). Let  $M = M(N) = \frac{\log N - \log 10}{\log \frac{1}{\gamma}}$ . First we calculate the probability that

$$\chi_j(x) > k + M \text{ for some } j \leq \beta_k(x). \tag{15}$$

If this occurs, then, since a point can jump back no more than two states under one iteration of  $F$ ,  $j < \beta_k(x) - M/2$ . Since  $F^i(x) \notin V_m$  for  $j < i < \beta_k(x)$  and  $k < m \leq k + M$  (cf. formula (14)), the probability that  $\chi_j > k + M$  is less than  $(1 - \eta_0)^{M/2}$ .

Next we verify, using (14) again, that  $P(\#\{j; \chi_j = m\} \geq t) \leq (1 - \eta_0)^{t-1}$ . Indeed, because points have to avoid the set  $V_m$   $t - 1$  times to make  $t$  returns to  $U_m$  possible, it follows that

$$\begin{aligned} P(\#\{j \leq \beta_k; \chi_j = m\} \geq 2M + k - m \text{ for some } 1 \leq m \leq k + M) \\ \leq \sum_{m=1}^{k+M} (1 - \eta_0)^{2M+k-m-1} \leq \frac{1}{\eta_0} (1 - \eta_0)^M. \end{aligned} \quad (16)$$

If neither (15) nor (16) occurs, i.e. if  $\#\{j \leq \beta_k(x); \chi_j(x) = m\} \leq 2M + k - m$  for all  $1 \leq m \leq k + M$  and  $\chi_j(x) \leq k + M$  for all  $j \leq \beta_k(x)$ , then

$$b_k(x) = \sum_{m=1}^{k+M} \#\{j \leq \beta_k(x); \chi_j(x) = m\} S_{m-1} \leq \sum_{m=1}^{k+M} (2M + k - m) S_{m-1} \leq N S_k.$$

This happens with probability at least  $(1 - (1 - \eta_0)^{M/2})(1 - \frac{1}{\eta_0} (1 - \eta_0)^M) = 1 - \xi(N)$  which tends to 1 uniformly in  $k$  as  $N \rightarrow \infty$ . This proves the lemma.  $\square$

Returning to the proof of Theorem 5.2, we copy the arguments of Theorem 4.4. That is, we find a subsequence  $\{k_i\}$  of  $\mathbb{N}$  which increases fast enough so that  $\sum_i P(b_{k_i} \geq \frac{1}{i^2} S_{k_{i+1}}) \leq \sum_i \xi(\frac{1}{i^2} \frac{S_{k_{i+1}}}{S_{k_i}}) < \infty$ . Then by the Borel Cantelli Lemma,


$$X := \{x; \exists j \forall i \geq j \ b_{k_i} \leq \frac{1}{i^2} S_{k_{i+1}}\}$$

has full measure in the basin of  $\omega(c)$ . Take  $N_1$  so large that  $\xi(N_1) \leq 3^{-3}$ . A proof similar to the one of Lemma 5.4 gives  $P(b_{k_i} - b_{k_{i-1}} \geq N_1 S_{k_i}) \leq \xi(N_1) \leq 3^{-3}$ . Then we can derive (cf. Proposition 4.6) that  $\liminf_i \frac{1}{i} \#\{j \leq i; b_{k_j}(x) \leq N S_{k_j}\} \geq \frac{2}{3}$  for  $N = N_1 + 2$  and  $\lambda$ -a.e.  $x \in B(\omega(c))$ . Given  $s \in \mathbb{S}^1$ , let  $I_s := \tilde{\pi}^{-1}$  be the fiber over  $s$  and let  $\lambda_s$  be the fiber measure that  $\lambda$  induces on  $I_s$ . Then (cf. Corollary 4.1) for  $\mu$ -a.e.  $s$  and  $\lambda_s \times \lambda_s$ -a.e.  $(x, x') \in I_s \times I_s$ , we have established that  $|b_k(x) - b_k(x')| \leq N S_k$  infinitely often.

The last step in the proof is to assume by contradiction that  $(\mathbb{S}^1, \mathcal{C}, \mu, R_\gamma^{-1})$  is not the maximal automorphic factor. Then there exist  $B, B' \in \mathcal{B}_+$  such that  $\pi(B) = \pi(B')$  and  $\lambda(f^n(B) \cap f^n(B')) = 0$  for all  $n \geq 0$ . We choose Lebesgue density points  $x$  and  $x'$  of  $B$  and  $B'$  respectively such that  $\tilde{\pi}(x) = \tilde{\pi}(x')$ . By the above arguments we can assume that  $|b_k(x) - b_k(x')| \leq N S_k$  infinitely often. Take such a  $k$ ; then  $\langle b_k(x) \rangle_i = \langle b_k(x') \rangle_i$  for all  $i > k + P$  for  $P = P(N) = -\frac{2 \log N}{\log \gamma}$ . Abbreviate  $\chi = \chi_{\beta_k(x)}$  and  $\chi' = \chi_{\beta_k(x')}$ , so  $F^{\beta_k(x)}(x) \in U_\chi$  and  $F^{\beta_k(x')}(x') \in U_{\chi'}$ . Because  $\tilde{\pi}(x) = \tilde{\pi}(x')$ ,

$$\langle b_k(x) + S_{\chi-1} \rangle_i = \langle b_k(x') + S_{\chi'-1} \rangle_i$$

for all  $i \leq k$ . By definition of  $\beta_k$ , both  $F^{\beta_k(x)+1}(x)$  and  $F^{\beta_k(x')+1}(x')$  are contained in  $(u_k, \bar{u}_k)$ . Let  $W_k$  be the component of  $U_\chi \cap F^{-1}([u_k, \bar{u}_k])$  that contains  $F^{\beta_k(x)}(x)$ ; similarly, let  $W'_k$  be the component of  $U_{\chi'} \cap F^{-1}([u_k, \bar{u}_k])$  that contains  $F^{\beta_k(x')}(x')$ .



Construction of  $T, T', W_k$  and  $W'_k$

Take

$$d := S_{\chi'_{-1}} + \sum_{i=k+1}^{k+P} \langle b_k(x') + S_{\chi'_{-1}} \rangle_i S_{i-1} \text{ and } d' := S_{\chi_{-1}} + \sum_{i=k+1}^{k+P} \langle b_k(x) + S_{\chi_{-1}} \rangle_i S_{i-1}.$$

Then  $D := b_k(x) + d = b_k(x') + d'$ . Find the unique interval  $T \subset W_k$  whose orbit is given by  $\langle d \rangle$  in the following sense: Let  $i_0, i_1, \dots, i_n$  (with  $1 \leq n \leq P$ ) be the indices for which  $\langle d \rangle_i = 1$ . Then  $T$  is taken such that

$$F^1(T) \subset U_{i_1}, F^2(T) \subset U_{i_2}, \dots, F^n(T) \subset U_{i_n} \text{ and } F^{n+1}(T) = [u_{k+P}, \bar{u}_{k+P}].$$

The interval  $T' \subset W'_k$  is chosen similarly. Then  $F^{n+1}(T) = F^{n'+1}(T')$ . Recall that the distortion of the branches of iterates of  $F$  is uniformly bounded by  $K$ . From this, one can derive that there exists  $\varepsilon = \varepsilon(P, K) > 0$  such that  $\lambda(T) \geq \varepsilon \lambda(W_k)$  and  $\lambda(T') \geq \varepsilon \lambda(W'_k)$ .

We now take intervals  $J \ni x$  and  $J' \ni x'$  such that the maps  $f^{b_k(x)} : J \rightarrow W_k$  and  $f^{b_k(x')} : J' \rightarrow W'_k$  are monotone onto, and such that

$$\frac{\lambda(J \setminus B)}{\lambda(J)}, \frac{\lambda(J' \setminus B')}{\lambda(J')} \leq \frac{\varepsilon}{3K^2}.$$

Since  $x$  and  $x'$  are density points, such intervals can be found for  $k$  sufficiently large. Therefore

$$\frac{\lambda(T \setminus f^{b_k(x)}(J \cap B))}{\lambda(T)}, \frac{\lambda(T' \setminus f^{b_k(x')}(J' \cap B'))}{\lambda(T')} \leq \frac{1}{3K}.$$

Applying  $F^{n+1}$  to  $f^{b_k(x)}(J) \cap T$  and  $F^{n'+1}$  to  $f^{b_k(x')}(J') \cap T'$ , and using the distortion bound  $K$  once more, we obtain

$$\begin{aligned} 0 < \frac{1}{3}\lambda(F^{n+1}(T)) &\leq \lambda(F^{n+1} \circ f^{b_k(x)}(J \cap B) \cap F^{n'+1} \circ f^{b_k(x')}(J' \cap B')) \\ &\leq \lambda(f^D(B) \cap f^D(B')). \end{aligned}$$

This contradicts the choice of  $B$  and  $B'$ .  $\square$

**Remark 1:** We now have  $\tilde{\pi} : I \rightarrow \mathbb{S}^1$  (defined  $\lambda$ -a.e.) such that  $\tilde{\pi} \circ f = R_\gamma^{-1} \circ \tilde{\pi}$  and  $\pi : \omega(c) \rightarrow \mathbb{S}^1$  such that  $\pi \circ f = R_\gamma \circ \pi$ . One can show that  $\tilde{\pi}$  is defined on  $\omega(c)$  and that  $\tilde{\pi}(y) = -\pi(y)$  for all  $y \in \omega(c)$ . This relation plays no role in our results, so we omit the proof.

**Remark 2:** In view of the previous remark,  $(\tilde{\pi}|_{\omega(c)})^{-1} \circ \tilde{\pi}$  gives a factor map from  $I$  directly onto the attractor. One might expect that for  $\lambda$ -a.e.  $x$  there exists  $y \in \omega(c)$  such that  $|f^n(x) - f^n(y)| \rightarrow 0$ . This is not true, in spite of the fact that for the candidate  $y \in \tilde{\pi}^{-1} \circ \tilde{\pi}(x) \cap \omega(c)$  there is a sequence  $\{n_i\}$  such that  $|f^{n_i}(x) - c|, |f^{n_i}(y) - c|$  simultaneously tend to 0. The reason is as follows. Assume that  $f^{n_i}(x) \in U_k$  is so close to  $c_{S_k}$  or  $\bar{c}_{S_k}$  that  $f^{n_i+S_{k-1}}(x) \in U_{k+1}$ . Then  $f^{n_i+S_{k-1}+S_{k-1}}(x) = f^{n_i+S_{k+1}-1}(x)$  is close to  $c_{S_{k+2}-1}$ . Checking the kneading invariant of  $f$  shows that  $c_{S_{k+2}-1}$  is close to  $f^{-1}(c) \cap [c, c_1]$  if  $k$  is even and close to  $f^{-1}(c) \cap [c_2, c]$  if  $k$  is odd. If at the same time  $f^{n_i}(y) \in \cup_{j \geq k+2} U_j$ , then  $f^{n_i+S_{k+1}-1}(y)$  is close to  $c_{S_{k+1}-1}$ , which is close to  $f^{-1}(c) \cap [c, c_1]$  if  $k$  is odd and close to  $f^{-1}(c) \cap [c_2, c]$  if  $k$  is even. Therefore  $\limsup |f^n(x) - f^n(y)| \geq \text{diam}(f^{-1}(c)) > 0$ .

## 6 A Dissipative Exact Unimodal Map

The results of Section 5 can be generalized to other unimodal maps with absorbing Cantor sets. In [4] combinatorial conditions are given under which a unimodal map  $f$  with sufficiently large critical order has an absorbing Cantor set. The main condition on the kneading map  $Q$  is that  $k - Q(k)$  is bounded. This applies to many examples from [6], for which  $(\omega(c), \rho, f)$  ( $\rho$  being the unique invariant probability measure) is shown to be isomorphic to some circle and torus rotation with Haar measure. In particular, for the maps with kneading maps  $Q(k) = \max(k - d, 0)$  for  $d = 2, 3, 4$ ,  $(\omega(c), \rho, f)$  is shown to be isomorphic to a rotation on a  $d - 1$ -dimensional torus.

However, for  $d \geq 5$ ,  $(\omega(c), \rho, f)$  has no nontrivial group as factor [6]. In this section we show that the unimodal map  $f$  with kneading map  $Q(k) = k - 5$  for  $k \geq 5$  is exact on  $I$ , even in the presence of an absorbing Cantor set. The proof that  $f$ , for a sufficiently large critical order, has a Cantor attractor is similar to the proof for the Fibonacci map. The inequalities (10) and (11) can be proven. The difference with

the Fibonacci map is that the leading root  $\alpha$  of the equation  $x^5 - x^4 - 1 = 0$  is not a Pisot-Vijayaraghavan number. More precisely, this equation has two roots on, two roots inside, and one root outside the unit circle. Therefore  $|\text{frac}(\alpha S_i)|$  is not summable and the map  $\pi_0$  cannot be defined. Related to this is the following lemma.

**Lemma 6.1.** *Suppose  $S_k$  are the cutting times corresponding to the kneading map  $Q(k) = \max(k - 5, 0)$ , i.e.  $S_k = k + 1$  for  $0 \leq k \leq 4$  and  $S_k = S_{k-1} + S_{k-5}$  otherwise. Then for  $k \geq 3$*

$$S_k = \begin{cases} S_{k-2} + S_{k-3} & \text{if } k \equiv 1 \text{ or } 4 \pmod{6}, \\ S_{k-2} + S_{k-3} + 1 & \text{if } k \equiv 2 \text{ or } 3 \pmod{6}, \\ S_{k-2} + S_{k-3} - 1 & \text{if } k \equiv 0 \text{ or } 5 \pmod{6}. \end{cases}$$

*Proof.* Straightforward by induction. □

**Theorem 6.1.** *Let  $f$  be the unimodal map with kneading map  $Q(k) = \max(0, k - 5)$ . Suppose that the critical order  $\ell$  is so large that  $f$  has a Cantor attractor, and a fortiori, (10) and (11) hold. Then  $f$  is Lebesgue exact.*

*Proof.* Take  $A$  arbitrary such that  $\lambda(A) > 0$ . Without loss of generality, we can assume that  $A \subset (u_1, \bar{u}_1)$ . We will show that Proposition 2.1 applies. Because  $f$  has a Cantor attractor, and a fortiori  $\chi_n(x) \rightarrow \infty$   $\lambda$ -a.e., we can assume that  $\chi_n(x) \rightarrow \infty$  for all  $x \in A$ . Let  $x \in A$  be a density point of  $A$ , such that  $f(x)$  is a density point of  $f(A)$ .

The proof of the existence of Cantor attractors [4] gives rise to the following distortion estimate: For any  $n$  and any  $J$  on which  $F^n|_J = f^n|_J$  is continuous, we have  $\text{dist}(f^n, J) \leq K$ , where  $K$  depends only on  $\ell$ . The proofs also yield that there exists  $C > 0$  such that  $\frac{1}{C\ell} \leq \frac{|u_k - u_{k+1}|}{|u_k - c|} \leq \frac{C}{\ell}$  for all  $k$ . Let  $\varepsilon = \frac{1}{6C^2\ell^2K^3} > 0$ . Because  $x$  is a density point, there exists  $J \ni x$  such that

$$\frac{\lambda(A \cap J')}{\lambda(J')} \geq 1 - \varepsilon$$

for any subinterval  $J'$  such that  $x \in J' \subset J$ . Take from now on  $n$  so large that  $J_n \subset J$  whenever  $J_n \ni x$  is the maximal interval on which  $F^n|_{J_n}$  is continuous.

Assume that  $\chi_n(x) = k$  where  $k \equiv 2, 3 \pmod{6}$ . Let  $U \subset U_k$  be the component containing  $F^n(x) = f^n(x)$ . Define  $\tilde{W}_1 \subset U$  to be the maximal interval such that  $F(\tilde{W}_1) \subset U_k$  and  $F^2(\tilde{W}_1) \subset U_{k+1}$ . Similarly  $\tilde{W}_2 \subset U$  will be the maximal interval such that  $F(\tilde{W}_2) \subset U_{k+3}$ . Because  $(u_{k+5}, \bar{u}_{k+5}) \subset F(U_{k+1})$ ,  $F(U_{k+3}) \subset (u_{k-20}, \bar{u}_{k-20})$ , the overlap  $F(U_{k+1}) \cap F(U_{k+3})$  satisfies  $\lambda(F(U_{k+1}) \cap F(U_{k+3})) \geq \frac{1}{2}\lambda(F(U_{k+1}))$  for  $\ell$  sufficiently large. We can find maximal intervals  $W_1 \subset \tilde{W}_1$  and  $W_2 \subset \tilde{W}_2$  such that  $F^3(W_1) = F^2(W_2)$ , and

$$\frac{\lambda(W_1)}{\lambda(U)}, \frac{\lambda(W_2)}{\lambda(U)} \geq \frac{1}{2C^2\ell^2K}.$$

Let  $V_1 = f^{-m}(W_1) \cap J_n$  and  $V_2 = f^{-m}(W_2) \cap J_n$ . Using the distortion argument once more, we derive that

$$\frac{\lambda(V_1)}{\lambda(J_n)}, \frac{\lambda(V_2)}{\lambda(J_n)} \geq \frac{1}{2C^2\ell^2K^2}.$$

By the choice of  $\varepsilon$ ,

$$\frac{\lambda(A \cap V_1)}{\lambda(V_1)}, \frac{\lambda(A \cap V_2)}{\lambda(V_2)} \geq 1 - \frac{1}{3K}.$$

Remember that  $k \equiv 2, 3 \pmod{6}$ , so by Lemma 6.1,  $S_k + S_{k+1} = S_{k+3} - 1$ . Let  $N := m + S_k + S_{k+1} = m - 1 + S_k + S_{k+3}$ . Then

$$\lambda(f^N(A) \cap f^{N+1}(A)) \geq \frac{1}{3}\lambda(F(U_{k+1}) \cap F(U_{k+3})) > 0.$$

It follows that  $\lambda(f^{-N} \circ f^{N+1}(A) \cap A) > 0$ . This is the assumption of Proposition 2.1.

The cases  $k \equiv 0, 1, 4, 5 \pmod{6}$  can be dealt with in a similar way.  $\square$

## References

- [1] J. Aaronson, *An introduction to infinite ergodic theory*, Math. Surveys & Monographs, **50** AMS, Providence RI, (1997).
- [2] A.M. Blokh, M. Lyubich, *Measurable dynamics of S-unimodal maps*, Ann. Sc. Éc. Norm. Sup. **24** (1991), 545-573.
- [3] H. Bruin, *Combinatorics of the kneading map*, Int. Jour. of Bifur. and Chaos **5** (1995), 1339-1349.
- [4] H. Bruin, *Topological conditions for the existence of absorbing Cantor sets*, Trans. AMS **350** (1998), 2229-2263.
- [5] H. Bruin, G. Keller, T. Nowicki, S. van Strien, *Wild Cantor Attractors exist*, Ann. Math. **143** (1996), 97-130.
- [6] H. Bruin, G. Keller, M. St.Pierre, *Adding machines for wild attractors*, Ergod. Th. & Dyn. Sys. **17** (1997), 1267-1287.
- [7] P. Couillet, C. Tresser, *Itération d'endomorphismes et groupe de renormalisation*, J. de Physique **C5** (1978), 25.
- [8] M. J. Feigenbaum, *Qualitative universality for a class of nonlinear transformations*, J. Stat. Phys. **19** (1978), 19-52.

- [9] A. S. Fraenkel, *Systems of numeration*, Amer. Math. Monthly **92** (1985), 105-114.
- [10] F. Hofbauer, *The topological entropy of the transformation  $x \mapsto ax(1-x)$* , Monatsh. Math. **90** (1980), 117-141.
- [11] G. Keller, *Exponents, attractors and Hopf decompositions for interval maps*, Ergod. Th. & Dyn. Sys. **10** (1990), 717-744.
- [12] F. Ledrappier, *Some properties of absolutely continuous invariant measures of an interval*, Ergod. Th. & Dyn. Sys. **1** (1981), 77-93.
- [13] M. Lyubich, J. Milnor, *The Fibonacci unimodal map*, Journ. AMS **6** (1993), 425-457.
- [14] M. Martens, *Interval Dynamics*, Ph.D. Thesis, Delft (1990).
- [15] M. Martens, *Distortion results and invariant Cantor sets of unimodal maps* Ergod. Th. & Dyn. Sys. **14** (1994), 331-349.
- [16] M. Martens, *The existence of  $\sigma$ -finite measures, applications to real one-dimensional dynamics*, Stony Brook Preprint **1992-1**
- [17] W. de Melo, S. van Strien, *One-Dimensional Dynamics*, Springer, Berlin Heidelberg New York, (1993).
- [18] J. Milnor, *On the concept of attractor*, Commun. Math. Phys. **99** (1985), 177-195.
- [19] M. Misiurewicz, *Attractive Cantor sets of positive measure for  $C^\infty$  maps of an interval*, Ergod. Th. & Dyn. Sys. **2** (1982), 405-415.
- [20] A. Renyi, *Representations for real numbers and their ergodic properties*, Acta Math. Akad. Sci. Hungar. **8** (1957), 477-493.
- [21] V. Rohlin, *Exact endomorphisms of a Lebesgue space*, AMS Transl. Ser. 2 **39** (1963), 1-36.