

## Liouville's Theorem in Conformal Geometry

W. Kühnel  
H.-B. Rademacher

Vienna, Preprint ESI 1862 (2006)

October 25, 2006

Supported by the Austrian Federal Ministry of Education, Science and Culture  
Available via <http://www.esi.ac.at>

# Liouville's theorem in conformal geometry

W. KÜHNEL and H.-B. RADEMACHER

ABSTRACT: Liouville's theorem states that all conformal transformations of  $E^n$  and  $S^n$  ( $n \geq 3$ ) are restrictions of Möbius transformations. As a generalization, we determine all conformal mappings of semi-Riemannian manifolds preserving pointwise the Ricci tensor. It turns out that, up to isometries, they are essentially of the same types as in the classical case but the metric may be different.

MSC 2000: 53A30, 53B30, 53C50

In two dimensions conformal mappings are nothing but holomorphic functions between two open parts of the complex numbers. In higher dimensions, the situation is much more rigid. The classical theorem of Liouville [28] states that a conformal mapping between two open parts of Euclidean 3-space is the composition of a similarity and an inversion. This work was motivated by conformal maps in cartography. Lie [27] generalized this to the case of Euclidean  $n$ -space, for expositions in textbooks see [34, p.173] and [2, Thm.A.3.7], other references are [30], [17], [13] and, under the assumption of minimal possible regularity, [32], [4]. In a slightly modified form it holds also in the case of pseudo-Euclidean  $n$ -space of arbitrary signature, see [15], [35, p.209]. It is also well known that the conformal mappings of (pseudo-)Euclidean space are precisely those preserving the sets of hyperspheres and hyperplanes. This aspect of Liouville's theorem is emphasized in [24], [25], [3].

In this note we point out a more general version of Liouville's theorem in the context of semi-Riemannian geometry. It turns out that an analogous theorem holds for arbitrary conformal mappings preserving pointwise the Ricci tensor. Particular cases are the flat case (pseudo-Euclidean space) and the Ricci-flat case (special Einstein spaces). In any case we have the same type of phenomenon, sketched in [11]: Up to isometries, the mapping is necessarily of one of the following three types:

1. a *similarity* (or dilatation) which fixes one point or limit point,

2. an *inversion* sending one point or limit point to infinity,
3. a special type of a transformation with a parallel isotropic vector field as the gradient of the conformal factor.

In the ordinary flat Minkowski space  $\mathbb{R}_1^4 = \{(u, v, x, y)\}$  with the metric  $g = -2dudv + u^2(dx^2 + dy^2)$  a combination of a similarity with a boost isometry leads to a specific homothety of type 1' which fixes a null geodesic and which was investigated in more generality by Alekseevskii [1]. A simple example for type 3 is the mapping  $F(u, v, x, y) = (-\frac{1}{u}, v + \frac{u}{2}(x^2 + y^2), ux, uy)$ . It is conformal with  $F^*g = \frac{1}{u^2}g$ . The gradient of the function  $u$  is the null vector  $\frac{\partial}{\partial v}$ . In General Relativity type 3 occurs also for *pp*-waves, see [29], [23]. In flat Minkowski space it can also be defined as a combination of a translation with two different inversions, compare [15]. Type 1 and type 2 occur also on cones where the limit points need not be part of the manifold. Normally, the apex of the cone is a singularity. Only in a flat space the apex can be a removable singularity, compare Corollary 2. A special example for type 1 is the flat Lorentzian warped product metric  $ds^2 = -dr^2 + r^2ds_{-1}^2$  where  $ds_{-1}^2$  denotes a 3-dimensional compact Riemannian manifold of constant negative curvature. The mapping  $F(r, x) = (cr, x)$  is such a similarity. By the transformation  $r = e^t$  the metric can also be written in the form  $ds^2 = e^{2t}(-dt^2 + ds_{-1}^2)$ . This is called the *expanding hyperbolic spacetime* in [9]. The types 1 and 3 occur also as members of 1-parameter groups generated by conformal vector fields. However, in this paper we focus on the point transformations (or just one conformal change of the metric) because Liouville's theorem does also.

The analysis and the classification of these cases is based on an ODE reduction of the differential equation  $Ric_{\bar{g}} = Ric_g$  for the two Ricci tensors of two conformally equivalent metrics  $\bar{g}$  and  $g$ . Global results about complete spaces admitting such conformal mappings were given in [10] and in our previous paper [19]. Here the results are semi-local in some sense because only the homotheties can be defined on a complete space.

NOTATIONS: Let  $(M, g)$  be an  $n$ -dimensional semi-Riemannian manifold with a metric tensor  $g$  of arbitrary signature  $(k, n - k)$ . A *conformal diffeomorphism*  $f: (M, g) \rightarrow (\bar{M}, \bar{g})$  is a mapping preserving the angles (or the orthogonality) between any two directions and the type (space-like or time-like) of vectors. An

equivalent formulation is that the induced metric  $f^*\bar{g}$  is everywhere a positive multiple of  $g$ . It is convenient to introduce this positive factor as  $\varphi^{-2}$  with a function  $\varphi: M \rightarrow \mathbb{R}_+$ . So on  $M$  we can consider two conformally equivalent metrics  $g$  and  $\bar{g} = \varphi^{-2}g$ . Such a transformation  $f$  is called an *isometry* if  $\varphi = 1$ , it is called a *homothety* or *similarity* if  $\varphi$  is constant. The classical case of a dilatation in Euclidean space is the mapping  $x \mapsto cx$  with a real constant  $c$ , the standard inversion is the mapping  $x \mapsto x/\|x\|^2$ , other inversions are the mappings  $x \mapsto cx/\|x\|^2$  for a real constant  $c$ .  $Ric$  or  $Ric_g$  denotes the Ricci tensor of the metric  $g$ . By definition  $Ric(X, Y)$  is nothing but the trace of the transformation  $R(-, X)Y$  where  $R$  denotes the curvature tensor of type (1,3). Note that the Ricci tensor is invariant under scaling, i.e.,  $Ric_{cg} = Ric_g$  for any positive constant  $c$ . Let  $\nabla$  denote the Levi-Civita connection induced by  $g$ . For any given smooth function  $\varphi$  on  $(M, g)$  let  $\nabla\varphi$  denote the *gradient* of  $\varphi$ ,  $\nabla^2\varphi$  denotes the *Hessian* (0,2)-tensor,  $\Delta\varphi = trace_g\nabla^2\varphi$  is the *Laplacian* of  $\varphi$ . The expression  $(\nabla^2\varphi)^0 = \nabla^2\varphi - \frac{\Delta\varphi}{n} \cdot g$  denotes the traceless part of the Hessian. A *conformal vector field*  $V$  is generated by a local 1-parameter group of conformal mappings.  $V$  is conformal if and only if  $\mathcal{L}_Vg = 2\psi \cdot g$  for a real function  $\psi$ . By the equation  $\mathcal{L}_{\nabla\varphi}g = 2\nabla^2\varphi$  a gradient field  $\nabla\varphi$  is conformal if and only if  $(\nabla^2\varphi)^0 = 0$ . This equation has been studied in many papers. One crucial case is the equation  $\nabla^2\varphi = \pm\varphi \cdot g$ , compare [36], [18]. For general facts about conformal transformations of semi-Riemannian manifolds we refer to the monographs [8] and [16].

DEFINITION: We call a non-isometric conformal diffeomorphism  $f: (M, g) \rightarrow (\bar{M}, \bar{g})$  a *Liouville mapping* if the equation  $Ric_{f^*\bar{g}} = Ric_g$  holds as a pointwise identity on  $M$ . If  $M = \bar{M}$  then we call a conformal transformation  $g \mapsto \bar{g}$  a *Liouville transformation* of the metric if the equation  $Ric_{\bar{g}} = Ric_g$  holds everywhere. A conformal vector field inducing a 1-parameter group of Liouville mappings is called a *conformal Liouville vector field* (an example of type 3 will given at the end of the paper). A Liouville vector field  $V$  is a particular case of what is called a *Ricci collineation*  $\mathcal{L}_VRic = 0$ , compare [22]. If in a spacetime the stress-energy momentum tensor is preserved by a certain Liouville mapping  $\bar{g}$  then the Einstein field equations are also preserved since  $Ric_{\bar{g}} = Ric_g$  implies  $\bar{S}\bar{g} = Sg$ . This gives a certain conformal interpretation of these equations even if they are not conformally invariant in general. In the vacuum case this reduces to the statement that

any conformal mapping between vacuum spacetimes is a Liouville mapping.

Obviously, the identity map between  $(M, g)$  and  $(M, cg)$  is a Liouville transformation if  $c$  is constant. Moreover, any homothetic mapping is a Liouville mapping. Therefore, throughout this paper we focus on the case of non-homothetic conformal mappings. The homothetic case was studied in [1] and [7]. In [11] a Liouville mapping is called a *quasi-similarity*, a Liouville transformation is called a quasi-homothety. We prefer our terminology because it is the aim of the present paper to show that all these considerations are in fact generalizations of Liouville's theorem in conformal geometry.

**Key example:** If we look at Euclidean space we find that there are translations and inversions simultaneously. Normally we describe inversions by polar coordinates. So it is instructive to look at translations in polar coordinates. In the Euclidean plane with  $x = r \cos \varphi, y = r \sin \varphi$  a fixed vector  $V = x_0 \partial_x + y_0 \partial_y$  can be expressed as  $V = (x_0 \cos \varphi + y_0 \sin \varphi) \partial_r + \frac{1}{r}(y_0 \cos \varphi - x_0 \sin \varphi) \partial_\varphi$ . In particular the coefficient in  $r$ -direction is independent of  $r$ , the one in  $\varphi$ -direction is  $r^{-1}$  times some expression which is independent of  $r$ . This does not only generalize to higher dimensional Euclidean spaces but also to any situation where we have a Liouville transformation and in addition a parallel vector field, see Proposition 3 below. The metric does not have to be flat in this case, compare the example after Proposition 6.

In the sequel we throughout consider *semi-Riemannian manifolds*  $M$  of dimension  $n$  which are sufficiently smooth in order to admit all the derivatives we are using.

**Lemma 1:** *For a conformal transformation  $\bar{g} = \varphi^{-2}g$  of class  $C^2$  the following conditions are equivalent:*

1.  $Ric_{\bar{g}} - Ric_g$  is a scalar multiple of  $g$
2.  $\nabla^2 \varphi$  is a scalar multiple of  $g$
3.  $(\nabla^2 \varphi)^0 = 0$
4.  $\varphi$  is a concircular scalar field in the sense of [36], [11].

*Proof.* The proof follows directly from the following standard formula for the relation between the two Ricci tensors of  $\bar{g}$  and  $g$ :

$$Ric_{\bar{g}} - Ric_g = \varphi^{-2} \left( (n-2) \cdot \varphi \cdot \nabla^2 \varphi + [\varphi \cdot \Delta \varphi - (n-1) \cdot g(\nabla \varphi, \nabla \varphi)] \cdot g \right).$$

The notion of a *concircular scalar field* is motivated by the so-called *concircular transformations* preserving circles. Here a *circle* is defined to be a curve with constant geodesic curvature and vanishing geodesic torsion.  $\square$

**Lemma 2:** *Under the same assumptions the following conditions are equivalent:*

1.  $Ric_{\bar{g}} - Ric_g = 0$
2.  $\nabla^2 \varphi = \frac{\Delta \varphi}{n} \cdot g$  and  $2\varphi \cdot \Delta \varphi = n \cdot g(\nabla \varphi, \nabla \varphi)$
3. *the conformal transformation  $g \mapsto \bar{g}$  is a Liouville transformation (or quasi-homothetic in the sense of [11]).*

Lemma 2 follows from Lemma 1 by taking the trace of the equation  $Ric_{\bar{g}} - Ric_g = 0$ .

By Lemma 2 a Liouville transformation is characterized by the equations  $(\nabla^2 \varphi)^0 = 0$  and  $2\varphi \cdot \Delta \varphi = n \cdot g(\nabla \varphi, \nabla \varphi)$ . From the second equation it becomes immediately clear that we have to distinguish between two cases (unless  $\varphi$  is constant):

- Case 1:  $g(\nabla \varphi, \nabla \varphi) \neq 0$ ,  
Case 2:  $g(\nabla \varphi, \nabla \varphi) = 0$  and  $\nabla \varphi \neq 0$ .

In the second case the gradient of  $\varphi$  is isotropic and necessarily parallel because  $Xg(\nabla \varphi, \nabla \varphi) = 2g(\nabla_X \nabla \varphi, \nabla \varphi) = 0$  and  $g(\nabla_X \nabla \varphi, Y) = 0$  for  $X$  orthogonal to  $\nabla \varphi$ . In the context of conformal mappings between Einstein spaces Case 1 was called *proper* in [5], Case 2 was called *improper*. The results for these two types are quite different.

**Lemma 3:** *In a certain neighborhood of any point with  $g(\nabla\varphi, \nabla\varphi) \neq 0$  the following conditions are equivalent:*

1.  $(\nabla^2\varphi)^0 = 0$
2.  $\nabla\varphi$  is a conformal vector field
3.  $g$  is a warped product metric  $g = \eta dt^2 + (\frac{d\varphi}{dt})^2 g_*$  where  $\eta = \pm 1$  denotes the sign of  $g(\nabla\varphi, \nabla\varphi)$  and where the metric  $g_*$  is independent of  $t$ .

The condition 3. implies that  $\varphi$  is a function only of the real parameter  $t$  and that

$$\nabla\varphi = \frac{d\varphi}{dt} \cdot \frac{\partial}{\partial t} \quad \text{and} \quad \mathcal{L}_{\nabla\varphi}g = 2\nabla^2\varphi = 2\frac{d^2\varphi}{dt^2} \cdot g.$$

*Proof.* Actually Lemma 3 is a standard result, originally due to Brinkmann [5] and Fialkow [12], compare [36], [11], [20].

1.  $\Leftrightarrow$  2. is trivial, 3.  $\Rightarrow$  1. is easily obtained by the calculation of  $\nabla \frac{\partial}{\partial t}$  in terms of Christoffel symbols. For the proof of 1.  $\Rightarrow$  3. one introduces  $\varphi$  as one coordinate function. Then for one particular  $\varphi$ -level  $M_*$  one takes geodesic parallel coordinates. It is easily seen that the parallel levels of the same distance coincide with the  $\varphi$ -levels. If we choose  $t$  as the arclength on the trajectories of  $\nabla\varphi$ , then we obtain a form  $g = \eta dt^2 + g_{ij}(t)$  for the metric,  $i, j = 1, \dots, n-1$ . The proof is completed by calculating  $\frac{\partial}{\partial t}((\frac{d\varphi}{dt})^{-2}g_{ij}(t)) = 0$  from the equation  $\nabla^2\varphi = \frac{d^2\varphi}{dt^2} \cdot g$ . This implies that  $g_{*ij} = (\frac{d\varphi}{dt})^{-2}g_{ij}$  is independent of  $t$ .  $\square$

**Corollary 1:** *In a certain neighborhood of any point with  $g(\nabla\varphi, \nabla\varphi) \neq 0$  the following conditions are equivalent:*

1.  $g \mapsto \varphi^{-2}g$  is a Liouville transformation
2.  $g$  is a warped product metric  $g = \eta dt^2 + (at + b)^2 g_*$  where  $a \neq 0$  and  $b$  are real constants.

In particular, in this case  $\nabla\varphi$  is a homothetic vector field satisfying  $\mathcal{L}_{\nabla\varphi}g = 2a \cdot g$ .

*Proof.* By combining Lemma 3 with Lemma 2, we find that the differential equation for the function  $\varphi(t)$  can be reduced to the ODE  $2\varphi\varphi'' = (\varphi')^2$ . The solutions are precisely the squares of linear functions:  $\varphi(t) = (At + B)^2$  with  $A \neq 0$  since  $\varphi$  is not constant by assumption. Consequently,  $\varphi'(t) = 2A(At + B)$  is linear, say  $\varphi'(t) = at + b$ , and there is a common zero of  $\varphi$  and  $\varphi'$ . The equation  $\nabla^2\varphi = a \cdot g$  is easily verified.  $\square$

In some sense this phenomenon of a common zero of  $\varphi$  and  $\varphi'$  is the crucial point in Liouville's theorem. It leads to the following generalization:

**Theorem 1 (Generalized Liouville Theorem):** *Let  $(M, g)$  be a connected semi-Riemannian manifold and let  $f: (M_1, g) \rightarrow (M, g)$  be a Liouville mapping of class  $C^3$  for some open  $M_1 \subset M$ . Assume that the induced conformal factor  $\varphi$  with  $f^*g = \varphi^{-2}g$  satisfies  $g(\nabla\varphi, \nabla\varphi) \neq 0$  everywhere. Then  $(M_1, g)$  is isometric to an open subset of a warped product  $M_2 = (0, \infty) \times_t M_*$  with a cone-like metric  $g = \eta dt^2 + t^2 g_*$  with  $\eta = \pm 1$  and where  $(M_*, g_*)$  is independent of  $t$ . Furthermore, up to an isometry, in these coordinates the mapping  $f$  appears as  $f(t, x) = (\frac{2}{t}, x)$ . Consequently, the mapping  $f$  is the composition of an isometry and an inversion.*

*If in addition  $M$  is assumed to be Einstein then it is Ricci flat and, consequently, any conformal mapping is a Liouville mapping. In this case it follows that  $(M_*, g_*)$  is an Einstein metric with the same Ricci curvature as a space of constant sectional curvature  $\eta$ . If in addition the dimension of  $M$  is not greater than 4 then  $g$  is flat.*

**Corollary 2:** *Let  $(M, g)$  be a complete Riemannian manifold and assume that there is a Liouville mapping  $f: M \setminus \{p\} \rightarrow M \setminus \{p\}$  for one point  $p \in M$ . Then  $(M, g)$  is isometric with the Euclidean space.*

*Proof. Theorem 1:* From Corollary 1 we obtain a local representation of the metric as  $g = \eta dt^2 + (at + b)^2 g_*$  with  $a \neq 0$ . By a shift of the parameter we may assume that  $b = 0$  and that the factor  $a$  is incorporated into the metric  $g_*$ . This leads to a local expression  $g = \eta dt^2 + t^2 g_*$  in the neighborhood of every point which satisfies the assumptions. Since  $g_*$  is independent of  $t$ , it cannot change on a connected open subset. Therefore there is one maximal "level space"  $(M_*, g_*)$ .



The maximum possible interval for  $t$  is either  $(0, \infty)$  or  $(-\infty, 0)$ . Hence  $M_1$  can be regarded as an open subset of a warped product  $M_2 := (0, \infty) \times_t M_*$  with a cone-like metric. Then the function  $\varphi$  is a function of  $t$  only, more precisely we have  $\varphi(t) = t^2/2$ . The conformal factor of the mapping is  $\varphi^{-2} = 4t^{-4}$ , hence the apex  $t = 0$  of the cone cannot belong to the domain of  $\varphi$  in any case. Note however, that this apex is a removable singularity only if  $(M_*, g_*)$  is isometric with the standard sphere in Euclidean space if  $\eta = 1$  or with a standard quadric in pseudo-Euclidean space if  $\eta = -1$ . In any case by removing the singularity we obtain the (pseudo-)Euclidean metric in geodesic polar coordinates around the origin, see [20]. If  $M$  is assumed to be Ricci flat then we obtain information about the metric  $g_*$  from the equation  $Ric = t^{-2}(Ric_* - (n-2)g_*)$ . In any case the vector field  $\nabla\varphi = t \cdot \frac{\partial}{\partial t}$  is a homothetic vector field satisfying the equation  $\mathcal{L}_{t \frac{\partial}{\partial t}} g = 2\nabla^2\varphi = 2g$ .

Now let us consider the conformal point transformation  $f: M_1 \rightarrow M$  with the conformal factor  $\varphi^{-2}$ . The inversion  $f_0: M_2 \rightarrow M_2$  defined by  $f_0(t, x) = (\frac{2}{t}, x)$  is also conformal with the same conformal factor  $\varphi^{-2} = 4t^{-4}$ . This implies that  $f_0 \circ f^{-1}: f(M) \rightarrow M_2$  and  $f \circ f_0^{-1}: f_0(M_1) \rightarrow M$  are isometric mappings.

Now assume that in addition  $g$  is Einstein. It is well known which Einstein spaces admit a representation as a warped product. In any case the level  $M_*$  has to be an Einstein space, too. In our case the normalized scalar curvatures  $S$  and  $S_*$  of  $g$  and  $g_*$  satisfy the equation  $1 + \eta S t^2 = \eta S_*$ , see [21], Lemma 3.3. This is possible only for  $S = 0$  and  $S_* = \eta$ . A peculiar example for such an Einstein space is given by the non-standard Einstein metrics on spheres (appropriately scaled), see [22]. In this case the cone has no topological singularity at the apex, just a metrical singularity. If the dimension of  $M$  is not greater than 4 then the level metric  $g_*$  is of constant curvature and, therefore,  $g$  is of constant curvature.

*Proof of Corollary 2:* In the Riemannian case we have  $g(\nabla\varphi, \nabla\varphi) \neq 0$  on an open subset  $M'$  unless  $\varphi$  is constant. If  $\varphi$  is globally constant then  $f$  is a homothety, and the assertion is well known. Otherwise, by Theorem 1  $M' \setminus \{p\}$  is an open part of a warped product  $(0, \infty) \times_t M_*$  with  $g = dt^2 + t^2g_*$ . Here  $(M_*, g_*)$  must be complete because otherwise  $g$  would not be complete. Again by the assumption of completeness the apex  $t = 0$  of the cone must be an ordinary (non-singular) point of  $M$ , hence  $M_*$  is a standard unit sphere and  $M$  is the Euclidean space in standard polar coordinates.  $\square$

- REMARKS: 1. There are, however, examples in higher dimensions which are Ricci flat but not flat. Take any 4-dimensional level metric  $g_*$  which is Einstein with scalar curvature  $\eta$  and then take  $M^5 = (0, \infty) \times M_*$  with the metric  $g = \eta dt^2 + t^2 g_*$ .
2. If the isometry mentioned in theorem 1 preserves the  $t$ -levels then it is easily seen to be induced by an isometry of  $M_*$ . Otherwise the metric is of a special type and admits a parallel vector field, see Corollary 4 below. If the metric is Riemannian and if  $M_*$  is complete then this cannot happen unless the metric is flat, see [14]. The case of a conformal mapping moving the apex  $p$  and not preserving the  $t$ -levels can really occur: Consider a Euclidean translation composed by an ordinary inversion at  $p$ . This defines a conformal mapping of  $E^n \setminus \{p\}$  into  $E^n$  which does not preserve the levels of the associated conformal factor. These levels are concentric spheres.
3. In addition to the obvious self-similarity of cones there are other types of self-similarities which do not satisfy the assumptions of Theorem 1, see [1], [7].

**Corollary 3 (Classical Liouville Theorem)** [35, p.209], [2, Thm.A.3.7]:

*Any conformal mapping  $f$  of class  $C^3$  from a connected and open part of Euclidean space into Euclidean space can be written (up to motions in source and target) as a dilatation or as an inversion.*

*Similarly (see [15], [33]), any conformal mapping of class  $C^3$  from a connected and open part of pseudo-Euclidean space into pseudo-Euclidean space can be written (up to motions) as either a dilatation or an inversion or the composition of two distinct inversions.*

*Proof.* Let  $f: U \rightarrow \mathbb{R}^n$  be a conformal mapping with the conformal factor  $\varphi^{-2}$  which is a function of class  $C^2$  on  $U$ . If  $\varphi$  is constant, then the mapping  $f_0(x) = \varphi \cdot f(x)$  defines an isometric mapping  $f_0$  which, consequently, can be globally extended. Hence  $f$  can be written as  $f(x) = \varphi^{-1} \cdot f_0(x)$ , a composition of a (pseudo-)Euclidean motion with a dilatation. If  $\varphi$  is not constant, then we have  $\nabla\varphi \neq 0$  on an open set. If  $\nabla\varphi$  is not a null vector then we can apply Theorem 1. The result is that in certain coordinates the mapping can be written as an inversion at a certain sphere. If  $\nabla\varphi$  is a (parallel) null vector field on an open set then we form the compositions  $j \circ f$  and  $f \circ j$  of  $f$  with an inversion  $j$ . For

a generic  $j$  the gradients of  $j \circ f$  and of  $f \circ j$  are no longer parallel on open sets. Therefore we can apply Theorem 1 to  $j \circ f$  and  $f \circ j$ . This proves Corollary 3.  $\square$

**Proposition 1:** *If  $\varphi : M \rightarrow \mathbb{R}$  is a function satisfying  $\nabla^2\varphi = c \cdot g$  with a constant  $c \neq 0$  (i.e.,  $\nabla\varphi$  is homothetic) and if  $F: M \rightarrow M$  is an isometry which does not preserve  $\varphi$  (i.e.,  $\varphi \circ F \neq \varphi$ ) then  $M$  carries a parallel vector field.*

*Proof.* Let  $\psi = \varphi \circ F$ . Then we have  $\nabla\psi|_p = DF^{-1}|_p(\nabla\varphi|_{F(p)})$  and  $\Delta\psi|_p = \Delta\varphi|_{F(p)}$ . It follows that  $\nabla^2\psi = c \cdot g$  and, consequently,  $\nabla^2(\psi - \varphi) = 0$ . Hence  $\nabla(\psi - \varphi)$  is parallel vector field. It remains to show that it does not vanish. If  $\nabla\psi = \nabla\varphi$  then we have  $2\varphi(F(p)) \cdot c = \frac{2}{n}\varphi(F(p))\Delta\varphi(F(p)) = g(\nabla\varphi|_{F(p)}, \nabla\varphi|_{F(p)}) = g(\nabla\psi|_p, \nabla\psi|_p) = g(\nabla\varphi|_p, \nabla\varphi|_p) = \frac{2}{n}\varphi(p)\Delta\varphi(p) = 2\varphi(p) \cdot c$  since  $F$  is an isometry. Consequently  $\nabla\psi = \nabla\varphi$  implies that  $\psi = \varphi$  which was excluded by our assumption.  $\square$

**Proposition 2:** *Assume that a cone metric  $g = \eta dt^2 + t^2 g_*$  is given. Then locally the following conditions are equivalent:*

1. *There exists a parallel vector field  $V$*
2. *On the level  $M_*$  there exists a non-constant function  $\alpha$  with  $g_*(\nabla_*\alpha, \nabla_*\alpha) \neq 0$  which satisfies the equation  $\nabla_*^2\alpha = -\eta\alpha g_*$ .*

*Proof.* (1)  $\Rightarrow$  (2) Let  $V$  be a parallel vector field. We can decompose it into  $V(t, x) = \alpha(t, x)\partial_t + W(t, x)$  where  $W$  is orthogonal to  $\partial_t$  and where  $\alpha$  is not the zero function because  $V$  cannot be orthogonal on every  $t$ -line simultaneously. Since the  $t$ -lines are geodesics,  $V$  has a constant angle with  $\partial_t$  along them. Consequently,  $\alpha$  does not depend on  $t$ . So we can regard  $\alpha$  as a function on any of the levels. From  $\nabla_{\partial_t}V = 0$  we obtain  $\nabla_{\partial_t}W = 0$ . Therefore  $W(t, x)$  is parallel along the  $t$ -lines, and we can set  $W(t, x) = \frac{1}{t}W(x)$  where  $W(x)$  is the lift of a vector field on  $M_*$ . Altogether we obtain  $V(t, x) = \alpha(x)\partial_t + \frac{1}{t}W(x)$ .

Next, if  $X$  denotes the lift of a vector field on  $M_*$ , we have for any fixed  $t$

$$\begin{aligned}
\nabla_X^* W &= \nabla_X W + g(X, W)t^{-1}\eta\partial_t \\
&= \nabla_X(tV(t, x) - t\alpha(x)\partial_t) + g_*(X, W)t\eta\partial_t \\
&= -tX(\alpha)\partial_t - t\alpha(x)\nabla_X\partial_t + g_*(X, W)t\eta\partial_t \\
&= t(g_*(X, W)\eta - g_*(\nabla^*\alpha, X))\partial_t - t\alpha(x)t^{-1}X.
\end{aligned}$$

Since the  $\partial_t$ -component of  $\nabla_X^* W$  must vanish for any  $X$  we obtain first the equation

$$\nabla^*\alpha = \eta W$$

and, consequently,

$$\nabla_X^* \nabla^*\alpha = \eta \nabla_X^* W = -\eta\alpha(x) \cdot X$$

which is the assertion. In particular,  $\alpha$  is not constant. Moreover, since

$$g(V, V) = \eta\alpha^2 + g_*(W, W) = \eta\alpha^2 + g_*(\nabla_*\alpha, \nabla_*\alpha)$$

is constant, it follows that  $g_*(\nabla_*\alpha, \nabla_*\alpha)$  cannot vanish on any open set. Consequently,  $\|W\|^2$  cannot vanish either.

For the converse direction (2)  $\Rightarrow$  (1) we assume that  $\alpha$  satisfies this differential equation and define  $V$  by the equations above. It then follows that  $V$  is parallel.  $\square$

**Corollary 4:** *If a cone metric  $g = \eta dt^2 + t^2 g_*$  carries a parallel vector field then  $g_*$  can be written as a warped product  $g_* = \epsilon du^2 + \alpha'^2(u)g_{**}$  where  $\epsilon = \pm 1$  and the function  $\alpha(u)$  satisfies the equation  $\alpha'' + \epsilon\eta\alpha = 0$ . Conversely, if a metric of the form  $g = \eta dt^2 + t^2(\epsilon du^2 + \alpha'^2(u)g_{**})$  with such an  $\alpha$  is given then  $g$  carries a parallel vector field.*

This follows from Lemma 3 in connection with Proposition 2. Along the trajectories of the gradient of  $\alpha$  is not a null vector. With  $\nabla_*\alpha = \epsilon\alpha'(u)\partial_u$  the function satisfies the ODE  $\alpha'' + \epsilon\eta\alpha = 0$ . The solutions are linear combinations of  $\sin, \cos$  if  $\epsilon\eta = 1$  and of  $\sinh, \cosh$  if  $\epsilon\eta = -1$ .

**Corollary 5** (see [14] ):

*If  $g = dt^2 + t^2g_*$  is positive definite and if  $g_*$  is complete then  $g$  is either irreducible or flat.*

*Proof.* This follows from the well known theorem of Tashiro [36] and Obata [31] that on a complete Riemannian manifold  $(M_*, g_*)$  the equation  $\nabla_*^2 \alpha = -\alpha g$  admits a non-constant solution only if the manifold is a round sphere of radius 1. So if  $g$  carries a parallel vector field then this equation on  $(M_*, g_*)$  is satisfied by Proposition 2. It follows that the level  $M_*$  is the unit sphere and, consequently,  $g$  is flat.  $\square$

**Corollary 6:** *Assume that the cone metric with  $g = \eta dt^2 + t^2g_*$  is an Einstein  $n$ -manifold (necessarily Ricci flat) and carries a parallel vector field. Then for  $n \leq 5$   $g$  is necessarily flat. In contrast, there is an example of a 6-dimensional Ricci flat cone which is not flat.*

**Example:** Let  $h$  be a complete Ricci flat space with a positive definite metric which is not flat (one example is the Calabi-Yau metric on the  $K3$  surface). Then the Lorentzian warped product metric

$$g = -dt^2 + t^2(du^2 + e^{2u}h)$$

is Ricci flat but not flat. It carries the homothetic vector field  $t\partial_t$  and, in addition, the parallel null vector field  $V = e^u(\partial_t + \frac{1}{t}\partial_u)$ . The level  $g_* = du^2 + e^{2u}h$  is complete if  $h$  is.

**Lemma 4:** *Assume that an  $n$ -dimensional manifold  $(M, g)$  admits a Liouville mapping such that the conformal factor  $\varphi$  has a gradient  $\nabla\varphi \neq 0$  everywhere. Then the following conditions are equivalent:*

1.  $g(\nabla\varphi, \nabla\varphi) = 0$  everywhere,
2.  $\nabla^2\varphi = 0$  everywhere,
3.  $\nabla\varphi$  is a parallel vector field,

4. in a suitable coordinate system the metric  $g$  can be written as  $g = 2d\varphi d\vartheta + g^\#(\varphi)$  where  $g^\#(\varphi)$  is an  $(n - 2)$ -dimensional metric which is independent of  $\vartheta$ .

*Proof.* 4.  $\Rightarrow$  1. holds because in these coordinates we have  $\frac{\partial}{\partial\vartheta} = \nabla\varphi$ .

1.  $\Leftrightarrow$  2. holds by Lemma 2,

2.  $\Leftrightarrow$  3. is trivial.

The main part is the proof of 1.  $\Rightarrow$  4 which is originally due to Brinkmann in [5, p.132-133], for a modern treatment see also [6, Sect.3.3]. Because  $\varphi$  is nowhere constant we can introduce  $\varphi$  as one of the coordinates. Then  $\nabla\varphi$  is a parallel vector field by 3., therefore we can introduce a second coordinate function  $\vartheta$  by the condition  $\frac{\partial}{\partial\vartheta} = \nabla\varphi$ . It follows that  $g(\frac{\partial}{\partial\varphi}, \frac{\partial}{\partial\vartheta}) = 1$  and, in particular, that  $\frac{\partial}{\partial\varphi}$  and  $\frac{\partial}{\partial\vartheta}$  are linearly independent everywhere. Now let us fix one particular  $\varphi$ - $\vartheta$ -level with a non-degenerate induced metric  $g_\#(\varphi, \vartheta)$ . Then introduce geodesic normal coordinates  $\varphi, \vartheta, x_1, \dots, x_{n-2}$  around this level. It follows that the metric  $g^\#$  is independent of  $\vartheta$  because  $\frac{\partial}{\partial\vartheta}$  is parallel. By a transformation of the metric one can assume that  $g(\frac{\partial}{\partial\varphi}, \frac{\partial}{\partial\varphi})$  is also zero. This leads to the expression  $g = 2d\varphi d\vartheta + g^\#(\varphi)$  for the metric.  $\square$

**Corollary 7** *If the conditions in Lemma 4 are satisfied then the  $\vartheta$ -lines (or the trajectories generated by the parallel vector field  $\nabla\varphi$ ) are null geodesics. The family of all of them forms a so-called null congruence. The Liouville mapping transforms this null congruence into a (possibly different) null congruence.*

*Proof.* We have to show that the parallel isotropic vector  $\nabla\varphi$  is mapped by the Liouville mapping onto a parallel isotropic vector. This follows directly from the formula

$$\bar{\nabla}_X V - \nabla_X V = -X(\log \varphi)V - V(\log \varphi)X + g(X, V)\nabla(\log \varphi)$$

for the conformal change of the metric  $\bar{g} = \varphi^{-2}g$  and any  $X, V$ . In our case we have  $V = \nabla\varphi$  which implies  $V(\log \varphi) = 0$  and  $X(\log \varphi)\nabla\varphi = g(X, \nabla\varphi)\nabla(\log \varphi)$ .  $\square$

**Lemma 5** *If an  $n$ -dimensional manifold  $(M, g)$  carries two linearly independent null congruences then it is (locally) reducible to a product  $(\mathbb{R}^2 \times M_*, 2dudv + g_*)$ . Moreover, the transition from one null congruence to the other can be realized by an isometry.*

This follows because the two linearly independent isotropic vector fields generate a 2-dimensional distribution which is integrable and parallel. Therefore, a 2-dimensional factor splits off which is a flat Lorentzian plane.

In view of Lemma 5 and up to an isometry, one has to consider only the case that the null congruence is preserved.

**Theorem 2 (The isotropic case in Liouville's theorem):** *Assume that an  $n$ -dimensional manifold  $(M, g)$  admits a Liouville mapping  $F: M \rightarrow M$  such that the conformal factor  $\varphi$  has an isotropic gradient  $\nabla\varphi \neq 0$  everywhere. Assume further that  $F$  preserves the null-congruence given by the parallel null vector  $\partial_v = \nabla\varphi$ . Then in certain coordinates  $u, v, x_k$  ( $k = 1, \dots, n-2$ ) the metric has the form*

$$g = -2dudv + \sum_{i,j} g_{ij}^\#(u, x_k) dx^i dx^j$$

and, up to an isometry,  $F$  has the form

$$F(u, v, x_k) = \left( -\frac{1}{cu}, cv + \zeta(u, x_k), \xi_1(u, x_k), \dots, \xi_{n-2}(u, x_k) \right)$$

with a constant  $c$  and with a certain function  $\zeta$ , where for any fixed  $u, v$  the transformation

$$(x_1, \dots, x_{n-2}) \mapsto (\xi_1, \dots, \xi_{n-2})$$

is a homothety with respect to the metric  $g^\#$ . The conformal factor of  $F$  is the function  $\varphi(u, v, x_k) = u$ , i.e.,  $F^*g = u^{-2}g$ .

Conversely, Let  $h$  be any metric on an  $(n-2)$ -dimensional space  $M_*$  admitting a 1-parameter group  $\Phi_u$  of similarities (homothetic transformations) with  $\Phi_u^*h = u^{-2}h$ . Then on  $M = \mathbb{R}_+ \times \mathbb{R} \times M_*$  the metric  $g = -2dudv + \Phi_u^*h$  admits a conformal mapping  $F$  such that the conformal factor  $u$  has an isotropic gradient  $\nabla u = \partial_v$ . In this case  $F$  acts on  $M_*$  by the similarities  $\Phi_u$ .

*Proof.* The particular form of the metric is obtained by Lemma 4 which also determines the conformal factor as  $\varphi = u$ . We compute the differential of  $F$  with respect to the basis  $\partial_u, \partial_v, \partial_1, \dots, \partial_{n-2}$  as follows. By assumption  $F$  preserves the null-congruence given by the  $v$ -lines. This implies that at every point the differential of  $F$  transforms the vector  $\partial_v$  into some vector linearly dependent on  $\partial_v$ . This implies that the component functions  $\kappa, \xi_1, \dots, \xi_{n-2}$  in

$$F(u, v, x_1, \dots, x_{n-2}) = (\kappa, \lambda, \xi_1, \dots, \xi_{n-2})$$

are independent of the variable  $v$ . Consequently, we have

$$\begin{aligned} F_u &= (\kappa_u, \lambda_u, (\xi_1)_u, \dots, (\xi_{n-2})_u) \\ F_v &= (0, \lambda_v, 0, \dots, 0) \\ F_k &= (\kappa_k, \lambda_k, (\xi_1)_k, \dots, (\xi_{n-2})_k) \text{ for } k = 1, \dots, n-2 \end{aligned}$$

where the indices indicate the partial derivatives with respect to the corresponding coordinates.

From the equation  $g(F_u, F_v) = u^{-2}g(\partial_u, \partial_v) = -u^{-2}$  we obtain  $\kappa_u \lambda_v = u^{-2}$ . Differentiating once more leads to the equation  $\kappa_u \lambda_{vv} = 0$ .

Similarly, from the equation  $g(F_k, F_v) = u^{-2}g(\partial_k, \partial_v) = 0$  we obtain  $\kappa_k \lambda_v = 0$ . Since  $\lambda_v$  cannot vanish identically, we get  $\kappa_k = 0$  for all  $k$ . As a consequence,  $\kappa_u$  does not vanish identically. It follows that  $\lambda_{vv}$  must vanish identically. Hence  $\lambda_v = c$  is constant and  $\kappa_u = \frac{1}{cu^2}$  and  $\kappa = -\frac{1}{cu}$ . It also follows that  $\lambda = cv + \eta(u, x_k)$  with some function  $\eta$ .

Now let  $u_0$  and  $v_0$  be fixed, and consider the mapping  $\Phi$  defined by

$$(x_1, \dots, x_{n-2}) \mapsto (\xi_1(u_0, x_1, \dots, x_{n-2}), \dots, \xi_{n-2}(u_0, x_1, \dots, x_{n-2})).$$

From the equation

$$\Phi^* g^\#(\partial_i, \partial_j) = g^\#(F_i, F_j) = g(F_i, F_j) = u_0^{-2}g(\partial_i, \partial_j) = u_0^{-2}g^\#(\partial_i, \partial_j)$$

we see that  $\Phi$  is homothetic.

Conversely, if  $(M_*, g_*)$  is given with a 1-parameter group  $\Phi_u$  of similarities then we can define  $F$  by

$$F(u, v, \mathbf{x}) = \left( -\frac{1}{cu}, cv + \zeta(u, \mathbf{x}), \Phi_u(\mathbf{x}) \right).$$



The function  $\zeta$  is determined by the condition that  $F_u$  must be orthogonal to the derivative of  $F$  by any coordinate in the  $(n - 2)$ -dimensional space  $M_*$  and, furthermore, that  $g(F_u, F_u) = 0$ .  $\square$

Remark: In the case of a vacuum spacetime our extra assumption on preserving the null-congruence is always satisfied unless the manifold is flat.

**Examples:** 1. Let  $g = -2dudv + \sum_i \epsilon_i x_i^2$  be the metric of the semi-euclidean space. Then the following mapping is a conformal involution:

$$F(u, v, x_1, \dots, x_{n-2}) = (-u^{-1}, v + (2u)^{-1} \sum_i \epsilon_i x_i^2, u^{-1}x_1, \dots, u^{-1}x_{n-2}).$$

A local 1-parameter group of conformal mappings is

$$\Phi_t(u, v, x_1, \dots, x_{n-2}) = \frac{1}{1-tu}(u, v(1-tu) + \frac{t}{2} \sum_i \epsilon_i x_i^2, x_1, \dots, x_{n-2}).$$

The expression in the first coordinate differs from the form given in Theorem 2 but that is just up to a conjugation with a shift  $u \mapsto u + u_0$ . It generates the following conformal vector field

$$\frac{d}{dt}\Phi_t = (u^2, \frac{1}{2} \sum_i \epsilon_i x_i^2, ux_1, \dots, ux_{n-2})$$

which is a standard special conformal vector field also in the general context of  $pp$ -waves, see [23]. Here it can be considered as a Liouville vector field where every member of the 1-parameter family is of type 3.

2. Let  $g = -2dudv + u^2 \sum_i \epsilon_i x_i^2$ . Then the following involution is conformal:

$$F(u, v, x_1, \dots, x_{n-2}) = (-u^{-1}, v + \frac{u}{2} \sum_i \epsilon_i x_i^2, ux_1, \dots, ux_{n-2}).$$

3. More generally, let  $h$  be any metric and let  $g = -2dudv + dt^2 + t^2h$ . Then the following involution is conformal:

$$F(u, v, t, x_k) = (-u^{-1}, v - \frac{1}{2}u^{-1}t^2, u^{-1}t, x_k).$$

Similarly, if  $g = -2dudv + u^2(dt^2 + t^2h)$ . Then the following involution is conformal:

$$F(u, v, t, x_k) = (-u^{-1}, v + \frac{u}{2}t^2, tu, x_k).$$

Acknowledgement: This work was partially supported by the Erwin Schrödinger International Institute for Mathematical Physics at Vienna.

## References

- [1] D. V. Alekseevskii: *Selfsimilar Lorentzian manifolds*, Ann. Glob. Anal. Geom. **3** (1985), 59–84
- [2] R. Benedetti and V. Petronio: *Lectures on hyperbolic geometry*. Springer, Berlin 1992.
- [3] W. Benz: *Der Satz von Liouville über räumliche orthogonaltreue (winkeltreue) Abbildungen für beliebige Signatur*, Aequationes Math. **21** (1980), 257–282
- [4] B. Bojarski and T. M. Iwaniec: *Another approach to Liouville theorem*, Math. Nachr. **107** (1982), 253–262
- [5] H. W. Brinkmann: *Einstein spaces which are mapped conformally on each other*, Math. Ann. **94** (1925), 119–145
- [6] D. A. Catalano: *Concircular diffeomorphisms of pseudo-Riemannian manifolds*. Doctoral thesis ETH Zürich 1999
- [7] A. A. Coley: *Self-Similarity and Cosmology*, Fields Institute Communications **15** (1997), 19–42
- [8] K. L. Duggal and R. Sharma: *Symmetries of Spacetimes and Riemannian Manifolds*, Kluwer, Dordrecht 1999
- [9] D. Eardley, J. Isenberg, J. Marsden and V. Moncrief: *Homothetic and conformal symmetries of solutions to Einstein's equations*, Commun. Math. Phys. **106** (1986), 137–158
- [10] J. Ferrand: *Sur une classe de morphismes conformes*, C. R. Acad. Sci. Paris **296** (1983), 681–684
- [11] J. Ferrand: *Concircular transformations of Riemannian manifolds*, Ann. Acad. Sci. Fenn., Ser. A.I. Math. **10** (1985), 163–171
- [12] A. Fialkow: *Conformal geodesics*, Trans. Amer. Math. Soc. **45** (1939), 443–473
- [13] C. Frances, *Une preuve du théorème de Liouville en géométrie dans le cas analytique*, L'Enseign. Math. (2) **49** (2003), 95–100
- [14] S. Gallot: *Équations différentielles caractéristiques de la sphère*, Ann. Scient. Ec. Norm. Sup (4) **12** (1979), 235–267

- [15] J. Haantjes: *Conformal representations of an  $n$ -dimensional euclidean space with a non-definite fundamental form on itself*, Proc. Kon. Ned. Akad. Amsterdam **40** (1937), 700–705
- [16] G. S. Hall: *Symmetries and Curvature Structure in General Relativity*, World Scientific, Singapore 2004
- [17] H. Jacobowitz, *Two notes on conformal geometry*, Hokkaido Math. J. **20** (1991), 313–329
- [18] Y. Kerbrat: *Transformations conformes des variétés pseudo-Riemanniennes*. J. Diff. Geometry **11** (1976), 547–571
- [19] W. Kühnel and H.-B. Rademacher: *Conformal diffeomorphisms preserving the Ricci tensor*, Proc. Amer. Math. Soc. **123** (1995), 2841–2848
- [20] W. Kühnel and H.-B. Rademacher: *Essential conformal fields in pseudo-Riemannian geometry*, J. Math. Pures Appl. (9) **74** (1995), 453–481, Part II: J. Math. Sci. Univ. Tokyo **4** (1997), 649–662
- [21] W. Kühnel and H.-B. Rademacher: *Conformal vector fields on pseudo-Riemannian spaces*, Diff. Geom. Appl. **7** (1997), 237–250
- [22] W. Kühnel and H.-B. Rademacher: *Conformal Ricci collineations of space-times*. Gen. Rel. Grav. **33** (2001), 1905–1914
- [23] W. Kühnel and H.-B. Rademacher: *Conformal geometry of gravitational plane waves*, Geom. Dedicata **109** (2004). 175–188
- [24] N. H. Kuiper: *On conformally-flat spaces in the large*, Ann. of Math. (2) **50** (1949), 916–924
- [25] R. S. Kulkarni: *Conformal structures and Möbius structures*, in: *Conformal Geometry* (R.S.Kulkarni and U.Pinkall, eds.), pp. 1–39, Aspects Math. E 12, Vieweg, Wiesbaden 1988
- [26] J. Lafontaine: *Sur la géométrie d'une généralisation de l'équation différentielle d'Obata*. J. Math. Pures Appl. (9) **62** (1983), 63–72
- [27] S. Lie: *Ueber Complexe, insbesondere Linien- und Kugel-Complexe mit Anwendung auf die Theorie partieller Differentialgleichungen*, Math. Ann. **5** (1872), 145–246

- [28] J. Liouville: *Extension au cas des trois dimensions de la question du tracé géographique*, Note VI, pp. 609 - 617 in: G. Monge: *Applications de l'analyse à la géométrie*, Bachelier, Paris 1850
- [29] R. Maartens and S. D. Maharaj: *Conformal symmetries of pp-waves*, *Class. Quantum Grav.* **8** (1991), 503–514
- [30] W. Moore and J. D. Zund: *A note on Liouville's theorem in conformal geometry*, *Tensor (N.S.)* **44** (1987), 204–207
- [31] M. Obata: *Certain conditions for a Riemannian manifold to be isometric with a sphere*. *J. Math. Soc. Japan* **14** (1962), 333–340
- [32] Yu. G. Reshetnyak: *Liouville's theorem on conformal mappings for minimal regularity assumptions*, *Sib. Math. J.* **8** (1967), 631–634, transl. from *Sib. Mat. Zh.* **8** (1967), 835–840
- [33] M. Schottenloher: *A Mathematical Introduction to Conformal Field Theory*, Springer, Berlin 1997
- [34] J. A. Schouten: *Der Ricci-Kalkül*, Springer, Berlin 1924
- [35] J. A. Schouten und D. J. Struik: *Einführung in die neueren Methoden der Differentialgeometrie*, Band 2, Noordhoff, Groningen 1938
- [36] Y. Tashiro: *Complete Riemannian manifolds and some vector fields*, *Trans. Amer. Math. Soc.* **117** (1965), 251–275

Wolfgang Kühnel  
 Institut für Geometrie und Topologie  
 Universität Stuttgart  
 Pfaffenwaldring 57  
 70550 Stuttgart  
 Germany

Hans-Bert Rademacher  
 Mathematisches Institut  
 Universität Leipzig  
 Augustusplatz 10/11  
 04109 Leipzig  
 Germany

kuehnel@mathematik.uni-stuttgart.de

rademacher@mathematik.uni-leipzig.de