

RECONSTRUCTION OF THE TRANSMISSION COEFFICIENT FOR STEPLIKE FINITE-GAP BACKGROUNDS

IRYNA EGOROVA AND GERALD TESCHL

ABSTRACT. We consider scattering theory for one-dimensional Jacobi operators with respect to steplike quasi-periodic finite-gap backgrounds and show how the transmission coefficient can be reconstructed from minimal scattering data. This generalizes the Poisson–Jensen formula for the classical constant background case.

1. INTRODUCTION

In classical one-dimensional scattering theory the transmission coefficient can be reconstructed from the reflection coefficient via the well-known Poisson–Jensen formula. This formula plays a crucial role in inverse scattering theory since it shows how to compute the left scattering data from the right one and vice versa. Moreover, it is also one of the key ingredients for deriving the associated sum rules which have attracted an enormous amount of interest recently (see e.g. [17], [21], [23], [24], [31]). Furthermore, these sum rules are intimately connected with conserved quantities of the associated completely integrable lattices (see [26], [27]). Finally, the reconstruction formula can be viewed as the solution of a scalar Riemann–Hilbert factorization problem which arises in the nonlinear steepest descent method [5] when deriving the long-time asymptotics (see [13], [18], respectively [19]).

Moreover, the same is true in case of scattering with respect to a finite-gap background [6], [30]. In this situation the analogous formula was given in [28] including the associated sum rules (see also [7], [9], [22]). Again there is a close relation with the solution of a scalar Riemann–Hilbert factorization problem on the underlying Riemann surface which arises in the nonlinear steepest descent method [14], [15], [16], and [20].

However, while scattering theory with a steplike constant background is classical and goes back to the early sixties [4] (see [2] for the most recent results), even in this case the reconstruction formula is unknown to the best of our knowledge except for the case when the two spectra overlap. This might be related to the fact that the case where the two spectra do not overlap will not be solved in terms of elementary methods, but will already require tools from the theory of elliptic surfaces, as we will see below. In case of steplike finite-gap backgrounds, scattering theory is again well-understood by now [8], [10], however, the reconstruction formula is again unknown to the best of our knowledge (except for the case of two finite-gap backgrounds in the same isospectral class [8]). The main purpose of our paper is to fill this gap

1991 *Mathematics Subject Classification.* Primary 30E20, 30F30; Secondary 34L25, 47B36.

Key words and phrases. Jacobi operators, scattering theory, periodic, Abelian integrals.

Supported by Austrian Science Fund (FWF) under Grant No. Y330.

Oper. Matrices **3**, 205–214 (2009).

and provide a reconstruction formula for the left/right transmission coefficient in terms of the left/right scattering data.

2. NOTATION

We begin by introducing some required background from the theory of hyperelliptic curves to be used in the remainder of this article. For further information and proofs we refer for instance to [1], [3], [11], [12], or [27].

Let \mathbb{M} be the Riemann surface associated with the function $P^{1/2}(z)$, where

$$(2.1) \quad P(z) = \prod_{j=0}^{2g+1} (z - E_j), \quad E_0 < E_1 < \cdots < E_{2g+1},$$

$g \in \mathbb{N}$. \mathbb{M} is a compact, hyperelliptic Riemann surface of genus g . We will choose $P^{1/2}(z)$ as the fixed branch

$$(2.2) \quad P^{1/2}(z) = - \prod_{j=0}^{2g+1} \sqrt{z - E_j},$$

where $\sqrt{\cdot}$ is the standard root with branch cut along $(-\infty, 0)$.

A point on \mathbb{M} is denoted by $p = (z, \pm P^{1/2}(z)) = (z, \pm)$, $z \in \mathbb{C}$. The two points at infinity are denoted by $p = \infty_{\pm}$. We use $\pi(p) = z$ for the projection onto the extended complex plane $\mathbb{C} \cup \{\infty\}$. The points $\{(E_j, 0), 0 \leq j \leq 2g+1\} \subseteq \mathbb{M}$ are called branch points and the sets

$$(2.3) \quad \Pi_{\pm} = \{(z, \pm P^{1/2}(z)) \mid z \in \mathbb{C} \setminus \Sigma\} \subset \mathbb{M}, \quad \Sigma = \bigcup_{j=0}^g [E_{2j}, E_{2j+1}],$$

are called upper and lower sheet, respectively. Note that the boundary of Π_{\pm} consists of two copies of Σ corresponding to the two limits from the upper and lower half plane.

Let $\{a_j, b_j\}_{j=1}^g$ be loops on the Riemann surface \mathbb{M} representing the canonical generators of the fundamental group $\pi_1(\mathbb{M})$. We require a_j to surround the points E_{2j-1}, E_{2j} (thereby changing sheets twice) and b_j to surround E_0, E_{2j-1} counter-clockwise on the upper sheet, with pairwise intersection indices given by

$$(2.4) \quad a_j \circ a_k = b_j \circ b_k = 0, \quad a_j \circ b_k = \delta_{jk}, \quad 1 \leq j, k \leq g.$$

The corresponding canonical basis $\{\zeta_j\}_{j=1}^g$ for the space of holomorphic differentials can be constructed by

$$(2.5) \quad \zeta = \sum_{j=1}^g \underline{c}(j) \frac{\pi^{j-1} d\pi}{P^{1/2}},$$

where the constants $\underline{c}(\cdot)$ are given by

$$c_j(k) = C_{jk}^{-1}, \quad C_{jk} = \int_{a_k} \frac{\pi^{j-1} d\pi}{P^{1/2}} = 2 \int_{E_{2k-1}}^{E_{2k}} \frac{z^{j-1} dz}{P^{1/2}(z)} \in \mathbb{R}.$$

The differentials fulfill

$$(2.6) \quad \int_{a_j} \zeta_k = \delta_{j,k}, \quad \int_{b_j} \zeta_k = \tau_{j,k}, \quad \tau_{j,k} = \tau_{k,j}, \quad 1 \leq j, k \leq g.$$

For further information we refer to [11], [27, App. A].

In addition, we will need Green's function (in the potential theoretic sense) of the upper sheet Π_+ :

Lemma 2.1 ([28]). *The Green function of Π_+ with pole at z_0 is given by*

$$(2.7) \quad g(z, z_0) = -\operatorname{Re} \int_{E_0}^p \omega_{p_0 \bar{p}_0}, \quad p = (z, +), \quad p_0 = (z_0, +),$$

where $\bar{p}_0 = \overline{p_0}^*$ (i.e., the complex conjugate on the other sheet) and ω_{pq} is the normalized Abelian differential of the third kind with poles at p and q .

Clearly, we can extend $g(z, z_0)$ to a holomorphic function on $\mathbb{M} \setminus \{p_0\}$ by dropping the real part. By abuse of notation we will denote this function by $g(p, p_0)$ as well. However, note that $g(p, p_0)$ will be multivalued with jumps in the imaginary part across b -cycles. We will choose the path of integration in $\mathbb{C} \setminus [E_0, E_{2g+1}]$ to guarantee a single-valued function.

From the Green function we obtain the Blaschke factor (cf. [28])

$$(2.8) \quad B(p, \rho) = \exp\left(g(p, \rho)\right) = \exp\left(\int_{E_0}^p \omega_{\rho \rho^*}\right), \quad \pi(\rho) \in \mathbb{R},$$

which has the following properties:

Lemma 2.2. *The Blaschke factor satisfies*

$$(2.9) \quad B(E_0, \rho) = 1, \quad \text{and} \quad B(p^*, \rho) = B(p, \rho^*) = B(p, \rho)^{-1};$$

it is real-valued for $\pi(p) \in (-\infty, E_0)$.

Moreover,

$$(2.10) \quad |B(p, \rho)| = 1, \quad p \in \Sigma, \quad \arg(B(p, \rho)) = \delta_j(\rho), \quad \pi(p) \in [E_{2j-1}, E_{2j}],$$

where we set $E_{-1} = -\infty$, $E_{2g+2} = \infty$ and

$$(2.11) \quad \delta_j(\rho) = \begin{cases} 0, & j = 0, \\ \frac{1}{2} \int_{b_j} \omega_{\rho \rho^*}, & j = 1, \dots, g, \\ 0, & j = g + 1. \end{cases}$$

Proof. The first part including the fact that $|B(p, \rho)| = 1$, $p \in \Sigma$, is proven in [28, Lem. 3.3]. To see the formula for the argument first observe that $\omega_{\rho \rho^*}$ is real-valued on $\pi^{-1}(\mathbb{R} \setminus \Sigma)$ and purely imaginary on $\pi^{-1}(\Sigma)$. This can be seen from the explicit expression (2.13) for $\omega_{\rho \rho^*}$ given below. Hence, taking the path of integration along the lift of the real axis, we see that the integral is real for $p = (\lambda, \pm)$ with $\lambda < E_0$ or $\lambda > E_{2g+1}$. For $p = (\lambda, \pm)$ with $\lambda \in [E_{2j-1}, E_{2j}]$ the imaginary part is constant and given by half the b_j period. \square

The above Abelian differential is explicitly given by

$$(2.12) \quad \omega_{pq} = \left(\frac{P^{1/2} + P^{1/2}(p)}{2(\pi - \pi(p))} - \frac{P^{1/2} + P^{1/2}(q)}{2(\pi - \pi(q))} + P_{pq}(\pi) \right) \frac{d\pi}{P^{1/2}},$$

where $P_{pq}(z)$ is a polynomial of degree $g - 1$ which has to be determined from the normalization $\int_{a_\ell} \omega_{pp^*} = 0$. In particular,

$$(2.13) \quad \omega_{pp^*} = \left(\frac{P^{1/2}(p)}{\pi - \pi(p)} + P_{pp^*}(\pi) \right) \frac{d\pi}{P^{1/2}}.$$

3. RECONSTRUCTING THE TRANSMISSION COEFFICIENT

Let H_q^\pm be two quasi-periodic finite-band Jacobi operators,¹

$$(3.1) \quad H_q^\pm f(n) = a_q^\pm(n)f(n+1) + a_q^\pm(n-1)f(n-1) + b_q^\pm(n)f(n), \quad f \in \ell^2(\mathbb{Z}),$$

associated with the hyperelliptic Riemann surface of the square root

$$(3.2) \quad P_\pm^{1/2}(z) = - \prod_{j=0}^{2g_\pm+1} \sqrt{z - E_j^\pm}, \quad E_0^\pm < E_1^\pm < \dots < E_{2g_\pm+1}^\pm,$$

where $g_\pm \in \mathbb{N}$ and $\sqrt{\cdot}$ is the standard root with branch cut along $(-\infty, 0)$. In fact, H_q^\pm are uniquely determined by fixing a Dirichlet divisor $\sum_{j=1}^{g_\pm} \hat{\mu}_j^\pm$, where $\hat{\mu}_j^\pm = (\mu_j^\pm, \sigma_j^\pm)$ with $\mu_j^\pm \in [E_{2j-1}^\pm, E_{2j}^\pm]$ and $\sigma_j^\pm \in \{-1, 1\}$. The spectra of H_q^\pm consist of $g_\pm + 1$ bands

$$(3.3) \quad \sigma_\pm := \sigma(H_q^\pm) = \bigcup_{j=0}^{g_\pm} [E_{2j}^\pm, E_{2j+1}^\pm].$$

We are interested in scattering theory for the operator

$$(3.4) \quad Hf(n) = a(n-1)f(n-1) + b(n)f(n) + a(n)f(n+1), \quad n \in \mathbb{Z},$$

whose coefficients are asymptotically close to the coefficients of H_q^\pm on the corresponding half-axes:

$$(3.5) \quad \sum_{n=0}^{\pm\infty} |n| \left(|a(n) - a_q^\pm(n)| + |b(n) - b_q^\pm(n)| \right) < \infty.$$

Let $\psi_q^\pm(z, n)$ be the Floquet solutions of the spectral equations

$$(3.6) \quad H_q^\pm \psi(n) = z\psi(n), \quad z \in \mathbb{C},$$

that decay for $z \in \mathbb{C} \setminus \sigma_\pm$ as $n \rightarrow \pm\infty$. They are uniquely defined by the condition $\psi_q^\pm(z, 0) = 1$, $\psi_q^\pm(z, \cdot) \in \ell^2(\mathbb{Z}_\pm)$. The solution $\psi_q^+(z, n)$ (resp. $\psi_q^-(z, n)$) coincides with the upper (resp. lower) branch of the Baker–Akhiezer functions of H_q^+ (resp. H_q^-), see [27].

The two solutions $\psi_\pm(z, n)$ of the spectral equation

$$(3.7) \quad H\psi = z\psi, \quad z \in \mathbb{C},$$

which are asymptotically close to the Floquet solutions $\psi_q^\pm(z, n)$ of the background equations (3.6) as $n \rightarrow \pm\infty$, are called Jost solutions.

Next, we introduce the sets

$$(3.8) \quad \sigma^{(2)} = \sigma_+ \cap \sigma_-, \quad \sigma_\pm^{(1)} = \overline{\sigma_\pm \setminus \sigma^{(2)}}, \quad \sigma = \sigma_+ \cup \sigma_-,$$

where σ is the (absolutely) continuous spectrum of H and $\sigma_+^{(1)} \cup \sigma_-^{(1)}$, $\sigma^{(2)}$ are the parts which are of multiplicity one, two, respectively.

In addition to the continuous part, H has a finite number of eigenvalues situated in the gaps, $\sigma_d = \{\lambda_1, \dots, \lambda_s\} \subset \mathbb{R} \setminus \sigma$ (see, e.g., [25]). For every eigenvalue we introduce the corresponding norming constants

$$(3.9) \quad \gamma_{\pm, k}^{-1} = \sum_{n \in \mathbb{Z}} |\psi_\pm(\lambda_k, n)|^2, \quad 1 \leq k \leq s.$$

¹Everywhere in this paper the sub or super index “+” (resp. “-”) refers to the background on the right (resp. left) half-axis.

Note that this definition has to be slightly modified in the unlikely event that $\psi_q^\pm(z, n)$ and hence $\psi_\pm(z, n)$ has a pole at $z = \lambda_k$ (see [10] for details). The transmission and reflection coefficients are defined as usual via the scattering relations

$$(3.10) \quad T_\mp(\lambda)\psi_\pm(\lambda, n) = \overline{\psi_\mp(\lambda, n)} + R_\mp(\lambda)\psi_\mp(\lambda, n), \quad \lambda \in \sigma_\mp.$$

Here the values of $\psi_\pm(\lambda, n)$ for $\lambda \in \sigma_\pm$ are to be understood as limits from above $\psi_\pm(\lambda, n) = \lim_{\varepsilon \downarrow 0} \psi_\pm(\lambda + i\varepsilon, n)$ (the corresponding limits from below just give the complex conjugate values $\overline{\psi_\pm(\lambda, n)} = \lim_{\varepsilon \downarrow 0} \psi_\pm(\lambda - i\varepsilon, n)$).

The following result is an immediate consequence of [10, Lem. 5.1].

Theorem 3.1 ([10]). *Suppose $a(n), b(n)$ satisfy (3.4), then $a(n), b(n)$ are uniquely determined by one of the sets of its “partial” scattering data \mathcal{S}_+ or \mathcal{S}_- , where*

$$(3.11) \quad \mathcal{S}_\pm = \left\{ R_\pm(\lambda), \lambda \in \sigma_\pm; |T_\pm(\lambda)|^2, \lambda \in \sigma_\mp^{(1)}; \right. \\ \left. \lambda_1, \dots, \lambda_s \in \mathbb{R} \setminus (\sigma_+ \cup \sigma_-), \gamma_{\pm,1}, \dots, \gamma_{\pm,s} \in \mathbb{R}_+ \right\}.$$

This leads to the natural question if there is a simple way to compute \mathcal{S}_+ from \mathcal{S}_- and vice versa (i.e., without solving the inverse scattering problem). It turns out that this question reduces to the reconstruction of the transmission coefficient $T_\pm(z)$ from \mathcal{S}_\pm . In fact, this follows from the following lemma.

Lemma 3.2 ([10]). *The transmission coefficients $T_\pm(z)$ admit a meromorphic extension to $\mathbb{C} \setminus \sigma$. In general they have simple poles at the eigenvalues λ_k of H . In addition, there are simple poles at $\mu_j^\pm \in \mathbb{R} \setminus \sigma_\pm$ which are not poles of $\psi_q^\pm(z, 1)$ (i.e., $\sigma_j^\pm = \mp 1$) and simple zeros at $\mu_j^\mp \in \mathbb{R} \setminus \sigma_\mp$ which are poles of $\psi_q^\mp(z, 1)$ (i.e., $\sigma_j^\mp = \mp 1$). A pole at μ_j^\pm could cancel with a zero at μ_j^\mp or could give a second order pole if $\mu_j^\pm = \lambda_k$.*

Moreover, the entries of the scattering matrix have the following properties:

$$(a) \quad \rho_+(z)T_+(z) = \rho_-(z)T_-(z), \\ (b) \quad \frac{T_\pm(\lambda)}{\overline{T_\pm(\lambda)}} = R_\pm(\lambda), \quad \lambda \in \sigma_\pm^{(1)}, \\ (c) \quad 1 - |R_\pm(\lambda)|^2 = \frac{\rho_\pm(\lambda)}{\rho_\mp(\lambda)} |T_\pm(\lambda)|^2, \quad \lambda \in \sigma^{(2)}, \\ (d) \quad \overline{R_\pm(\lambda)}T_\pm(\lambda) + R_\mp(\lambda)\overline{T_\pm(\lambda)} = 0, \quad \lambda \in \sigma^{(2)},$$

where

$$(3.12) \quad \rho_\pm(z) = \frac{\prod_{j=1}^{g_\pm} (z - \mu_j^\pm)}{P_\pm^{1/2}(z)}.$$

Hence, the problem is to reconstruct the meromorphic function $T_+(z)$, $z \in \mathbb{C} \setminus \sigma$ from its boundary values

$$(3.13) \quad \begin{cases} |T_+(\lambda)|^2, & \lambda \in \sigma_-^{(1)}, \\ |T_+(\lambda)|^2 = \frac{\rho_-(\lambda)}{\rho_+(\lambda)} (1 - |R_+(\lambda)|^2), & \lambda \in \sigma^{(2)}, \\ \frac{T_+(\lambda)}{\overline{T_+(\lambda)}} = R_+(\lambda), & \lambda \in \sigma_+^{(1)}. \end{cases}$$

That is, we know its absolute value on σ_- and its argument on the rest $\sigma_+^{(1)}$. There will be three Riemann surfaces involved, the one corresponding to $\sigma = \sigma_+ \cup \sigma_-$ and the ones corresponding to σ_\pm . All objects corresponding to σ will be denoted

as in Section 2, while the objects associated with σ_{\pm} will have an additional \pm sub/superscript.

For the reconstruction we will need the solution for the following scalar Riemann–Hilbert problem on the Riemann surfaces associated with σ :

$$(3.14) \quad \begin{aligned} Q_+(z) &= -Q_-(z), \quad z \in \sigma_+^{(1)}, \\ (Q) &\geq -\mathcal{D}_{\underline{\nu}}, \\ Q(\infty_+) &= 1. \end{aligned}$$

Here $\mathcal{D}_{\underline{\nu}}$ is some nonspecial divisor of degree g . We will choose the points $\underline{\nu} = (\nu_1, \dots, \nu_g)$ to be one the lower sheet such that $\nu_j \in \mathbb{R} \setminus \sigma$.

Theorem 3.3. *The solution of (3.14) is given by*

$$(3.15) \quad Q(z) = \frac{\theta(\underline{z}(\infty_+))}{\theta(\underline{z}(\infty_+) - \underline{\delta})} \frac{\theta(\underline{z}(z) - \underline{\delta})}{\theta(\underline{z}(z))} \exp\left(\frac{1}{2} \int_{\sigma_+^{(1)}} \omega_{z\infty_+}\right),$$

where

$$(3.16) \quad \underline{z}(z) = \underline{A}_{E_0}(z) - \underline{\alpha}_{E_0}(\mathcal{D}_{\underline{\nu}}) - \underline{\Xi}_{E_0}, \quad \delta_{\ell} = \frac{1}{2} \int_{\sigma_+^{(1)}} \zeta_{\ell},$$

and the integrals are taken over the lift of the indicated spectra to the upper sheet Π_u . Here θ , \underline{A}_{E_0} , $\underline{\alpha}_{E_0}$, and $\underline{\Xi}_{E_0}$ are the Riemann theta function, Abel map, Abel map for divisors, and vector of Riemann constants for the Riemann surface associated with σ , respectively.

In addition, we have $Q(z) = \overline{Q(\bar{z})}$ and $Q(z)$ has square root type zeros on the upper sheet and square root type poles on the lower sheet near each open endpoint. We have $Q(z) \in \mathbb{R}$ for z in $\mathbb{R} \setminus \sigma_+^{(1)}$ and $Q(z) \in i\mathbb{R}$ for z in $\sigma_+^{(1)}$.

Proof. The first part follows as in [15, Thm. 4.3] and the behaviour near open endpoints as in [15, Lem. 4.4]. The last claim follows from $Q(z) = \overline{Q(\bar{z})}$ since $Q_+(z) = \overline{Q_+(z^*)} = \overline{Q_-(z)} = \pm Q_+(z)$ with the plus sign for $z \in \sigma \setminus \sigma_+^{(1)}$ and the minus sign for $z \in \sigma_+^{(1)}$. \square

Now we are ready to show

Theorem 3.4. *The transmission coefficient $T_+(z)$ can be reconstructed from the reflection coefficient $R_+(z)$ and the eigenvalues λ_j via*

$$(3.17) \quad \begin{aligned} T_+(z) &= \left(\prod_{\mu_j^- \in M^-} B_-(z, \mu_j^-) \right) \left(\prod_{\mu_j^+ \in M^+} B_-(z, \mu_j^+)^{-1} \right) \left(\prod_{k=1}^s B_-(z, \lambda_k)^{-1} \right) \times \\ &\exp\left(\frac{Q(z)^{-1}}{\pi i} \int_{\sigma_-^{(1)}} Q \log(|T_+|) \omega_{zz^*}\right) \\ &+ \frac{Q(z)^{-1}}{2\pi i} \int_{\sigma^{(2)}} Q \left(\log\left(\frac{\rho_-}{\rho_+}\right) + \log(1 - |R_+|^2) \right) \omega_{zz^*} \\ &+ \frac{Q(z)^{-1}}{2\pi} \int_{\sigma_+^{(1)}} Q \left(\arg(R_+) + \delta^- \right) \omega_{zz^*}, \end{aligned} \quad (3.18)$$

where the integrals are taken over the lift of the indicated spectra to the upper sheet Π_u (of the Riemann surface associated with σ). Moreover, we use the convention

that we identify z with $(z, +)$, and similarly for λ_k, μ_j^\pm , whenever used in the argument of a function defined on a Riemann surface. Here

$$(3.19) \quad M^\pm = \{\mu_j^\pm \mid \mu_j^\pm \in \mathbb{R} \setminus \sigma \text{ and } \sigma_j^\pm = -1\},$$

and

$$(3.20) \quad \delta^-(\lambda) = \sum_{\ell} \delta_{\ell}^- \chi_{[E_{2\ell-1}^-, E_{2\ell}^-]}(\lambda)$$

with (cf. Lemma 2.2)

$$(3.21) \quad \delta_{\ell}^- = - \sum_{\mu_j^- \in M^-} \delta_{\ell}^-(\mu_j^-) + \sum_{\mu_j^+ \in M^+} \delta_{\ell}^-(\mu_j^+) + \sum_{k=1}^s \delta_{\ell}^-(\lambda_k).$$

Proof. We start by considering the multivalued function

$$(3.22) \quad t_+(z) = \left(\prod_{\mu_j^- \in M^-} B_-(z, \mu_j^-)^{-1} \right) \left(\prod_{\mu_j^+ \in M^+} B_-(z, \mu_j^+) \right) \left(\prod_{k=1}^s B_-(z, \lambda_k) \right) T_+(z)$$

which has neither zeros nor poles on Π_u and satisfies

$$(3.23) \quad \begin{cases} |t_+(\lambda)|^2 = |T_+(\lambda)|^2, & \lambda \in \sigma_-, \\ \arg(t_+(\lambda)) = \arg(T_+(\lambda)) + \delta_{\ell}^-, & \lambda \in \sigma_+^{(1)} \cap [E_{2\ell-1}^-, E_{2\ell}^-]. \end{cases}$$

Moreover, the absolute value of $t_+(z)$ is single-valued and hence its logarithm is a harmonic function on Π_u which can be reconstructed from its boundary values. To accommodate the fact that we know its absolute value on σ_- and its argument on $\sigma_+^{(1)}$ we consider

$$(3.24) \quad Q(z) \log(t_+(z)).$$

Note that since $t_+(z)$ might still have zeros and poles on σ , the function $\log(t_+(z))$ might have logarithmic singularities on σ .

Since $Q(\lambda)$ is real-valued for $\lambda \in \mathbb{R} \setminus \sigma_+^{(1)}$ and purely imaginary for $\lambda \in \sigma_+^{(1)}$, we infer that the real part of $Q(z) \log(t_+(z))$ is harmonic on Π_u and can be reconstructed from its boundary values

$$(3.25) \quad \operatorname{Re}(Q(\lambda) \log(t_+(\lambda))) = \begin{cases} Q(\lambda) \log(|T_+(\lambda)|), & \lambda \in \sigma_-, \\ iQ(\lambda) (\arg(T_+(\lambda)) + \delta_{\ell}^-), & \lambda \in \sigma_+^{(1)} \cap [E_{2\ell-1}^-, E_{2\ell}^-], \end{cases}$$

using Green's function:

$$(3.26) \quad \operatorname{Re}(Q(z) \log(t_+(z))) = \operatorname{Re} \left(\frac{1}{\pi i} \int_{\sigma_-} Q \log(|T_+|) \omega_{zz^*} + \frac{1}{\pi} \int_{\sigma_+^{(1)}} Q (\arg(T_+) + \delta^-) \omega_{zz^*} \right).$$

Dropping the real part we get

$$(3.27) \quad T_+(z) = \left(\prod_{\mu_j^- \in M^-} B_-(z, \mu_j^-) \right) \left(\prod_{\mu_j^+ \in M^+} B_-(z, \mu_j^+)^{-1} \right) \left(\prod_{k=1}^s B_-(z, \lambda_k)^{-1} \right) \times \\ \exp \left(\frac{Q(z)^{-1}}{\pi i} \int_{\sigma_-} Q \log(|T_+|) \omega_{zz^*} + \frac{Q(z)^{-1}}{\pi} \int_{\sigma_+^{(1)}} Q(\arg(T_+) + \delta^-) \omega_{zz^*} \right).$$

In fact, by [29, Thm. 1] both the left-hand and the right-hand side have the same absolute value and hence can only differ by a constant with absolute value one (in particular, the right-hand side is single-valued since the left-hand side is). This constant must be one since both sides are real-valued for real-valued z to the left of σ . \square

Acknowledgments. We want to thank Peter Yuditskii for helpful discussions. We thank Dmitry Shepelsky for pointing out an error in the original version of this paper.

REFERENCES

- [1] N. I. Akhiezer, *Elements of the Theory of Elliptic Functions*, Amer. Math. Soc., Providence, 1990.
- [2] A. Boutet de Monvel, I. Egorova, and G. Teschl, *Inverse scattering theory for one-dimensional Schrödinger operators with steplike finite-gap potentials*, J. Analyse Math. **106:1**, 271–316 (2008).
- [3] W. Bulla, F. Gesztesy, H. Holden, and G. Teschl *Algebro-Geometric Quasi-Periodic Finite-Gap Solutions of the Toda and Kac-van Moerbeke Hierarchies*, Mem. Amer. Math. Soc. **135-641**, (1998).
- [4] V.S. Buslaev and V.N. Fomin, *An inverse scattering problem for the one-dimensional Schrödinger equation on the entire axis*, Vestnik Leningrad. Univ. **17**, no. 1, 56–64 (1962).
- [5] P. Deift and X. Zhou, *A steepest descent method for oscillatory Riemann–Hilbert problems*, Ann. of Math. (2) **137**, 295–368 (1993).
- [6] I. Egorova, J. Michor, and G. Teschl, *Scattering theory for Jacobi operators with quasi-periodic background*, Comm. Math. Phys. **264-3**, 811–842 (2006).
- [7] I. Egorova, J. Michor, and G. Teschl, *Inverse scattering transform for the Toda hierarchy with quasi-periodic background*, Proc. Amer. Math. Soc. **135**, 1817–1827 (2007).
- [8] I. Egorova, J. Michor, and G. Teschl, *Scattering theory for Jacobi operators with steplike quasi-periodic background*, Inverse Problems **23**, 905–918 (2007).
- [9] I. Egorova, J. Michor, and G. Teschl, *Soliton solutions of the Toda hierarchy on quasi-periodic background revisited*, Math. Nach. **282:4**, 526–539 (2009).
- [10] I. Egorova, J. Michor, and G. Teschl, *Scattering theory for Jacobi operators with general steplike quasi-periodic background*, Zh. Mat. Fiz. Anal. Geom. **4-1**, 33–62 (2008).
- [11] H. Farkas and I. Kra, *Riemann Surfaces*, 2nd edition, GTM 71, Springer, New York, 1992.
- [12] F. Gesztesy, H. Holden, J. Michor, and G. Teschl, *Soliton Equations and their Algebro-Geometric Solutions. Volume II: (1+1)-Dimensional Discrete Models*, Cambridge Studies in Advanced Mathematics **114**, Cambridge University Press, Cambridge, 2008.
- [13] S. Kamvissis, *On the long time behavior of the doubly infinite Toda lattice under initial data decaying at infinity*, Comm. Math. Phys., **153-3**, 479–519 (1993).
- [14] S. Kamvissis and G. Teschl, *Stability of periodic soliton equations under short range perturbations*, Phys. Lett. A, **364-6**, 480–483 (2007).
- [15] S. Kamvissis and G. Teschl, *Stability of the periodic Toda lattice under short range perturbations*, arXiv:0705.0346.

- [16] S. Kamvissis and G. Teschl, *Stability of the periodic Toda lattice: Higher order asymptotics*, arXiv:0805.3847.
- [17] R. Killip and B. Simon, *Sum rules for Jacobi matrices and their applications to spectral theory*, Ann. of Math. (2) **158**, 253–321 (2003).
- [18] H. Krüger and G. Teschl, *Long-time asymptotics for the Toda lattice in the soliton region*, Math. Z. **262**, 585–602 (2009).
- [19] H. Krüger and G. Teschl, *Long-time asymptotics of the Toda lattice for decaying initial data revisited*, Rev. Math. Phys. **21:1**, 61–109 (2009).
- [20] H. Krüger and G. Teschl, *Long-time asymptotics for the periodic Toda lattice in the soliton region*, Int. Math. Res. Not. (to appear).
- [21] A. Laptev, S. Naboko, and O. Safronov, *On new relations between spectral properties of Jacobi matrices and their coefficients*, Comm. Math. Phys. **241**, no. 1, 91–110 (2003).
- [22] J. Michor and G. Teschl, *Trace formulas for Jacobi operators in connection with scattering theory for quasi-periodic background*, in Operator Theory, Analysis and Mathematical Physics, J. Janas, et al. (eds.), 51–57, Oper. Theory Adv. Appl., 174 Birkhäuser, Basel, 2007.
- [23] F. Nazarov, F. Peherstorfer, A. Volberg, and P. Yuditskii, *On generalized sum rules for Jacobi matrices*, Int. Math. Res. Not. **2005** no. 3, 155–186 (2005).
- [24] B. Simon and A. Zlatoš, *Sum rules and the Szegő condition for orthogonal polynomials on the real line*, Comm. Math. Phys. **242**, no. 3, 393–423 (2003).
- [25] G. Teschl, *Oscillation theory and renormalized oscillation theory for Jacobi operators*, J. Diff. Eqs. **129**, 532–558 (1996).
- [26] G. Teschl, *Inverse scattering transform for the Toda hierarchy*, Math. Nachr. **202**, 163–171 (1999).
- [27] G. Teschl, *Jacobi Operators and Completely Integrable Nonlinear Lattices*, Math. Surv. and Mon. **72**, Amer. Math. Soc., Providence, R.I. 2000.
- [28] G. Teschl, *Algebraic constraints on solitons with respect to quasi-periodic backgrounds*, Bull. London Math. Soc. **39-4**, 677–684 (2007).
- [29] V. Voichick and L. Zalcman, *Inner and outer functions on Riemann surfaces*, Proc. Amer. Math. Soc. **16**, 1200–1204 (1965).
- [30] A. Volberg and P. Yuditskii, *On the inverse scattering problem for Jacobi Matrices with the Spectrum on an Interval, a finite system of intervals or a Cantor set of positive length*, Comm. Math. Phys. **226**, 567–605 (2002).
- [31] A. Zlatoš, *Sum rules for Jacobi matrices and divergent Lieb-Thirring sums*, J. Funct. Anal. **225**, no. 2, 371–382 (2005).

B.VERKIN INSTITUTE FOR LOW TEMPERATURE PHYSICS, 47 LENIN AVENUE, 61103 KHARKIV, UKRAINE

E-mail address: egorova@ilt.kharkov.ua

FACULTY OF MATHEMATICS, UNIVERSITY OF VIENNA, NORDBERGSTRASSE 15, 1090 WIEN, AUSTRIA, AND INTERNATIONAL ERWIN SCHRÖDINGER INSTITUTE FOR MATHEMATICAL PHYSICS, BOLTZMANNGASSE 9, 1090 WIEN, AUSTRIA

E-mail address: Gerald.Teschl@univie.ac.at

URL: <http://www.mat.univie.ac.at/~gerald/>