



Hook length property of d-complete posets via q-integrals

Sungkyunkwan University


Jang Soo Kim

(Joint work with Meesue Yoo)

Hook length formula for shapes


- $\lambda = (\lambda_1, \dots, \lambda_L)$ is a **partition** of n
if $\lambda_1 \geq \dots \geq \lambda_L > 0$ and $\lambda_1 + \dots + \lambda_L = n$
- Young diagram of $\lambda = (4, 3, 1)$ is 
- A standard Young tableau of shape $\lambda = (4, 3, 1)$
is .
- hook length of $x \in \lambda$:

Thm $f^\lambda = \frac{n!}{\prod_{x \in \lambda} h(x)}$



Hook length formula for shifted shapes

- $\lambda = (\lambda_1, \dots, \lambda_L)$ is a strict partition of n
if $\lambda_1 > \dots > \lambda_L > 0$ and $\lambda_1 + \dots + \lambda_L = n$
- shifted Young diagram of $\lambda = (8, 7, 5, 2)$



$$h(x) = 13$$

1	2	3	6	7	12	15	16
4	5	8	13	14	20	21	
9	10	17	19	22			
11	18						


SYT

Thm λ is a strict partition of n

$$\Rightarrow q^\lambda = \frac{n!}{\prod_{x \in \lambda} h(x)}$$

Hook length formula for trees

(3)












P: tree poset.

The **hook length** of $x \in P$
is $\#\{y \in P : y \leq x\}$

Thm # linear extensions of P is



$$\frac{n!}{\prod_{x \in P} h(x)}.$$

ex






$$\frac{5!}{1 \cdot 1 \cdot 1 \cdot 3 \cdot 5} = 8$$

- reverse SYTs are linear extensions.



poset rotated 45°



Thm P : poset (shape, shifted shape or tree)

⇒ # linear extensions of P

$$= \frac{n!}{\prod_{x \in P} h(x)}$$

Proctor generalized this to d -complete posets.


P-partitions

Def P : poset.

A **P-partition** is an order-reversing map.



($\sigma: P \rightarrow \mathbb{N}$ s.t. $\sigma(x) \geq \sigma(y)$ if $x \leq_P y$)

ex




$\begin{matrix} 1 \\ 3 \\ 3 \\ 7 \end{matrix}$

(partition)



(reverse plane partition)



\Leftrightarrow

0	0	0	1	3	4
1	1	2	3		
2					

(shifted RPP)

- The **size** of $\sigma: P \rightarrow \mathbb{N}$ is $|\sigma| = \sum_{x \in P} \sigma(x)$.

- $GF_q(P) = \sum_{\sigma: P \rightarrow \mathbb{N}} q^{|\sigma|}$

q-Hook length formula

Thm P : shape, shifted shape or tree

$$GF_q(P) = \sum_{\sigma: P \rightarrow N} q^{|\sigma|} = \prod_{x \in P} \frac{1}{1 - q^{h(x)}}$$

FACT: $\sum_{\sigma: P \rightarrow N} q^{|\sigma|} = \frac{\sum_{\pi \in \mathcal{L}(P)} q^{\text{maj}(\pi)}}{(1-q)(1-q^2)\cdots(1-q^n)}$

Thm (Proctor)

P : d-complete poset.

$$GF_q(P) = \sum_{\sigma: P \rightarrow N} q^{|\sigma|} = \prod_{x \in P} \frac{1}{1 - q^{h(x)}}$$

Goal: Prove this theorem.

Note: This theorem has been proved Proctor (with Peterson), Ishikawa and Tagawa, and Nakada.

Proctor proved more general colored-hook length formula.

Okada conjectured even more general hook length formula with Macdonal weights.




Outline of Proof

- (Proctor) Every d-complete poset is a slant-sum of irreducible d-complete posets.
 - (Proctor) There are 15 classes of irr. d.c.p.
 - (Kim, Stanton) $GF_q(P)$ can be written as q-integrals.
1. Show that irr. d.c.p are enough to consider.
 2. Express $GF_q(P)$ for each irr. d.c.p P as q-integral.
 3. Evaluate the q-integral.

(Among 15 integrals, 2 of them are known,
11 of them can be evaluated by computer.)

d-complete poset

Def P is **d-complete** if




$\begin{bmatrix} y \\ x \end{bmatrix}$ means $[x,y]$




Def P : d-complete.

The **hook length** of $x \in P$ is


$$h(x) = \begin{cases} h(y) + h(z) - h(w) & \text{if } \\ & \# y \leq x \\ & \# z \leq x \\ & \# w \leq x \end{cases} \quad \text{otherwise}$$




ex).





Irreducible d-complete posets









shapes



shifted shapes.






Slant sums

Def P : d-complete.

$x \in P$: acyclic if

① x is covered by at most one element,


②



(Every irr. d.c.p has
at most 2 acyclic elements.)

Def P_1, P_2 : d-complete. $x \in P_1, y \in P_2$, x : acyclic
 y : max of P_2

Slant sum $P_1^x \setminus_y P_2$ is





Thm (Proctor)

Every d-complete poset is a slant sum
of irr. d-complete posets.

Semi-irreducible d-complete posets

Def A **semi-irreducible** d-complete poset is a poset obtained from an irr. d.c.p. by adding a chain below each acyclic element.

ex).




irr.

semi-irr.

Lem P, Q : d-complete. $x \in P$: acyclic, $y \in Q$: max.

$$GF_q(P^x \setminus y Q) = GF_q(P') GF_q(Q) \cdot \prod_{i=1}^{|Q|} (1 - q^i)$$


Cor It is enough to consider semi-irr. d.c.p.



q-integrals

- $\int_0^a f(x) dx = \text{area of}$

$$= \lim_{n \rightarrow \infty} \sum_{i=0}^n f(x_i) (x_i - x_{i+1})$$



$$x_i = aq^i \quad (0 < q < 1)$$

Def **q-integral**

$$\int_0^a f(x) d_q x = \sum_{i=0}^{\infty} f(aq^i) (aq^i - aq^{i+1})$$

$$\int_a^b f(x) d_q x = \int_0^b f(x) d_q x - \int_0^a f(x) d_q x$$

Note

$$\lim_{q \rightarrow 1^-} \int_a^b f(x) d_q x = \int_a^b f(x) dx.$$

FACT


$$\int_a^b x^n d_q x = \frac{b^{n+1} - a^{n+1}}{1+q+\dots+q^{n+1}}$$

Order polytopes

Def P : poset on $\{x_1, \dots, x_n\}$. (naturally labeled)

The **order polytope** of P is

$$O(P) = \{(x_1, \dots, x_n) \in [0, 1]^n : x_i \leq x_j \text{ if } x_i \leq_P x_j\}$$



$$O(P) = \{(x_1, x_2, x_3, x_4) \in [0, 1]^4 \mid x_1 \leq x_2, x_2 \leq x_3, x_2 \leq x_4\}$$

- We can consider q-integrals over $O(P)$.

$$\begin{aligned} & \int_{O(P)} f(x_1, x_2, x_3, x_4) d_q x_1 d_q x_2 d_q x_3 d_q x_4 \\ &= \int_0^1 \int_0^1 \int_0^{\min(x_3, x_4)} \int_0^{x_2} f(x_1, x_2, x_3, x_4) d_q x_1 d_q x_2 d_q x_3 d_q x_4 \end{aligned}$$

q -integral over order polytope

Thm (Kim, Stanton, 2016)

P : poset on $\{x_1, \dots, x_n\}$ (naturally labeled)

$$\int_{O(P)} f(x_1, \dots, x_n) d_q x_1 \dots d_q x_n \\ = (1-q)^n \sum_{\sigma: P \rightarrow N} f(q^{\sigma(1)}, \dots, q^{\sigma(n)}) q^{|\sigma|}$$

In particular,

$$\int_{O(P)} d_q x_1 \dots d_q x_n = (1-q)^n GF_q(P)$$

Note: If P is not naturally labeled,

we have (P, ω) -partitions. (The label ω is related to order of integration.)

Attaching shifted shapes

Lem (Kim, Stanton, 2016)


P : poset on $\{u_1, \dots, u_n\}$

Q : poset on $\{v_1, \dots, v_N\}$

$y_1 < \dots < y_k$: chain in P

(with certain conditions)

P



$$\delta_k = (k-1, \dots, 1, 0)$$

$$\int_{O(Q)} f(x_1, \dots, x_n) dq u_1 \cdots dq u_N$$




$$= \frac{q^c}{(1-q)^k} \int_{O(P)} f(x_1, \dots, x_n) \frac{\alpha_{\lambda+\delta_k}(y_1, \dots, y_k)}{\prod_{i=1}^k (q;q)_{\lambda+k-i}} dq u_1 \cdots dq u_n$$

$$\alpha_\lambda(x_1, \dots, x_n) = \det \left(x_i^{j-1} \right)_{i,j=1}^n, \quad (q;q)_n = (1-q)(1-q^2) \cdots (1-q^n)$$

Expressing $GF_q(P)$ as q -integral.

(17)

Observation: Every semi-irreducible d.e.p is obtained as follows.



Attach $\lambda + \delta_3$ and a chain above the top.

$$GF_q(P) = \int_{O(P)} dq y_1 \cdots dq y_N \sim \int_{0 \leq x_1, \dots \leq x_4 \leq 1} a_{\mu + \delta_3}(x_2, x_3, x_4) a_{\delta_2}(x_1, x_2) a_{\delta_2}(x_3, x_4) \\ \cdot a_{\lambda + \delta_3}(x_1, x_2, x_3) dq x_1 \cdots dq x_n.$$

Evaluation of q -integral

FACT

$$\int_{0 \leq x_1 \leq \dots \leq x_n \leq 1} x_1^{a_1} \dots x_n^{a_n} d_q x_1 \dots d_q x_n = \prod_{i=1}^n \frac{1-q}{1-q^{a_i+1+i}}$$

Thus,

$$\int_{0 \leq x_1 \leq \dots \leq x_4 \leq 1} a_{\mu+\delta_3}(x_2, x_3, x_4) a_{\delta_2}(x_1, x_2) a_{\delta_2}(x_3, x_4) a_{\lambda+\delta_3}(x_1, x_2, x_3) d_q x_1 \dots d_q x_4$$

can be evaluated by computer!

There are 19 classes of semi-itr. d.c.p.


\Rightarrow " 19 q -integrals to evaluate.

15 of them can be done by computer. (Took 11 hrs.)

2 of the remaining are known.

\Rightarrow We need to evaluate 2 q -integrals.

* Semi-irr. d.c.p. of class 4.





$$\hat{\mu}_i^{(l)} = \begin{cases} \mu_i + 1 & \text{if } i < l \\ \mu_{i+1} & \text{if } i \geq l \end{cases}$$

$$\begin{aligned}
 GF_q(P_4) &\sim \int_{0 \leq x_1 \leq \dots \leq x_{n+1} \leq 1} x_n^k a_{\lambda + \delta_{n-1}}(x_1, \dots, x_{n-1}) \underbrace{a_{\mu + \delta_{n+1}}(x_1, \dots, x_{n+1})}_{= \sum_{l=1}^{n+1} (-1)^{n+l-l} x_{n+1}^{\mu_l + n+1 - l}} dq x_1 \cdots dq x_{n+1} \\
 &= \sum_{l=1}^{n+1} \int_0^1 (-1)^{n+l-l} x_{n+1}^{\mu_l + n+1 - l} a_{\hat{\mu}_i^{(l)} + \delta_n}(x_1, \dots, x_n) dq x_1 \cdots dq x_n dq x_{n+1} \\
 &= \sum_{l=1}^{n+1} \int_0^1 (-1)^{n+l-l} x_{n+1}^{\mu_l + n+1 - l} \int_{0 \leq x_1 \leq \dots \leq x_n \leq x_{n+1}} x_n^k a_{\lambda + \delta_{n-1}}(x_1, \dots, x_{n-1}) a_{\hat{\mu}_i^{(l)} + \delta_n}(x_1, \dots, x_n) dq x_1 \cdots dq x_n dq x_{n+1} \\
 &= \sum_{l=1}^{n+1} \int_0^1 (-1)^{n+l-l} x_{n+1}^{\mu_l + n+1 - l + k + |\lambda| + |\hat{\mu}_i^{(l)}| + \binom{n-1}{2} + \binom{n}{2}} dq x_{n+1} \cdot Y_l.
 \end{aligned}$$

$$GF_q(P_4) \sim \sum_{\ell=1}^{m+1} A_\ell Y_\ell, \quad (A_\ell: \text{easy constant}) \quad 20$$

$$Y_\ell = \int_{0 \leq x_1 \leq \dots \leq x_n \leq x_{n+1}} x_n^k a_{\lambda + \delta_{n-1}}(x_1, \dots, x_{n-1}) a_{\hat{\mu}^{(\ell)} + \delta_n}(x_1, \dots, x_n) d_q x_1 \dots d_q x_n$$

$$Y_\ell \sim GF_q(Q_\ell) = \frac{1}{1-q^N} GF_q(Q_\ell^-) = \frac{1}{1-q^N} \cdot \frac{1}{(q;q)_k} \cdot \prod_{x \in V^{(\ell)}} \frac{1}{1-q^{h(x)}}$$



The hook length formula is equivalent to the following identity.

$$\frac{\prod_{j=1}^{n-1} (1 - q^{|\lambda|+|\mu|+\lambda_j+n^2+n-j+k+1})}{\prod_{i=1}^{n+1} (1 - q^{|\lambda|+|\mu|-\mu_i+n^2-n+k+i})} = \sum_{\ell=1}^{n+1} \frac{q^{-|\lambda|-|\mu|+\mu_\ell-n^2+n-k-\ell}}{1 - q^{|\lambda|+|\mu|-\mu_\ell+n^2-n+k+\ell}} \cdot \frac{\prod_{j=1}^{n-1} (1 - q^{\mu_\ell+\lambda_j+2n-\ell-j+1})}{\prod_{j=1, j \neq \ell}^{n+1} (1 - q^{\mu_\ell-\mu_j+j-\ell})}.$$

This can be obtained by the known partial fraction expansion.

$$\frac{\prod_{j=1}^{n+1} (1 - b_j/t)}{\prod_{j=1}^n (1 - a_j/t)} = \sum_{\ell=1}^n \frac{\prod_{j=1}^{n+1} (1 - a_\ell/b_j)}{(1 - a_\ell/t) \prod_{j=1, j \neq \ell}^n (1 - a_\ell/a_j)}, \quad \text{for } b_1 \cdots b_{n+1} = a_1 \cdots a_n t.$$

The hook length formula for the remaining semi-irreducible d-complete poset can be proved similarly. In this case we need the following partial fraction expansion.

$$\prod_{i=1}^n \frac{1 - tx_i y_i}{1 - tx_i} = y_1 \cdots y_n + \sum_{\ell=1}^n \frac{1 - y_\ell}{1 - tx_\ell} \prod_{i=1, i \neq \ell}^n \frac{1 - x_i y_i / x_\ell}{1 - x_i / x_\ell}.$$

T

H

A

N

K

Y

C

L