

## Tensor fields and connections on holomorphic orbit spaces of finite groups

Andreas Kriegl, Mark Losik, and Peter W. Michor \*

**Abstract.** For a representation of a finite group  $G$  on a complex vector space  $V$  we determine when a holomorphic  $\binom{p}{q}$ -tensor field on the principal stratum of the orbit space  $V/G$  can be lifted to a holomorphic  $G$ -invariant tensor field on  $V$ . This extends also to connections. As a consequence we determine those holomorphic diffeomorphisms on  $V/G$  which can be lifted to orbit preserving holomorphic diffeomorphisms on  $V$ . This in turn is applied to characterize complex orbifolds.

**Keywords:** complex orbifolds, orbit spaces of complex finite group actions

**Subject Classification:** 32M17

### 1. Introduction

Locally, an orbifold  $Z$  can be identified with the orbit space  $B/G$ , where  $B$  is a  $G$ -invariant neighborhood of the origin in a vector space  $V$  with a finite group  $G \subset GL(V)$  and, using this identification, one can easily define local (and then global) tensor fields and other differential geometrical objects in  $Z$  as appropriate  $G$ -invariant tensor fields and objects on  $B \subset V$ . In particular, one can naturally define Riemannian orbifolds, Einstein orbifolds, symplectic orbifolds, Kähler-Einstein orbifolds etc.

We study complex orbifolds, that is, orbifolds modeled on orbit spaces  $V/G$ , where  $G$  is a finite subgroup of  $GL(V)$  for a complex vector space  $V$ . In particular, the orbit spaces  $Z = M/G$  of a discrete proper group  $G$  of holomorphic transformations of a complex manifold  $M$  are complex orbifolds.

An orbifold  $X$  has a structure defined by the sheaf  $\mathfrak{F}_X$  of local invariant holomorphic functions in a local uniformizing system.  $X$  has also a stratification by strata  $S$  which are glued from local isotropy type strata of local uniformizing systems. In particular, the regular stratum  $X_0$  is an open dense complex manifold in  $X$ .

Holomorphic geometric objects on  $X$  (e.g. tensor fields and connections) are locally defined as invariant objects on the uniformizing system. Their restrictions to the regular stratum  $X_0$  are usual holomorphic geometric objects on the complex manifold  $X_0$ .

A natural question is to characterize these restrictions, i.e. to describe tensor fields and connections on  $X_0$  which are extendible to  $X$ . We look at the lifting problem for connections because this allows a very elegant approach to the lifting

---

\*M.L. and P.W.M. were supported by ‘Fonds zur Förderung der wissenschaftlichen Forschung, Projekt P 14195 MAT’

problem for holomorphic diffeomorphisms. And the last problem has immediate consequences for characterizing complex orbifolds, i.e., for answering the following question: Which data does one need besides  $\mathfrak{F}_X$  and  $X_0$  to characterize a complex orbifold  $X$ ? The main goal of the paper is to answer these questions.

We have first to investigate the local situation, thus we consider a finite subgroup  $G \subset GL(V)$  and the orbit space  $Z = V/G$  with the structure given by the sheaf  $\mathfrak{F}_{V/G}$  of invariant holomorphic functions on  $V$ , and the orbit type stratification. The prime role is played by strata of codimension 1 with the orders of the corresponding stabilizer groups, which are arranged in the *reflection divisor*  $D_{V/G}$  which keeps track of all complex reflections in  $G$ . It turns out that the union  $Z_1$  of  $Z_0$  and of all codimension 1 strata is a complex manifold, see 3.5. We characterize all  $G$ -invariant holomorphic tensor fields and connections on  $V$  in terms of the *reflection divisor* of the corresponding meromorphic tensor field and connection on  $Z_1$ , see 3.7 and 4.2. Our result gives a generalization 3.9 of Solomon's theorem [10], see 3.10. Using the lifting property of connections we are able to prove that a holomorphic diffeomorphism  $Z = V/G \rightarrow V/G' = Z'$  between two orbit spaces has a holomorphic lift to  $V$  which is equivariant over an isomorphism  $G \rightarrow G'$  if and only if  $f$  respects the regular strata and the reflection divisors, i.e.  $f(Z_0) \subset Z'_0$  and  $f_*(D_Z) \subset D_{Z'}$ . In fact we give two proofs of this result, which in [4] is carried over to the algebraic geometry setting for algebraically closed ground fields of characteristic 0. The related problem of lifting (smooth) homotopies from (general) orbit spaces has been treated in [1] and [9].

Applying the local results we prove that a complex orbifold  $X$  is uniquely determined by the sheaf  $\mathfrak{F}_X$ , the regular stratum  $X_0$ , and the reflection divisor  $D_X$  alone, see 6.6.

## 2. Preliminaries

**2.1. The orbit type stratification.** Let  $V$  be an  $n$ -dimensional complex vector space,  $G$  a finite subgroup of  $GL(V)$ , and  $\pi : V \rightarrow V/G$  the quotient projection. The ring  $\mathbb{C}[V]^G$  has a minimal system of homogeneous generators  $\sigma^1, \dots, \sigma^m$ . We will use the map  $\sigma = (\sigma^1, \dots, \sigma^m) : V \rightarrow \mathbb{C}^m$ . Denote by  $Z$  the affine algebraic variety in  $\mathbb{C}^m$  defined by the relations between  $\sigma^1, \dots, \sigma^m$ . It is known that  $\sigma(V) = Z$ .

We consider the orbit space  $V/G$  endowed with the quotient topology as a local ringed space defined by the following sheaf of rings  $\mathfrak{F}_{V/G}$ : if  $U$  is an open subset of  $V/G$ ,  $\mathfrak{F}_{V/G}(U)$  is equal to the space of  $G$ -invariant holomorphic functions on  $\pi^{-1}(U)$ . Clearly one may consider sections of  $\mathfrak{F}_{V/G}$  on  $U$  as functions on  $U$ . We call these functions holomorphic functions on  $U$ . It is known that the map of the orbit space  $V/G$  to  $Z$  induced by the map  $\sigma$  is a homeomorphism. Moreover, this homeomorphism induces an isomorphism of the sheaf  $\mathfrak{F}_{V/G}(U)$  and the structure sheaf of the complex algebraic variety  $Z$  (see [7]). Via the above isomorphism we identify the local ringed spaces  $V/G$  and  $Z$ . Under this identification the projection  $\pi$  is identified with the map  $\sigma$ . Let  $G$  and  $G'$  be finite subgroups of  $GL(V)$  and let  $Z = V/G$  and  $Z' = V/G'$  be the corresponding orbit spaces. By definition a holomorphic diffeomorphism of the orbit space  $Z$  to the orbit space  $Z'$  is an isomorphism of  $Z$  to  $Z'$  as local ringed spaces.

Let  $K$  be a subgroup of  $G$ ,  $(K)$  the conjugacy class of  $K$ . Denote by  $V_{(K)}$  the set of points of  $V$  whose isotropy groups belong to  $(K)$  and put  $Z_{(K)} = \pi(V_{(K)})$ . It is known that  $\{Z_{(K)}\}$  is a finite stratification of  $Z$ , called the isotropy type stratification, into locally closed irreducible smooth algebraic subvarieties (see [5]). Denote by  $Z^i$  the union of the strata of codimension greater than  $i$  and put  $Z_i = Z \setminus Z^i$ . Then  $Z_0$  is the principal stratum of  $Z$ , i.e.  $Z_0 = Z_{(K)}$  for  $K = \{\text{id}\}$ . It is known that  $Z_0$  is a Zariski open subset of  $Z$  and a complex manifold. It is clear that the restriction of the map  $\sigma$  to the set  $V_{\text{reg}}$  of regular points of  $V$  is a tale map onto  $Z_0$ .

In this paper we consider the orbit space  $Z = V/G$  with the above structure of local ringed space and the stratification  $\{Z_{(K)}\}$ .

**2.2. The divisor of a tensor field.** We shall use divisors of meromorphic functions on a complex manifold  $X$ . For technical reasons (see e.g. the last formula of this section) we define  $\text{div}(0) = \sum_S \infty \cdot S$ , where the sum runs over all complex subspaces of  $X$  of codimension 1.

Let  $f$  and  $g$  be two meromorphic functions on  $X$ . Then we have  $\text{div}(f + g) \geq \min\{\text{div}(f), \text{div}(g)\}$ , where  $\text{div}(f)$  denote the divisor of  $f$ . Taking the minimum means: For each irreducible complex subspace  $S$  of  $X$  of codimension 1 belonging to the support of  $f$  or  $g$  take the minimum of the coefficients in  $\mathbf{Z}$  of  $S$  in  $\text{div}(f)$  and  $\text{div}(g)$ .

Let  $P$  be a meromorphic tensor field (i.e., with meromorphic coefficient functions in local coordinates) on  $X$ . In local holomorphic coordinates  $y^1, \dots, y^n$  on an open subset  $U \subset X$  the tensor field  $P$  can be written as

$$P|_U = \sum_{i_1, \dots, i_p, j_1, \dots, j_q} P_{j_1 \dots j_q}^{i_1 \dots i_p} \frac{\partial}{\partial y^{i_1}} \otimes \dots \otimes \frac{\partial}{\partial y^{i_p}} \otimes dy^{j_1} \otimes \dots \otimes dy^{j_q}.$$

and we define the *divisor* of  $P$  on  $U$  as the minimum of all divisors  $\text{div}(P_{j_1 \dots j_q}^{i_1 \dots i_p}) \in \text{Div}(U)$  for all coefficient functions of  $P$ . The resulting coefficient of the complex subspace  $S$  of codimension 1 in  $\text{div}(P) \in \text{Div}(U)$  does not depend on the choice of the holomorphic coordinate system; e.g., for a vector field  $\sum_i X^i \frac{\partial}{\partial y^i} = \sum_{i,k} X^i \frac{\partial u^k}{\partial y^i} \frac{\partial}{\partial u^k}$  we have

$$\text{div}\left(\sum_i X^i \frac{\partial u^k}{\partial y^i}\right) \geq \min_i \text{div}\left(X^i \frac{\partial u^k}{\partial y^i}\right) = \min_i \left(\text{div}(X^i) + \text{div}\left(\frac{\partial u^k}{\partial y^i}\right)\right) \geq \min_i \text{div}(X^i).$$

Finally we define the divisor of  $P$  on  $X$  by gluing the local divisors for any holomorphic atlas of  $X$ . Note that a tensor field  $P$  is holomorphic if and only if  $\text{div}(P) \geq 0$ .

### 3. Invariant tensor fields

**3.1.** Let  $P$  be a  $G$ -invariant holomorphic tensor field of type  $\binom{p}{q}$  on  $V$ . Since  $\sigma$  is an tale map on  $V_{\text{reg}}$ , there is a unique holomorphic tensor field  $Q$  on  $Z_0$  of type  $\binom{p}{q}$  such that the pullback  $\sigma^*(Q)$  coincides with the restriction of  $P$  to  $V_{\text{reg}}$ . It is clear that the tensor field  $P$  is uniquely defined by  $Q$ .

Consider a holomorphic tensor field  $Q$  of type  $\binom{p}{q}$  on  $Z_0$  and its pullback  $\sigma^*(Q)$  which is a  $G$ -invariant holomorphic tensor field on  $V_{\text{reg}}$ . Then by the Hartogs extension theorem,  $\sigma^*(Q)$  has a  $G$ -invariant holomorphic extension to  $V$  iff it has a holomorphic extension to  $\sigma^{-1}(Z_1)$ .

Denote by  $\mathfrak{H}$  the set of all reflection hyperplanes corresponding to all complex reflections in  $G$  and, for each  $H \in \mathfrak{H}$ , by  $e_H$  the order of the cyclic subgroup of  $G$  fixing  $H$ . It is clear that  $\sigma(\cup_{H \in \mathfrak{H}} H)$  contains all strata of codimension 1. This implies immediately the following

**3.2. Proposition.** *If  $\mathfrak{H} = \emptyset$ , for each holomorphic tensor field  $P_0$  on  $Z_0$  the pullback  $\sigma^*(P_0)$  has a  $G$ -invariant holomorphic extension to  $V$ .  $\blacksquare$*

**3.3. The reflection divisor of the orbit space.** Consider the set  $R_Z$  of all hyper surfaces  $\sigma(H)$  in  $Z$ , where  $H$  runs through all reflection hyperplanes in  $V$ . Note that  $\sigma(H)$  is a complex subspace of  $Z_1$  of codimension 1. We endow each  $S = \sigma(H) \in R_Z$  with the label  $e_H$  of the hyperplane  $H$ . It is easily seen that this label does not depend on the choice of  $H$ , we denote it by  $e_S$  and we consider  $e_S \cdot S$  as an effective divisor on  $Z$  and we consider the effective divisor in  $Z_1$

$$D = D_{V/G} = D_Z = \sum_{S \in R_Z} e_S \cdot S,$$

which we call the *reflection divisor*.

**3.4. Basic example.** Let the cyclic group  $\mathbb{Z}_r = \mathbb{Z}/r\mathbb{Z}$  with generator  $\zeta_r = e^{2\pi i/r}$  act on  $\mathbb{C}$  by  $z \mapsto e^{2\pi i k/r} z$  for  $r \geq 2$ . The generating invariant is  $\tau(z) = z^r$ .

We consider first a holomorphic tensor field  $P = f(z)(dz)^{\otimes q} \otimes (\frac{\partial}{\partial z})^{\otimes p}$  on  $\mathbb{C}$ . It is invariant,  $\zeta_r^* P = P$ , if and only if  $f(\zeta_r z) = \zeta_r^{p-q} f(z)$ , so that in the expansion  $f(z) = \sum_{k \geq 0} f_k z^k$  at 0 of  $f$  the coefficient  $f_k \neq 0$  at most when  $k \cong p - q \pmod{r}$ . Writing  $p - q = rs + t$  with  $s \in \mathbb{Z}$  and  $0 \leq t < r$  we see that  $P$  is invariant if and only if  $f(z) = z^t g(z^r)$  for holomorphic  $g$ .

We use the coordinate  $y = \tau(z) = z^r$  on  $\mathbb{C}/\mathbb{Z}_r = \mathbb{C}$ ,  $\tau^* dy = r z^{r-1} dz$  and  $\tau^*(\frac{\partial}{\partial y}|_{\mathbb{C} \setminus 0}) = \frac{1}{r z^{r-1}} \frac{\partial}{\partial z}|_{\mathbb{C} \setminus 0}$ , and we write

$$\begin{aligned} P|_{\mathbb{C} \setminus 0} &= g(z^r) z^t (dz)^{\otimes q} \otimes (\frac{\partial}{\partial z})^{\otimes p} \\ &= g(y) z^t (r z^{r-1})^{p-q} (dy)^{\otimes q} \otimes (\frac{\partial}{\partial y})^{\otimes p} \\ &= g(y) z^{-rs} (r z^r)^{p-q} (dy)^{\otimes q} \otimes (\frac{\partial}{\partial y})^{\otimes p} \\ &= g(y) r^{p-q} y^{p-q-s} (dy)^{\otimes q} \otimes (\frac{\partial}{\partial y})^{\otimes p} \end{aligned}$$

(we omitted  $\tau^*$ ). Thus a holomorphic tensor field  $P$  of type  $\binom{p}{q}$  on  $\mathbb{C}$  is  $\mathbb{Z}_r$ -invariant if and only if  $P|_{\mathbb{C} \setminus 0} = \tau^* Q$  for a meromorphic tensor field

$$Q = g(y) y^m (dy)^{\otimes q} \otimes (\frac{\partial}{\partial y})^{\otimes p}$$

on  $\mathbb{C}$  with  $g$  holomorphic with  $g(0) \neq 0$  and with

$$m \geq p - q - s.$$

It is easily checked that the above inequality is equivalent to the following one

$$mr + (q - p)(r - 1) \geq 0.$$

**3.5.** Suppose  $\mathfrak{H} \neq \emptyset$ . Let  $z \in Z_1 \setminus Z_0$  and  $v \in \sigma^{-1}(z)$ . Then there is a unique hyperplane  $H \in \mathfrak{H}$  such that  $v \in H$  and the isotropy group  $G_v$  is isomorphic to a cyclic group. It is evident that the order  $r_z = e_H$  of  $G_v$  depends only on  $z = \sigma(v)$  and is locally constant on  $Z_1 \setminus Z_0$ .

By the holomorphic slice theorem (see [5], [6]) there is a  $G_v$ -invariant open neighborhood  $U_v$  of  $v$  in  $V$  such that the induced map  $U_v/G_v \rightarrow V/G$  is a local biholomorphic map at  $v$ .

Choose orthonormal coordinates  $z^1, \dots, z^n$  in  $V$  with respect to a  $G$ -invariant Hermitian inner product on  $V$ , so that  $H = \{z^n = 0\}$ . Then the ring  $\mathbb{C}[V]^{G_v}$  is generated by  $z^1, \dots, z^{n-1}, (z^n)^r$ , where  $r = r_z$ .

Put  $\tau^1 = z^1, \dots, \tau^{n-1} = z^{n-1}$ ,  $\tau^n = (z^n)^r$ , and  $\tau = (\tau^1, \dots, \tau^n) : U_v \rightarrow \mathbb{C}^n$ . Then there are holomorphic functions  $f^i$  ( $i = 1, \dots, n$ ) in an open neighborhood  $W_z$  of  $z \in \mathbb{C}^m$  such that  $\tau^a = f^a \circ \sigma|_{U_v}$ . On the other hand, we know that in an open neighborhood of  $v$  all  $\sigma^a$  for  $(a = 1, \dots, m)$  are holomorphic functions of the  $\tau^i$ . We denote by  $y^i$  the holomorphic function on  $Z$  such that  $\tau^i = y^i \circ \sigma$ . Then we can use  $y^i$  as coordinates of  $Z$  defined in the open neighborhood  $W_z \subseteq \mathbb{C}^m$  of  $z$ . Note that we found holomorphic coordinates near each point of  $Z_1$ , so we have:

**Corollary.** *The union  $Z_1$  of all codimension  $\leq 1$  strata, with the restriction of the sheaf  $\mathfrak{F}_{V/G}$ , is a complex manifold.  $\blacksquare$*

**3.6. The reflection divisor of a meromorphic tensor field on  $Z_1$ .** Let  $\Gamma_{\mathcal{M}}(T_q^p(Z_1))$  be the space of meromorphic tensor fields (i.e. with meromorphic coefficient functions in local holomorphic coordinates on the complex manifold  $Z_1$ ), and let  $P \in \Gamma_{\mathcal{M}}(T_q^p(Z_1))$ .

Let  $S$  be an irreducible component of  $Z_1 \setminus Z_0$  and let  $z \in S$ . Local coordinates  $y^1, \dots, y^n$  on  $U \subset Z_1$ , centered at  $z$ , are called adapted to the stratification of  $Z_1$  if  $S = \{y^n = 0\}$  near  $z$ . By definition the coordinates  $y^1, \dots, y^n$  from 3.5 have this property. Denote by  $\mathcal{O}_z$  the ring of germs of holomorphic functions and by  $\mathcal{M}_z$  the field of germs of meromorphic functions, both at  $z \in Z_1$ .

Let  $y^1, \dots, y^n$  be local coordinates on  $U \subset Z_1$ , centered at  $z$ , adapted to the stratification of  $Z_1$ . Then on  $U$  the meromorphic tensor field  $P$  is given by

$$P|_U = \sum_{i_1, \dots, i_p, j_1, \dots, j_q} P_{j_1 \dots j_q}^{i_1 \dots i_p} \frac{\partial}{\partial y^{i_1}} \otimes \dots \otimes \frac{\partial}{\partial y^{i_p}} \otimes dy^{j_1} \otimes \dots \otimes dy^{j_q}.$$

where the  $P_{j_1 \dots j_q}^{i_1 \dots i_p}$  are meromorphic on  $U$ . Let us fix one nonzero summand of the right hand side: for the coefficient function we have  $P_{j_1 \dots j_q}^{i_1 \dots i_p} = (y^n)^m f$  for some integer  $m$  such that the germs at  $z$  of  $y^n$ ,  $g$ , and  $h$  are pairwise relatively prime in  $\mathcal{O}_z$  where  $f = g/h \in \mathcal{M}_z$ . Suppose that the factor  $\frac{\partial}{\partial y^n}$  appears exactly  $p'$  times and the factor  $dy^n$  appears exactly  $q'$  times in this summand. The integer

$$\mu = mr + (q' - p')(r - 1),$$

a priori depending on  $z$ , is constant along an open dense subset of  $S$  and it is called the *reflection residuum* of the summand at  $S$ . Finally let  $\mu_S(P)$  be the minimum of the reflection residua at  $S$  of all summands of  $P$  in the representation of  $P$ .

Let  $\tilde{y}^1, \dots, \tilde{y}^n$  be arbitrary local coordinates on  $U \subset Z_1$ , centered at  $z$ , adapted to the stratification of  $Z_1$ . In a neighborhood of  $z$  we have  $y^n = f\tilde{y}^n$ ,

where  $f$  is a holomorphic function such that  $f(z) \neq 0$ . Remark that  $\tilde{y}^n$  divides  $\frac{\partial y^n}{\partial \tilde{y}^i}$  and  $\frac{\partial \tilde{y}^n}{\partial y^i}$  ( $i = 1, \dots, n$ ) in  $\mathcal{O}_z$ . A straightforward calculation using the above remark shows that the values of  $\mu_S(P)$  calculated in the coordinates  $\tilde{y}^i$  and in the coordinates  $y^i$  are the same. Then  $\mu_S(P)$  does not depend on the choice of the system of local coordinates adapted to the stratification of  $Z_1$ . For details see [4]: there we checked this in the algebraic geometry setting where the use of tensor fields is less familiar.

We now can define the *reflection divisor*

$$\operatorname{div}_D(P) = \operatorname{div}_{D_{V/G}}(P) \in \operatorname{Div}(U)$$

as follows: take the divisor  $\operatorname{div}(P)$ , and for each irreducible component  $S$  of  $Z_1 \setminus Z_0$  do the following: if  $S$  appears in the support of  $\operatorname{div}(P) \in \operatorname{Div}(U)$ , replace its coefficient by  $\mu_S(P)$ ; if it does not appear, add  $\mu_S(P) \cdot S$  to it. If  $S$  is not contained in  $Z_1 \setminus Z_0$ , we keep its coefficient in  $\operatorname{div}(P)$ .

Finally we glue the global *reflection divisor*  $\operatorname{div}_D(P) \in \operatorname{Div}(Z_1)$  from the local ones, using a holomorphic atlas for  $Z_1$ .

**3.7. Theorem.** *Let  $G \subset GL(V)$  be a finite group, with reflection divisor  $D = D_{V/G} = D_Z$ . Then we have:*

- *Let  $P$  be a holomorphic  $G$ -invariant tensor field on  $V$ . Then the reflection divisor  $\operatorname{div}_D(\pi_*P) \geq 0$ .*
- *Let  $Q \in \Gamma_{\mathcal{M}}(T_q^p(Z_1))$  be a meromorphic tensor field on  $Z_1$ . Then the  $G$ -invariant meromorphic tensor field  $\pi^*Q$  extends to a holomorphic  $G$ -invariant tensor field on  $V$  if and only if  $\operatorname{div}_D(Q) \geq 0$ .*

*The above remains true for  $G$ -invariant holomorphic tensor fields defined in a  $G$ -stable open subset of  $V$ .*

**Proof.** This follows directly from Hartogs' extension theorem, the basic example 3.4 using  $y^1, \dots, y^{n-1}$  as dummy variables, and the definition of the reflection divisor  $\operatorname{div}_D(P)$  as explained in 3.6. ■

**3.9. Corollary.** *The mapping  $\sigma$  establishes an injective correspondence between the space of holomorphic  $G$ -invariant tensor fields of type  $\binom{p}{q}$  on  $V$  which are skew-symmetric with respect to the covariant entries, and the space of holomorphic tensor fields on  $Z_1$  of the same type and the same skew-symmetry condition. If  $p = 0$  the correspondence is bijective.*

*The above remains true for  $G$ -invariant holomorphic tensor fields defined in a  $G$ -stable open subset of  $V$ .*

**Proof.** Let  $P$  be a holomorphic  $G$ -invariant tensor field on  $V$  satisfying the conditions of the corollary. For each nonzero decomposable summand of  $\pi_*P$  take the integers  $m$ ,  $p'$ , and  $q'$  defined in 3.6. By skew symmetry of  $P$  we have  $q' \leq 1$ . By Theorem 3.7 we get  $\operatorname{div}_D(\pi_*P) \geq 0$  and thus  $mr \geq (p' - q')(r - 1) > -r$ . So  $m \geq 0$  and the summand is holomorphic on  $Z_1$ .

If  $Q$  is a holomorphic differential form on  $Z_1$  its pullback  $\sigma^*Q$  is a  $G$ -invariant holomorphic form on  $\sigma^{-1}(Z_1)$  and then has a holomorphic extension to the whole of  $V$ . ■

**3.10. Remarks.** Note that Corollary 3.9 is a generalization of Solomon's theorem (see [10]): *If  $G \subset GL(V)$  is a finite complex reflection group then every  $G$ -invariant polynomial exterior  $q$ -form  $\omega$  on  $V$  can be written as  $\omega = \sigma^* \varphi$  for a polynomial  $q$ -form  $\varphi$  on  $\mathbb{C}^n$ , where  $\sigma = (\sigma^1, \dots, \sigma^n) : V \rightarrow \mathbb{C}^n$  is the mapping consisting of a minimal system of homogeneous generators of  $\mathbb{C}[V]^G$ .*

Actually, in the case of a reflection group  $Z = \mathbb{C}^n$  and each holomorphic  $\binom{p}{q}$ -tensor field  $Q$  on  $Z_1$  has a holomorphic extension to  $Z$  by Hartogs' extension theorem.

#### 4. Invariant complex connections

**4.1.** Let  $\Gamma$  be a holomorphic  $G$ -invariant complex connection on  $V$ . Then the image  $\sigma_* \Gamma$  of  $\Gamma$  under the map  $\sigma$  defines a holomorphic complex connection on  $Z_0$ .

Let  $z \in Z_1 \setminus Z_0$ ,  $v \in \sigma^{-1}(z)$ , and  $r$  the order of  $G_v$ . Consider the coordinates  $z^i$  in  $V$  defined in 3.5. Denote by  $\Gamma_{jk}^i$  the components of the connection  $\Gamma$  with respect to these coordinates. By assumption, the  $\Gamma_{jk}^i$  are holomorphic functions on  $V$ . Recall the standard formula for the image  $\gamma$  of  $\Gamma$  under a holomorphic diffeomorphism  $f = (y^a(x^i))$

$$\gamma_{bc}^a \circ f = \frac{\partial y^a}{\partial x^i} \frac{\partial x^j}{\partial y^b} \frac{\partial x^k}{\partial y^c} \Gamma_{jk}^i(x^l) - \frac{\partial^2 y^a}{\partial x^i \partial x^j} \frac{\partial x^i}{\partial y^b} \frac{\partial x^j}{\partial y^c}.$$

Remark that the similar formula is true for the transformation of the components of connection under the change of coordinates.

Consider the generator  $g$  of the cyclic group  $G_v$  given by 3.5. Since  $g$  acts linearly, the connection reacts to it like a  $\binom{1}{2}$ -tensor field. Thus by the considerations of 3.4 we get in the notation of 3.5, where  $i, j, k = 1, \dots, n-1$ :

$$\begin{aligned} \Gamma_{jk}^i &= \tilde{\Gamma}_{jk}^i \circ \sigma, & \Gamma_{jk}^n &= \frac{1}{r} z^n \tilde{\Gamma}_{jk}^n \circ \sigma, & \Gamma_{jn}^i &= r(z^n)^{r-1} \tilde{\Gamma}_{jn}^i \circ \sigma, \\ \Gamma_{nk}^i &= r(z^n)^{r-1} \tilde{\Gamma}_{nk}^i \circ \sigma, & \Gamma_{jn}^n &= \tilde{\Gamma}_{jn}^n \circ \sigma, & \Gamma_{nk}^n &= \tilde{\Gamma}_{nk}^n \circ \sigma, \\ \Gamma_{nn}^i &= r^2(z^n)^{r-2} \tilde{\Gamma}_{nn}^i \circ \sigma, & \Gamma_{nn}^n &= r(z^n)^{r-1} \tilde{\Gamma}_{nn}^n \circ \sigma, \end{aligned}$$

where the  $\tilde{\Gamma}_{bc}^a$  are holomorphic functions of the coordinates  $y^a$  ( $a = 1, \dots, n$ ) introduced in 3.5.

Using the transformation formula for connections, we get the following formulas for the components  $\gamma_{bc}^a$  of the meromorphic connection  $\sigma_* \Gamma$  with respect to the coordinates  $y^a$

$$\begin{aligned} \gamma_{jk}^i &= \tilde{\Gamma}_{jk}^i, & \gamma_{jk}^n &= y^n \tilde{\Gamma}_{jk}^n, & \gamma_{jn}^i &= \tilde{\Gamma}_{jn}^i, & \gamma_{nk}^i &= \tilde{\Gamma}_{nk}^i, & (4.1.1) \\ \gamma_{jn}^n &= \tilde{\Gamma}_{jn}^n, & \gamma_{nk}^n &= \tilde{\Gamma}_{nk}^n, & \gamma_{nn}^i &= \frac{1}{y^n} \tilde{\Gamma}_{nn}^i, & \gamma_{nn}^n &= \tilde{\Gamma}_{nn}^n - \frac{r-1}{ry^n}. \end{aligned}$$

Let  $\tilde{y}^a$  for  $a = 1, \dots, n$  be other local coordinates centered at  $z$  and adapted to the stratification of  $Z_1$ . Then in a neighborhood of  $z$  we have

$$y^n = f \tilde{y}^n, \quad \tilde{y}^n = \tilde{f} y^n,$$

where  $f$  and  $\tilde{f}$  are holomorphic functions in a neighborhood of  $z$  and  $\tilde{f}f = 1$ . Then we have

$$\frac{\partial y^n}{\partial \tilde{y}^i} = \frac{\partial f}{\partial \tilde{y}^i} \tilde{y}^n, \quad \frac{\partial \tilde{y}^n}{\partial y^i} = \frac{\partial \tilde{f}}{\partial y^i} y^n \quad (i = 1, \dots, n-1)$$

and on  $S = \{y^n = 0\}$

$$\frac{\partial y^n}{\partial \tilde{y}^n} = f, \quad \frac{\partial \tilde{y}^n}{\partial y^n} = \tilde{f}.$$

Using these formulas one can check that in the coordinates  $\tilde{y}^a$  the formulas 4.1.1 have the same form as in the coordinates  $y^a$ . For example, for the new component  $\tilde{\gamma}_{nn}^n$  we have

$$\tilde{\gamma}_{nn}^n + \frac{r-1}{r\tilde{y}^n} = \frac{(r-1) \left( 1 - \tilde{f} \frac{\partial \tilde{y}^n}{\partial y^n} \left( \frac{\partial y^n}{\partial \tilde{y}^n} \right)^2 \right)}{r y^n \tilde{f}} + h,$$

where  $h$  is a holomorphic function near  $z$ . Since on  $S = \{y^n = 0\}$  we have

$$1 - \tilde{f} \frac{\partial \tilde{y}^n}{\partial y^n} \left( \frac{\partial y^n}{\partial \tilde{y}^n} \right)^2 = 1 - \tilde{f}^2 f^2 = 0,$$

$y^n$  divides in  $\mathcal{O}_z$  the function

$$1 - \tilde{f} \frac{\partial \tilde{y}^n}{\partial y^n} \left( \frac{\partial y^n}{\partial \tilde{y}^n} \right)^2.$$

Thus

$$\tilde{\gamma}_{nn}^n + \frac{r-1}{r\tilde{y}^n}$$

is holomorphic in a neighborhood of  $z$ .

**4.2. Theorem.** *Let  $\gamma$  be a holomorphic complex linear connection on  $Z_0$  such that for each  $z \in Z_1 \setminus Z_0$  it has an extension to a neighborhood of  $z$  whose components in the coordinates adapted to the stratification of  $Z_1$  are defined by the formulas 4.1.1 where  $\tilde{\Gamma}_{bc}^a$  are holomorphic. Then there is a unique  $G$ -invariant holomorphic complex linear connection  $\Gamma$  on  $V$  such that  $\sigma_*\Gamma$  coincides with  $\gamma$  on  $Z_0$ . This remains true if we replace  $V$  by a  $G$ -open subset of  $G$ .*

**Proof.** Since  $\sigma$  is tale on the principal stratum, there is a unique  $G$ -invariant complex linear connection  $\Gamma_0$  on  $\sigma^{-1}(Z_0)$  such that  $\sigma_*\Gamma_0 = \gamma$ . The condition of the theorem implies that the connection  $\Gamma_0$  has a holomorphic extension to  $\sigma^{-1}(Z_1)$ . Then by Hartogs' extension theorem the connection  $\Gamma_0$  has a unique holomorphic extension  $\Gamma$  to the whole of  $V$ .  $\blacksquare$

## 5. Lifts of diffeomorphisms of orbit spaces

**5.1.** Let  $G$  and  $G'$  be finite subgroups of  $GL(V)$  and  $GL(V')$  and let  $F$  be a holomorphic diffeomorphism  $V \rightarrow V'$  which maps  $G$ -orbits to  $G'$ -orbits bijectively. Then the map  $F$  induces an isomorphism  $f$  of the sheaves  $\mathfrak{F}_{V/G} \rightarrow \mathfrak{F}_{V'/G'}$ , i.e. a holomorphic diffeomorphism of orbit spaces  $V/G$  and  $V'/G'$ .

**Lemma.** *There is a unique isomorphism  $a : G \rightarrow G'$  such that  $F \circ g = a(g) \circ F$  for every  $g \in G$ .*

Note that  $a$  and its inverse  $a^{-1}$  map complex reflections to complex reflections.

**Proof.** The cardinalities of the two groups are the same since  $F$  maps a generic regular orbit to a regular orbit. Consequently, it maps regular points to regular points and we have  $\sigma' \circ F = f \circ \sigma : V \rightarrow V'/G'$  for a holomorphic diffeomorphism  $f : V/G \rightarrow V'/G'$ , where  $\sigma : V \rightarrow V/G$  and  $\sigma' : V' \rightarrow V'/G'$  are the quotient projections.

Fix some  $G$ -regular  $v \in V$ . Then  $F(v)$  and  $F(gv)$  for  $g \in G$  are regular points of  $V'$  of the same orbit. Therefore, there is a unique  $a(g) \in G'$  such that  $F(gv) = a(g)(F(v))$ . We have  $\sigma' \circ F \circ g = f \circ \sigma \circ g = f \circ \sigma = \sigma' \circ F = \sigma' \circ a(g) \circ F$ . Since  $\sigma'$  is tale on  $V'_{\text{reg}}$  we see that  $F \circ g = a(g) \circ F$  locally near  $v$  and thus globally. By uniqueness, the map  $g \rightarrow a(g)$  is an isomorphism of  $G$  onto  $G'$ . ■

In this section we study when a diffeomorphism  $f$  of the orbit spaces  $Z \rightarrow Z'$  has a holomorphic lift  $F$ .

**5.2. Corollary.** *Let  $F : V \rightarrow V$  be a holomorphic diffeomorphism which maps  $G$ -orbits onto  $G'$ -orbits, and  $f : Z \rightarrow Z'$  the corresponding holomorphic diffeomorphism of the orbit spaces. Then  $f$  maps the isotropy type stratification of  $Z$  onto that of  $Z'$  and, moreover, it maps  $D_Z$  to  $D_{Z'}$ .*

**Proof.** This follows from Lemma 5.1 and the definition 3.3 of the reflection divisor. ■

**5.3. Theorem.** *Let  $G$  and  $G'$  be two finite subgroups of  $GL(V)$  and let  $f : Z \rightarrow Z'$  be a holomorphic diffeomorphism of the corresponding orbit spaces such that  $f(Z_0) = Z'_0$  and  $f_*(D_Z) = D_{Z'}$ . If  $Q$  is a holomorphic tensor field of type  $\binom{p}{q}$  on  $Z_0$  which satisfies the conditions of Theorem 3.7, then  $f_*(Q)$  also satisfies these conditions on  $Z'_0$  and thus there exists a unique  $G'$ -invariant holomorphic tensor field  $Q'$  of type  $\binom{p}{q}$  such that  $\sigma'_*Q'$  coincides with  $f_*Q$  on  $Z'_0$ .*

*This is also true for holomorphic connections if we replace Theorem 3.7 by Theorem 4.2. The theorem remains true if we replace  $V$  by invariant open subsets of  $V$ .*

**Proof.** Since  $f(Z_0) = Z'_0$  the tensor field  $f_*Q$  is also holomorphic on  $Z'_0$ . Let  $z \in Z_1 \setminus Z_0$ . Then there is a complex space  $S \in R_Z$  of codimension 1 such that  $z \in S$ . By assumption  $f(z) \in Z'_1 \setminus Z'_0$  and  $f(z) \in f(S) \in R_{Z'}$  and  $r_z = e_S = e_{f(S)} = r_{f(z)}$ . Now, obviously  $f_*Q$  satisfies the conditions of Theorem 3.7 at  $f(x)$ . Thus there exists a  $G'$ -invariant holomorphic tensor field  $Q'$  on  $V$  with  $\sigma'_*Q' = f_*Q$ .

A similar argument applies to connections. ■

**5.4 Theorem.** *Let  $G$  and  $G'$  be two finite subgroups of  $GL(V)$ . Let  $f : Z \rightarrow Z'$  be a holomorphic diffeomorphism of the orbit spaces such that  $f(Z_0) = Z'_0$  and  $f_*(D_Z) = D_{Z'}$ .*

*Then  $f$  lifts to a holomorphic diffeomorphism  $F : V \rightarrow V$ , i.e.  $\sigma' \circ F = f \circ \sigma$ .*

The local version is also true. Namely, if  $B$  is a ball in the vector space  $V$  centered at 0 (for an invariant Hermitian metric),  $U = \sigma(B)$ , and  $f : U \rightarrow Z'$  is a local holomorphic diffeomorphism of  $U$  onto a neighborhood  $U'$  of  $\sigma'(0)$  such that  $f(U \cap Z_0) = U' \cap Z'_0$  and  $f$  maps  $D_Z \cap U$  to  $D_{Z'} \cap U'$ , then there is a holomorphic lift  $F : B \rightarrow V$ .

**Proof.** Let  $\Gamma$  be the natural flat connection on  $V$ . Then  $\Gamma$  is uniquely defined by the holomorphic connection  $\sigma_*\Gamma$  on  $Z_0$  which satisfies the conditions of Theorem 4.2. By Theorem 5.3 there is a unique  $G$ -invariant holomorphic complex linear connection  $\Gamma'$  on  $V$  such that  $\sigma'_*\Gamma'$  coincides with  $f_*(\sigma_*\Gamma)$  on  $Z'_0$ . It is evident that  $\Gamma'$  is a torsion free flat connection, since  $\Gamma$  is it and  $\Gamma'$  is locally isomorphic to  $\Gamma$  on an open dense subset.

Let  $v \in V$  be  $G$ -regular and let  $v' \in V$  be  $G'$ -regular, such that  $(f \circ \sigma)(v) = \sigma'(v')$ . Then there is a biholomorphic map  $F$  of a neighborhood  $W$  of  $v$  onto a neighborhood of  $v'$  such that  $\sigma' \circ F = f \circ \sigma$  on  $W$  and  $F(v) = v'$ . Moreover by construction  $F$  is a locally affine map of the affine space  $(V, \Gamma)$  into  $(V, \Gamma')$  equipped with the above structures of locally affine spaces, thus we have

$$F = \exp_{v'}^{\Gamma'} \circ T_v F \circ (\exp_v^\Gamma)^{-1} \quad (1)$$

where  $\exp_v^\Gamma : T_v V \rightarrow V$  is the holomorphic geodesic exponential mapping centered at  $v$  given by the connection  $\Gamma$  and its induced spray. It is globally defined, thus complete and a holomorphic diffeomorphism since  $\Gamma$  is the standard flat connection. Likewise  $\exp_{v'}^{\Gamma'}$  is the holomorphic exponential mapping of the flat connection  $\Gamma'$ . The formula above extends  $F$  to a globally defined holomorphic mapping if  $\exp_{v'}^{\Gamma'} : T_v V \rightarrow V$  is also globally defined (complete). Assume for contradiction that this is not the case. Let  $F$  be maximally extended by equation (1); it still projects to  $f : Z \rightarrow Z'$ . We consider  $\exp_{v'}^{\Gamma'}$  as a real exponential mapping, and then there is a real geodesic which reaches infinity in finite time and this is the image under  $F$  of a finite part  $\exp_v^\Gamma([0, t_0]w)$  of a real geodesic of  $\Gamma$  emanating at  $v$ . The sequence  $\exp_v^\Gamma((t_0 - 1/n)w)$  converges to  $\exp_v^\Gamma(t_0 w)$  in  $V$ , but its image under  $F$  diverges to infinity by assumption. On the other hand, the image under  $F$  is contained in the set  $(\sigma')^{-1}(f\sigma(\exp_v^\Gamma([0, t_0]w)))$  which is compact since  $\sigma'$  is a proper mapping. Contradiction.

Any holomorphic lift  $F$  of a holomorphic diffeomorphism  $f$  is a holomorphic diffeomorphism of  $V$  which maps  $G$ -orbits onto  $G'$  orbits, by the following argument: Let  $F'$  be a holomorphic lift of  $f^{-1}$ . Evidently the map  $F' \circ F$  preserves each  $G$ -orbit. Then, for a  $G$ -regular point  $v \in V$ , there is a  $g \in G$  such that  $F' \circ F = g$  in a neighborhood of  $v$  and, then, on the whole of  $V$ . Similarly  $F \circ F' = g' \in G'$ . This implies that  $F$  is a holomorphic diffeomorphism of  $V$ . By definition the lift  $F$  respects the partitions of  $V$  into orbits. ■

We give a second proof of Theorem 5.4 based on the known results about the fundamental groups of  $V_{\text{reg}}$  and  $Z_0$  for finite complex reflection groups. It is an extension of the proof of [8], using results of [2].

**5.5. Lemma.** *Let  $G$  and  $G'$  be two finite subgroups of  $GL(V)$  and let  $f : Z \rightarrow Z'$  be a holomorphic diffeomorphism of the corresponding orbit spaces. Suppose  $v_0 \in V_{\text{reg}}$ ,  $v'_0 \in V'_{\text{reg}}$ , and  $f \circ \sigma(v_0) = \sigma'(v'_0)$ . If the image of the fundamental group  $\pi_1(V_{\text{reg}}, v_0)$  under  $f \circ \sigma$  is contained in the subgroup  $\sigma'_*(\pi_1(V_{\text{reg}}, v'_0))$  of  $\pi_1(Z'_0, \sigma'(v'_0))$ , the holomorphic lift of  $f \circ \sigma$  mapping  $v_0$  to  $v'_0$  exists.*

**Proof.** Consider the restriction  $\varphi$  of the map  $f \circ \sigma$  to  $V_{\text{reg}}$ . Since the restriction of  $\sigma$  to  $V_{\text{reg}}$  is a covering map onto  $Z_0$ , the condition of the lemma implies that there is a holomorphic lift  $F_0$  of the map  $\varphi$  to  $V_{\text{reg}}$ . The map  $F_0$  is bounded on  $B \cap V_{\text{reg}}$  for each compact ball  $B$  in  $V$  since its image is contained in the compact set  $(\sigma')^{-1}(f(\sigma(B)))$ . Then by the Riemann extension theorem  $F_0$  has a holomorphic extension  $F$  to  $V$  which is the required holomorphic lift of  $f$ . ■

**5.6.** Next we prove Theorem 5.4 in the case when the group  $G$  is generated by complex reflections. Put

$$B := \pi_1(Z_0) \quad \text{and} \quad P := \pi_1(V_{\text{reg}}).$$

The groups  $B$  and  $P$  are called the *braid group* and the *pure braid group* associated to  $G$ , respectively. It is clear that the map  $\sigma$  induces an isomorphism of  $P$  onto a subgroup of  $B$ .

The following results about the groups  $B$  and  $P$  are well known (see, for example, [2]). The braid group  $B$  is generated by those elements which are represented by loops around the hypersurfaces  $\sigma(H)$  for  $H \in \mathfrak{H}$ . The pure braid group  $P$  is generated by the elements of  $B$  of the type  $s^{\epsilon_H}$ , where  $s$  is any of the above generators of  $B$  represented by a loop around the hypersurface  $\sigma(H)$ . This implies the following

**Proposition.** *Suppose the group  $G$  is generated by complex reflections. Let  $f$  be a holomorphic diffeomorphism of the orbit space  $Z = \mathbb{C}^n$  with  $f(Z_0) = Z_0$  which also preserves  $D_Z$ . Then  $f|_{Z_0}$  preserves the subgroup  $P$  of  $B$ .* ■

The following proposition is an immediate consequence of Lemma 5.5 and Proposition 5.6.

**5.7. Proposition.** *Suppose the groups  $G$  and  $G'$  are generated by complex reflections. Let  $f : Z \rightarrow Z'$  be a holomorphic diffeomorphism between the corresponding orbit spaces, such that  $f(Z_0) = Z'_0$  and  $f_*(D_Z) = D_{Z'}$ .*

*Then  $f$  has a holomorphic lift  $F$  to  $V$ .* ■

**Second proof of 5.4.** Now let  $G \subset GL(V)$  be a finite group and let  $G_1$  be the subgroup generated by all complex reflections in  $G$ . Clearly  $G_1$  is a normal subgroup of  $G$ . Let  $G_2 = G/G_1$ . Let  $\sigma_1^1, \dots, \sigma_1^n$  be a system of homogeneous generators of  $\mathbb{C}[V]^{G_1}$  and  $\sigma_1 : V \rightarrow \mathbb{C}^n$  the corresponding orbit map. Then the action of  $G$  on  $V$  induces the action of the group  $G_2$  on  $V_1 := \mathbb{C}^n = \sigma_1(V)$ . Since each representation of the group  $G_2$  is completely reducible, by standard arguments of invariant theory, we may assume that the generators  $\sigma_1^i$ 's are chosen in such a way that the above action of  $G_2$  on  $V_1 = \mathbb{C}^n$  is linear. Then the representation of  $G_2$  on  $V_1$  contains no complex reflections. Let  $\sigma_2^1, \dots, \sigma_2^m$  be a system of homogeneous generators of  $\mathbb{C}[V_1]^{G_2}$  and  $\sigma_2 : V_1 \rightarrow \mathbb{C}^m$  the corresponding orbit map. Then  $\sigma^i = \sigma_2^i \circ \sigma_1$  ( $i = 1, \dots, m$ ) is a system of generators of  $\mathbb{C}[V]^G$  with orbit map  $\sigma = \sigma_2 \circ \sigma_1$ . Similarly for  $G'$ .

Let  $f : Z \rightarrow Z'$  be a holomorphic diffeomorphism, such that  $f(Z_0) = Z'_0$  and  $f_*(D_Z) = D_{Z'}$ . Since the group  $G_2$  contains no complex reflections the set  $V_{1,\text{reg}}$  of regular points of the action of  $G_2$  on  $V_1$  is obtained from  $V_1$  by removing some subsets of codimension  $\geq 2$ . And similarly for  $G'$ . Then the fundamental

group  $\pi_1(V_{1,\text{reg}}) = \pi_1(V_1) = 0$  is trivial and by lemma 5.5 the diffeomorphism  $f$  has a holomorphic lift  $F_1 : V_1 \rightarrow V_1'$  which is a holomorphic diffeomorphism mapping the principal stratum to the principal stratum, and the reflection divisor to the reflection divisor, since  $G_2$  contains no complex reflections on  $V_1$ . Thus the diffeomorphism  $F_1$  has a holomorphic lift to  $V$  by Proposition 5.7, which is a holomorphic lift of  $f$ . ■

## 6. An intrinsic characterization of a complex orbifold

We recall the definition of orbifold.

**6.1. Definition.** [11] *Let  $X$  be a Hausdorff space. An atlas of a smooth  $n$ -dimensional orbifold on  $X$  is a family  $\{U_i\}_{i \in I}$  of open sets that satisfy:*

1.  $\{U_i\}_{i \in I}$  is an open cover of  $X$ .
2. For each  $i \in I$  we have a local uniformizing system consisting of a triple  $(\tilde{U}_i, G_i, \varphi_i)$ , where  $\tilde{U}_i$  is a connected open subset of  $\mathbb{R}^n$  containing the origin,  $G_i$  is a finite group of diffeomorphisms acting effectively and properly on  $\tilde{U}_i$ , and  $\varphi_i : \tilde{U}_i \rightarrow U_i$  is a continuous map of  $\tilde{U}_i$  onto  $U_i$  such that  $\varphi_i \circ g = \varphi_i$  for all  $g \in G_i$  and the induced map of  $\tilde{U}_i/G_i$  onto  $U_i$  is a homeomorphism. The finite group  $G_i$  is called a local uniformizing group.
3. Given  $\tilde{x}_i \in \tilde{U}_i$  and  $\tilde{x}_j \in \tilde{U}_j$  such that  $\varphi_i(\tilde{x}_i) = \varphi_j(\tilde{x}_j)$ , there is a diffeomorphism  $g_{ij} : \tilde{V}_j \rightarrow \tilde{V}_i$  from a neighborhood  $\tilde{V}_j \subseteq \tilde{U}_j$  of  $\tilde{x}_j$  onto a neighborhood  $\tilde{V}_i \subseteq \tilde{U}_i$  of  $\tilde{x}_i$  such that  $\varphi_j = \varphi_i \circ g_{ij}$ .

Two atlases are equivalent if their union is again an atlas of a smooth orbifold on  $X$ . An orbifold is the space  $X$  with an equivalence class of atlases of smooth orbifolds on  $X$ .

If we take in the definition of orbifold  $\mathbb{C}^n$  instead of  $\mathbb{R}^n$  and require that  $G_i$  is a finite group of holomorphic diffeomorphisms acting effectively and properly on  $\tilde{U}_i$  and the maps  $g_{ij}$  are biholomorphic, we get the definition of complex analytic  $n$ -dimensional orbifold.

**6.2. Theorem.** [11] *Let  $M$  be a smooth manifold and  $G$  a proper discontinuous group of diffeomorphisms of  $M$ . Then the orbit space  $M/G$  has a natural structure of smooth  $n$ -dimensional orbifold. If  $M$  is a complex  $n$ -dimensional manifold and  $G$  is a group of holomorphic diffeomorphisms of  $M$ , the orbit space  $M/G$  is a complex  $n$ -dimensional orbifold.*

**6.3 Definitions.** In the definition of atlas of a complex orbifold on  $X$  we can always take  $\tilde{U}_i$  to be balls of the space  $\mathbb{C}^n$  (with respect to some Hermitian metric) centered at the origin and the finite subgroups  $G_i$  to be subgroups of the  $GL(n)$  acting naturally on  $\mathbb{C}^n$ . In the sequel we consider atlases of complex orbifolds satisfying these conditions.

Let  $X$  be a complex orbifold with an atlas  $(\tilde{U}_i, G_i, \varphi_i)$ . A function  $f : U_i \rightarrow \mathbb{C}$  is called holomorphic if  $f \circ \varphi_i$  is a holomorphic function on  $\tilde{U}_i$ . The germs of holomorphic functions on  $X$  define a sheaf  $\mathfrak{F}_X$  on  $X$ . It is evident that the sheaf  $\mathfrak{F}_X$  depends only on the structure of complex orbifold on  $X$ .

Consider a uniformizing system  $(\tilde{U}_i, G_i, \varphi_i)$  of the above atlas and the corresponding action of  $G_i$  on  $\mathbb{C}^n$ . Then we have the isotropy type stratification of the orbit space  $\mathbb{C}^n/G_i$ , the induced stratification of  $U_i$ , and the divisor  $D_{U_i}$ .

By corollary 5.2 we get the *stratification on  $X$*  by gluing the strata on the  $U_i$ 's. Denote by  $X_0$  the principal stratum of this stratification. By definition, for each  $x \in X_0$ , for each uniformizing system  $(\tilde{U}_i, G_i, \varphi_i)$ , and for each  $y \in \tilde{U}_i$  such that  $\varphi_i(y) = x$ , the isotropy group  $G_y$  of  $y$  is trivial. Note that  $X_0$  is a complex manifold. Note that  $X_1$  is also a complex manifold since this holds locally as noted in 3.5.

Denote by  $R_X$  the *set of all strata of codimension 1* of  $X$ . Since the pullbacks of the reflection divisors  $D_{U_i}$  to  $U_i \cap U_j$  agree by 5.2 we may glue them into the reflection divisor  $D_X$  on  $X_1$ .

**6.4. Definition.** *Let  $X$  and  $\tilde{X}$  be two smooth orbifolds. The orbifold  $\tilde{X}$  is called a covering orbifold for  $X$  with a projection  $p : \tilde{X} \rightarrow X$  if  $p$  is a continuous map of underlying topological spaces and each point  $x \in X$  has a neighborhood  $U = \tilde{U}/G$  (where  $\tilde{U}$  is an open subset of  $\mathbb{R}^n$ ) for which each component  $V_i$  of  $p^{-1}(U)$  is isomorphic to  $\tilde{U}/G_i$ , where  $G_i \subseteq G$  is some subgroup. The above isomorphisms  $U = \tilde{U}/G$  and  $V_i = \tilde{U}/G_i$  must respect the projections.*

Note that the projection  $p$  in the above definition is not necessarily a covering of the underlying topological spaces. It is clear that a covering orbifold for a complex orbifold is a complex orbifold. Hereafter we suppose that all orbifolds and their covering orbifolds are connected.

**6.5. Theorem.** [11] *An orbifold  $X$  has a universal covering orbifold  $p : \tilde{X} \rightarrow X$ . More precisely, if  $x \in X_0$ ,  $\tilde{x} \in \tilde{X}_0$  and  $p(\tilde{x}) = x$ , for any other covering orbifold  $p' : \tilde{X}' \rightarrow X$  and  $\tilde{x}' \in \tilde{X}'$  such that  $p'(\tilde{x}') = x$  there is a cover  $q : \tilde{X} \rightarrow \tilde{X}'$  such that  $p = p' \circ q$  and  $q(\tilde{x}) = \tilde{x}'$ . For any points  $\tilde{x}, \tilde{x}' \in p^{-1}(x)$  there is a deck transformation of  $\tilde{X}$  taking  $\tilde{x}$  to  $\tilde{x}'$ .*

Now we prove the main theorem of this section.

**6.6. Theorem.** *An  $n$ -dimensional complex orbifold  $X$  is uniquely determined by the sheaf of holomorphic functions  $\mathfrak{F}_X$ , the principal stratum  $X_0$ , and the reflection divisor  $D_X$ .*

**Proof.** For each  $x \in X$ , there exists  $V = \mathbb{C}^m$ , a finite group  $G \subset GL(m)$ , a ball  $B$  in  $V$  centered at 0, an open subset  $U$  of  $X$  containing  $x$ , and an isomorphism  $\psi : \pi(B) \rightarrow U$  between the sheaves  $\mathfrak{F}_Z|_{\pi(B)}$  and  $\mathfrak{F}_X|_U$ . Consider the map  $\pi : V \rightarrow Z = V/G$ , the stratum  $Z_0$  and the reflection divisor  $D_Z$ . We suppose also that  $\psi(Z_0 \cap B/G) \subseteq X_0$  and  $\psi_*(D_{\pi(B)}) = D_U$ . It suffices to prove that the germ of the uniformizing system  $\{B, G, \psi \circ \pi|_B\}$  at  $x$  is the germ of some uniformizing system of the orbifold  $X$ .

Let  $y \in V_{\text{reg}} \cap B$ . Then the ring  $\mathfrak{F}_Z(\pi(y))$  of germs of  $\mathfrak{F}_Z$  at  $\pi(y)$  is isomorphic to the ring of germs of holomorphic functions on  $\mathbb{C}^n$  at 0 and thus we have  $m = n$ .

Consider the uniformizing system  $(\tilde{U}_i, G_i, \varphi_i)$  of the orbifold  $X$ , where  $\tilde{U}_i$  is a ball in  $\mathbb{C}^n$  centered at the origin,  $G_i$  is a finite subgroup of the group  $GL(n)$  acting naturally on  $V = \mathbb{C}^n$ , and where  $\varphi_i(0) = x$ . Consider the map  $\pi_i : V \rightarrow V/G_i$  given by some system of generators of  $\mathbb{C}[V]^{G_i}$ . We may assume

that  $\varphi_i = \psi_i \circ \pi_i|_{\tilde{U}_i}$ , where  $\psi_i : \mathfrak{F}_{\tilde{U}_i/G_i} \rightarrow \mathfrak{F}_{U_i}$  is an isomorphism of sheaves.

$$\begin{array}{ccccc}
 \mathbb{C}^n & \longleftarrow & B & \xrightarrow{\quad F \quad} & \tilde{U}_i & \hookrightarrow & \mathbb{C}^n \\
 & & \downarrow \pi & & \downarrow \pi_i & & \\
 & & B/G & \xrightarrow{\quad f \quad} & \tilde{U}_i/G_i & & \\
 & & \searrow \psi & & \swarrow \psi_i & & \\
 & & & & U & & 
 \end{array}$$

Then the maps  $\psi$  and  $\psi_i$  define a map (germ)  $f$  of a holomorphic diffeomorphism  $B/G$  to  $U_i/G_i$  at  $0 := \pi(0)$  such that  $f(0) = 0 := \pi_i(0)$ . Then  $f$  induces an isomorphism  $\mathfrak{F}_{V/G}(0) \rightarrow \mathfrak{F}_{V/G_i}(0)$ , it maps  $(B/G)_0$  to  $(\tilde{U}_i/G_i)_0$  and  $f_*(D_{B/G}) = D_{\tilde{U}_i/G_i}$ . Thus by theorem 5.4 there is a germ of a holomorphic diffeomorphism  $F : B \rightarrow \tilde{U}_i$  which is equivariant for a suitable isomorphism  $G \rightarrow G_i$ . ■

**6.7. Corollary.** *Let  $M$  be a complex simply connected manifold,  $G$  a proper discontinuous group of holomorphic diffeomorphisms of  $M$ , and  $\mathfrak{F}_X$  the corresponding sheaf on the orbifold  $X = M/G$ . The  $G$ -manifold  $M$  is a universal covering orbifold for the orbifold  $X$  and it is defined uniquely up to a natural isomorphism of universal coverings by the sheaf  $\mathfrak{F}_X$ , the principal stratum  $X_0$ , and by the reflection divisor  $D_X$ .*

**Proof.** Evidently the manifold  $M$  is a covering orbifold for  $X$ . If  $\tilde{X}$  is a universal covering orbifold for  $X$ , by definition 6.4 there is a cover  $q : \tilde{X} \rightarrow M$ . By definition  $\tilde{X}$  should be a manifold and  $q$  a cover of manifolds. Therefore,  $q$  is a diffeomorphism. Then the statement of the corollary follows from theorem 6.6. ■

An automorphism of the sheaf  $\mathfrak{F}_X$  is called a holomorphic diffeomorphism of the orbit space  $X$ . Theorem 6.5 and corollary 6.7 imply the following analogue of Theorem 5.4.

**6.8. Theorem.** *Let  $M$  be a complex simply connected manifold,  $G$  a proper discontinuous group of holomorphic diffeomorphisms of  $M$ , and  $\mathfrak{F}_X$  the corresponding sheaf on the orbifold  $X = M/G$ . Each holomorphic diffeomorphism  $f$  of the orbit space  $X$  preserving  $X_0$  and  $D_X$  has a holomorphic lift  $F$  to  $M$ , which is  $G$ -equivariant with respect to an automorphism of  $G$ . The lift  $F$  is unique up to composition by an element of  $G$ .*

**Proof.** By theorem 6.6 and corollary 6.7 the manifold  $M$  with the map  $f \circ p : M \rightarrow X$ , where  $p : M \rightarrow X$  is the projection, is a universal covering orbifold for  $X$ . Then there is a holomorphic diffeomorphism  $F : M \rightarrow M$  such that  $p \circ F = f \circ p$ . The equivariance property holds locally by 5.1, thus globally. The lift is uniquely given by choosing  $F(x)$  for a regular point  $x$  in the orbit  $f(p(x))$ . ■

**6.9.** Let  $V$  be a complex vector space with a linear action of a finite group  $G$ . The group  $\mathbb{C}^*$  acts on  $V$  by homotheties and induces an action on  $Z = V/G$ .

**Corollary.** *In this situation, the  $G$ -module  $V$  is uniquely defined up to a linear isomorphism by the sheaf  $\mathfrak{F}_{V/G}$  with the action of  $\mathbb{C}^*$ , by  $Z_0$ , and the reflection divisor  $D_Z$ .* ■

**Proof.** Consider the orbit space  $Z = V/G$  of a  $G$ -module  $V$  with the sheaf  $\mathfrak{F}_{V/G}$ , regular stratum  $Z_0$ , reflection divisor  $D_Z$ , and the action of  $\mathbb{C}^*$  induced by the action of  $\mathbb{C}^*$  on  $V$  by homotheties. Suppose that we have another  $G'$ -module  $V'$  with the same data on  $Z' = V'/G'$  such that there is a biholomorphic map  $f : Z \rightarrow Z'$  preserving these data. By Theorem 4.5 there is a biholomorphic lift  $F : V \rightarrow V'$ , and by lemma 5.1 there is an isomorphism  $a : G \rightarrow G'$  such that  $F \circ g = a(g) \circ F$ . Thus we may assume that  $G = G'$ ,  $V = V'$ ,  $Z = Z'$ , and  $a$  is the identity map. By definition the pullback  $A$  of the vector field on the orbit space  $V/G$  defined by the action of the group  $\mathbb{C}^*$  on  $V/G$  coincides with the vector field on  $V$  defined by the above action of the group  $\mathbb{C}^*$  on  $V$ . By construction  $F^*A = A$  and then the map  $F$  commutes with the action of  $\mathbb{C}^*$  on  $V$ , i.e. for each  $t \in \mathbb{C}^*$  and  $v \in V$  we have  $F(tv) = tF(v)$ . Since  $F$  is biholomorphic it is a linear automorphism of the vector space  $V$ . By definition  $F$  is then an automorphism of the  $G$ -module  $V$ . ■

**6.10. Tensor fields and connections on orbifolds.** The local results in section 3 show that the correct definition of a  $\binom{p}{q}$ -tensor field  $Q$  on an orbifold  $X$  is as follows:  $Q$  is a meromorphic  $\binom{p}{q}$ -tensor field on  $X_1$  such that  $\operatorname{div}_{D_X}(Q) \geq 0$ .

Likewise, we can define connections on orbifolds by requiring the local conditions of section 4.

## References

- [1] E. Bierstone, *Lifting isotopies from orbit spaces*, Topology 14 (1975), 245–252..
- [2] M. Broué, G. Malle, R. Rouquier, *Complex reflection groups, braid groups, Hecke algebras*, J. reine angew. Math. 500 (1998), 127–190..
- [3] M. Losik, *Lifts of diffeomorphisms of orbit spaces for representations of compact Lie groups*, Geom. Dedicata, 88 (2001), 21–36..
- [4] M. Losik, P.W. Michor, V.L. Popov, *Invariant tensor fields and orbit varieties for finite algebraic transformation groups*, arXiv:math.AG/0206008..
- [5] D. Luna, *Slices étales*, Bull. Soc. Math. France, Mémoire 33 (1973), 81–105..
- [6] D. Luna, *Sur certaines opérations différentiables des groupes de Lie*, Amer. J. Math. 97 (1975), 172–181..
- [7] D. Luna, *Fonctions différentiables invariantes sous l'opération d'une groupe réductif*, Ann. Inst. Fourier 26 (1976), 33–49..
- [8] O. V. Lyashko, *Geometry of bifurcation diagrams*, J. Soviet Math. 27 (1984), 2736–2759..
- [9] G.W. Schwarz, *Lifting smooth homotopies of orbit spaces*, Publ. Math. IHES 51 (1980), 37–136..
- [10] L. Solomon, *Invariants of finite reflection groups*, Nagoya J. Math. 22 (1963), 57–64..
- [11] W.P. Thurston, *The geometry and topology of tree-manifolds*, Lect. Notes, Princeton Univ. Press, Princeton (1978).

A. Kriegl: Institut für Mathematik,  
Universität Wien, Strudlhofgasse 4,  
A-1090 Wien, Austria  
Andreas.Kriegl@univie.ac.at

M. Losik: Saratov State University, ul.  
Astrakhanskaya, 83, 410026 Saratov,  
Russia  
LosikMV@info.sgu.ru

P. W. Michor: Institut für Mathe-  
matik, Universität Wien, Strudlhof-  
gasse 4, A-1090 Wien, Austria; *and*:  
Erwin Schrödinger Institut für Math-  
ematische Physik, Boltzmanngasse 9,  
A-1090 Wien, Austria  
Peter.Michor@esi.ac.at

Received  
and in final form