

Double Grothendieck polynomials and colored lattice models

Valentin Buciumas^{*1} and Travis Scrimshaw^{†1}

¹*School of Mathematics and Physics, The University of Queensland, St. Lucia, QLD 4072, Australia*

Abstract. We construct an integrable colored vertex model whose partition function is a double Grothendieck polynomial and relate it to bumpless pipe dreams. This gives a new proof of recent results of Weigandt. For vexillary permutations, we then construct a new model that we call the semidual version model. We use our semidual model and the five-vertex model of Motegi and Sakai to give a new proof that double Grothendieck polynomials for vexillary permutations are equal to flagged factorial Grothendieck polynomials. We then obtain a new proof that the stable limit is a factorial Grothendieck polynomial as defined by McNamara. The states of our semidual model naturally correspond to families of nonintersecting lattice paths, where we can then use the Lindström–Gessel–Viennot lemma to give a determinant formula for double Schubert polynomials corresponding to vexillary permutations.

Keywords: Grothendieck polynomial, colored lattice model, vexillary permutation

1 Introduction

The Yang–Baxter equation, also known as the star–triangle equation, has been the cornerstone of many aspects of modern mathematical physics. It plays a central role in (quantum) integrable systems, which naturally arise across a broad spectrum of mathematics and physics. Applications include explaining combinatorial phenomena and properties, describing probabilistic models and studying Whittaker functions, such as in [5, 3, 8, 2, 9, 15, 22, 19, 26]. In many of the papers cited, one equates an object of interest (like a Hall–Littlewood polynomial or a Whittaker function) with the partition function of a solvable lattice model. This naturally implies interesting properties such as branching rules, Cauchy-type identities, exchange relations, and combinatorial descriptions.

Our work is motivated by the geometry of Schubert varieties. Consider the general linear group $G = \mathrm{GL}(\mathbb{C}^n)$ and the subgroups of lower triangular matrices B and diagonal matrices T . A complete flag is an element in the (complete/full) flag variety G/B and

^{*}valentin.buciumas@gmail.com. V.B. was supported by the Australian Research Council DP180103150.

[†]tcscrim@gmail.com.

corresponds to a sequence of subspaces $\{0\} = V_0 \subseteq \cdots \subseteq V_n = \mathbb{C}^n$ such that $\dim V_i = i$. A Schubert cell is a (left) B -orbit in G/B , and a Schubert variety is the Zariski closure of a Schubert cell. The Schubert varieties are indexed by permutations and have many nice properties, such as Bruhat order corresponding to inclusion and forming a basis for the (T) -equivariant (connective) K-theory ring of the flag variety $K_T^\beta(G/B)$.

In this paper, we connect the equivariant K-theory ring $K_T^\beta(G/B)$ with solvable lattice models. To do this, we use the double (β) -Grothendieck polynomials G_w , where w is a permutation, as representatives for the Schubert varieties in $K_T^\beta(G/B)$ [9, 18]. Our main result (Theorem 3.3) is a colored lattice model whose partition function is (up to a trivial factor of the parameter β) G_w . We construct our model as a translation of the bumpless pipe dreams with a fixed key given in [25], where resolving multiple crossings of two strands precisely corresponds to the colorization performed in [5, 8]. Thus, we encode the permutation into the model by using the coloring. Our proof is showing the Yang–Baxter equation implies the partition function satisfies the same functional equations defining double Grothendieck polynomials (up to a β factor). Therefore, we have a new proof of [25, Thm. 1.1], which gives a formula for G_w as a sum over bumpless pipe dreams with key w proven using a combination of algebraic and combinatorial techniques. Our model is the colored version of the six-vertex model that Lascoux used in [17] to describe G_w using alternating sign matrices. By specializing $\beta = 0$, we recover the formula for double Schubert polynomials from [16]. Our results can be considered as the flag variety version of [8, 22] on the Grassmannian, the set of k -dimensional planes in \mathbb{C}^n .

In the second part of our paper, we consider the analog of $G_w(\mathbf{x}, \mathbf{y}; \beta)$ for the equivariant K-theory of the Grassmannian, which are the factorial Grothendieck polynomials. The stable limit of double Grothendieck polynomials decompose into finitely many factorial Grothendieck polynomials [6, 20]. When w is vexillary (it avoids the pattern 2143), the stable limit of G_w is a single factorial Grothendieck polynomial. Factorial Grothendieck polynomials are given as the sum over set-valued tableaux [20].

We look at what happens to our colored model restricted to vexillary permutations. To do this, we first construct a variation of the uncolored version of our model by swapping $0 \leftrightarrow 1$ on the horizontal lines, which we call the semidual model. To encode the permutation w , we modify the semidual model to not be on a rectangular grid, but instead on a grid given by a partition Λ_w that depends on w . By a key property of vexillary permutations, the partition function is the same as our original colored model and the semidual model becomes a five-vertex model. Hence we can change the weights of one of the vertices by removing a β factor and maintain integrability, and the resulting vertices become those of [22] (up to a gauge transformation). The fact that the shape of the semidual model is Λ_w corresponds to imposing a flagging on set-valued tableaux under the natural bijection between (marked) states of the [22] model given in [21, Sec. 4.2]. Therefore we obtain a new proof that double Grothendieck

polynomials are sums over flagged set-valued tableaux [11] (Theorem 4.4), which was proven using Gröbner geometry. Furthermore, by taking the stable limit, we recover [20] (Theorem 2.1), which was proven by examining the underlying combinatorics. By taking appropriate specializations, we obtain new proofs of results from [6, 22, 26].

A state of the semidual model corresponds to a family of nonintersecting lattice paths, which we then apply the Lindström–Gessel–Viennot lemma to. Yet, we can only obtain a determinant formula for flagged (factorial) Schur functions (Theorem 4.6). We were unable to extend this to the determinant formula for G_w from [1, 19]. Additionally, our NILPs are distinct from those in [13] referenced in [25, Sec. 7.3] and those in [12] and [14].

This extended abstract is organized as follows. In Section 2, we provide the necessary background on double Grothendieck polynomials. In Section 3, we give our integrable colored lattice model whose partition function is a double Grothendieck polynomial and connect it to bumpless pipe dreams. In Section 4, we give our semidual model and relate it to (flagged) set-valued tableaux. This is the extended abstract version of [7].

After [7] was written, we were made aware of [4], where a different lattice model interpretation for double Grothendieck polynomials by using regular pipe dreams. The Boltzmann weights for the states (and the L -matrices) are different as there is no weight preserving bijection between the two (cf. [25, Ex. 6.3]). Their applications are distinct from ours except for Corollary 3.4, which we realized we could prove after seeing [4].

2 Background

Fix a positive integer n . Let $\mathbf{x} = (x_1, \dots, x_n)$ (resp. $\mathbf{y} = (y_1, \dots)$) be a finite (resp. infinite) sequence of indeterminates. For $(\alpha_1, \dots, \alpha_n) \in \mathbb{Z}^n$ of length n , denote $\mathbf{x}^\alpha := x_1^{\alpha_1} \cdots x_n^{\alpha_n}$.

Let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$ be a *partition*, a sequence of weakly decreasing nonnegative integers (of length n). Let $\ell(\lambda) = \max\{k \mid \lambda_k > 0\}$ denote the *length* of λ . The Young diagram (in English convention) of λ is a drawing consisting of stacks of boxes with row i having λ_i boxes pushed into the upper-left corner. The *01-sequence* of λ is given by reading the boundary of λ , starting at the bottom, with horizontal steps being 0 and vertical steps being 1 (we ignore all trailing 0s in the 01-sequence). For example, the 01-sequence of $\lambda = (5, 2, 2, 1, 0, 0)$ for $n = 6$ is 11010110001.

Let S_n be the symmetric group on n elements with $s_i = (i \ i + 1)$. For $w \in S_n$, let $\ell(w)$ denote the length of w . Let w_0 denote the longest element in S_n . The *diagram* of w is

$$D(w) := \{(p, q) \in \{1, \dots, n\} \times \{1, \dots, n\} \mid w(p) > q \text{ and } w^{-1}(q) > p\},$$

(the boxes not in the Rothe diagram of w). Let $w' = 1^k \times w$ denote the permutation

$$w'(i) := \begin{cases} i & \text{if } i \leq k, \\ w(i - k) + k & \text{if } i > k. \end{cases}$$

Let \leq denote the (strong) Bruhat order on S_n . Define $w\mathbf{x} := (x_{w(1)}, \dots, x_{w(n)})$. A permutation w is *vexillary* if it avoids the pattern 2143; that is to say there does not exist $1 \leq i < j < k < \ell \leq n$ such that $w_j < w_i < w_\ell < w_k$.

We can compute λ_w by ordering the sequence whose i -th value is the number of boxes in row i of $D(w)$. We call λ_w the *partition associated to w* . Next, let Λ_w denote the smallest partition whose Young diagram contains the boxes $D(w)$. Following [11, Sec. 5.2], define $F_w = (F_i)_{i=1}^{\ell(\lambda_w)}$ with F_i the row index of the southeastern box of Λ_w that lies on the same diagonal as the last box (i, λ_i) in row i of $\lambda = \lambda_w$. We call F_w the *flagging associated to w* .

Define $x \oplus y := x + y + \beta xy$. Let $f \in \mathbb{Z}[\mathbf{x}]$. The *K-theoretic divided difference operator* is

$$\mathfrak{d}_i f := \frac{(1 + \beta x_{i+1})f - (1 + \beta x_i)s_i f}{x_i - x_{i+1}}.$$

These are normally denoted by π_i in the literature. We also require the *Demazure–Lascoux operator* $\omega_i f := \mathfrak{d}_i(x_i f)$ and the *Demazure–Lascoux atom operator* $\bar{\omega}_i := \omega_i - 1$. For any operator $D_i = \partial_i, \mathfrak{d}_i, \pi_i, \omega_i, \bar{\omega}_i$, define $D_w := D_{i_1} \cdots D_{i_k}$ for any reduced expression $w = s_{i_1} \cdots s_{i_k}$, which is well-defined as these operators all satisfy the braid relations.

Following [9], the *double Grothendieck polynomial* is defined recursively by

$$G_{w_0}(\mathbf{x}, \mathbf{y}; \beta) := \prod_{i+j \leq n} x_i \oplus y_j, \quad G_w(\mathbf{x}, \mathbf{y}; \beta) := \mathfrak{d}_i G_{ws_i}(\mathbf{x}, \mathbf{y}; \beta), \quad \text{for } \ell(ws_i) = \ell(w) + 1.$$

Let SVT_λ denote the set of *set-valued tableaux of shape λ* , fillings of the boxes of λ with non-empty (finite) subsets of $\{1, 2, \dots, n\}$ that satisfy locally

$$\begin{array}{|c|c|} \hline A & B \\ \hline C & \\ \hline \end{array}, \quad \max A \leq \min B, \quad \max A < \min C.$$

For $F = \{F_i \in \mathbb{Z}\}_{i=1}^{\ell(\lambda)}$, let $\text{SVT}_{\lambda, F} \subseteq \text{SVT}_\lambda$ be the subset of *flagged set-valued tableaux*, where every value in the i -th row of T is at most F_i .

Following [11], define the *flagged factorial Grothendieck polynomial* as

$$G_{\lambda, F}(\mathbf{x}|\mathbf{y}; \beta) := \sum_{T \in \text{SVT}_{\lambda, F}} \beta^{|T| - |\lambda|} \prod_{A \in T} \prod_{i \in A} x_i \oplus y_{i+c(A)},$$

where $c(A) = c - r$ is the *content* of the box A (which we have equated with its entry) in row r and column c . A determinant formula for $G_{\lambda, F}(\mathbf{x}, \mathbf{y}; \beta)$ was given in [19].

Theorem 2.1 ([11, Thm. 5.8]). *For a vexillary permutation w , $G_w(\mathbf{x}, \mathbf{y}; \beta) = G_{\lambda_w, F_w}(\mathbf{x}|\mathbf{y}; \beta)$.*

3 Double Grothendieck polynomial colored model

We now give our main result: an integrable colored vertex model whose partition function is $G_w(\mathbf{x}, \mathbf{y}; \beta)$ and relate it to the bumpless pipe dream formula from [25].

a_1	a_2	a_2^\dagger	b_1	b_2	c_1	c_2
$\beta(x \oplus y)$	1	1	1	1	$1 + \beta(x \oplus y)$	1

Figure 1: The Boltzmann weights with $c > c'$ and d being any color for the model \mathfrak{G}_w .

Fix colors $\mathbf{c} = (c_1 > c_2 > \dots > c_n)$ and a permutation $w \in S_n$. Let $w\mathbf{c} = (c_{w(1)}, c_{w(2)}, \dots, c_{w(n)})$ be the natural action of w on the colors. The *(lattice) model* is given by a rectangular grid of n horizontal and vertical lines with each crossing a *vertex* and the lines between two vertices (resp. from one vertex) are *edges* (resp. *half edges*). The *boundary conditions* are the top and left half edges are labeled by 0, the right half edges are labeled by $w\mathbf{c}$ from top-to-bottom, and the bottom half edges begin \mathbf{c} from left-to-right. A *state* is a labeling of the edges by $\{0\} \sqcup \mathbf{c}$, and we call a state *admissible* if each vertex is one of those in Figure 1. For examples of admissible states, see Figure 3.

We assign a non-zero *(Boltzmann) weight* with *spectral parameter* x to each of the vertices in Figure 1, the set of which is called an *L-matrix*. The *(Boltzmann) weight* $\text{wt}(S)$ of a state S is the product of all of the Boltzmann weights of all vertices. Let \mathfrak{G}_w denote the set of all possible admissible states for this model, which we call the *double Grothendieck model*. The *partition function* is $Z(\mathfrak{G}_w; \mathbf{x}, \mathbf{y}; \beta) := \sum_{S \in \mathfrak{G}_w} \text{wt}(S)$.

Proposition 3.1. *The colored vertex model \mathfrak{G}_w with R-matrix given by Figure 2 is integrable: it satisfies the RLL form of the Yang–Baxter equation, where the partition function of the following two models are equal for any boundary conditions $a, b, c, d, e, f \in \{0, c_1, c_2, \dots, c_n\}$:*

(3.1)

The Yang–Baxter relation only involves at most 3 colors, so Proposition 3.1 is an identity of $2^4 \times 2^4$ matrices as colors are conserved under the R-matrix and L-matrix. Thus Proposition 3.1 can be shown by direct computation using, e.g., SAGEMATH [23].





Figure 2: The colored R -matrix with $c > c'$ and d being any color for the double Grothendieck model. Note that the weights are not symmetric with respect to color.

Following [5, 8], we use the Yang–Baxter equation and the train argument to construct a functional equation for the partition function of our model \mathfrak{G}_w . The proof is standard and is the same as the one given in, e.g., [8, Lemma 3.3]; pictorially it is the equality of:



Lemma 3.2. Let $w \in S_n$, and consider any s_i such that $ws_i > w$. Then we have

$$\beta Z(\mathfrak{G}_w; \mathbf{x}, \mathbf{y}; \beta) = \frac{(1 + \beta x_{i+1}) \cdot Z(\mathfrak{G}_{ws_i}; \mathbf{x}, \mathbf{y}; \beta) - (1 + \beta x_i) \cdot Z(\mathfrak{G}_{ws_i}; s_i \mathbf{x}, \mathbf{y}; \beta)}{x_i - x_{i+1}}.$$

Theorem 3.3. We have $Z(\mathfrak{G}_w; \mathbf{x}, \mathbf{y}; \beta) = \beta^{\ell(w)} G_w(\mathbf{x}, \mathbf{y}; \beta)$.

The authors thank Anatol Kirillov for noting the following well-known symmetry can be seen from [9] or from $G_w(\mathbf{x}, \mathbf{y}; \beta)$ as a sum over usual pipe dreams (or RC-graphs) [9, 10]. A colored lattice model proof was first given in [4, Prop. 9.1].

Corollary 3.4. We have $Z(\mathfrak{G}_w; \mathbf{x}, \mathbf{y}; \beta) = Z(\mathfrak{G}_{w^{-1}}; \mathbf{y}, \mathbf{x}; \beta)$ and $G_w(\mathbf{x}, \mathbf{y}; \beta) = G_{w^{-1}}(\mathbf{y}, \mathbf{x}; \beta)$.

In [25, Eq. (3.7)], each uncolored vertex corresponds to one of the tiles that defines a bumpless pipe dream. The key ∂P of a bumpless pipe dream P is given by the Demazure




Figure 3: The unique state for the colored model \mathfrak{G}_{w_0} (left), a state in $\mathfrak{G}_{s_1 s_3}$ (center), and a state in $\mathfrak{G}_{s_2 s_3 s_2}$ (right) with $n = 4$ and $c_1 > c_2 > c_3 > c_4$.

product of the simple transpositions (where instead $s_i^2 = s_i$). Indeed, color each line of P starting from the bottom as c_1, \dots, c_n from left-to-right where each color must cross (*i.e.*, the vertex a_2^\dagger or with the colors swapped). Moving from the bottom-left to the top-right along diagonals, when we replace a color flipped a_2^\dagger with a_2 (see also [25, Eq. (2.4), Lemma 2.1]). Thus states of \mathfrak{G}_w are bumpless pipe dreams with key w after forgetting the colors, and we obtain a new proof of [25, Thm. 1.1].

4 The semidual vertex model

In this section, we modify our model and remove the colors in the case w is a vexillary permutation. We draw the states using tiles. Thus the L -matrix given in Figure 1 becomes



By [25, Lemma 7.2], a bumpless pipe dream state has no a_2 vertex if and only if w is vexillary. Thus we set the corresponding Boltzmann weight to 0. We can forget the colors when drawing the lattice models, but it does not quite equate the colored model with the uncolored model as we still need to keep track of which strands cross to encode w . This is important as the model needs to remain colored in order to get the correct partition function corresponding to w . Let \mathfrak{G} denote the corresponding uncolored model.

We define the *semidual model* \mathfrak{D} to be the colored lattice model on an $n \times n$ grid using the L -matrix given in Figure 4 and boundary conditions as follows. The left (resp.

\bar{a}_1	\bar{a}_2	\bar{b}_1	\bar{b}_2	\bar{c}_1	\bar{c}_2
1	1	1	$\beta(x \oplus y)$	$1 + \beta(x \oplus y)$	1

Figure 4: Boltzmann weights of the semidual model.


bottom) boundary condition is the colors $c_n > \dots > c_1$ from top-to-bottom (resp. left-to-right); the top and right boundary edges are all 0. Since no colors cross, we forget the coloring, but unlike before, we obtain an equivalent model with no dependency on w .

Proposition 4.1. *There exists a bijection $\Phi: \mathfrak{G} \rightarrow \mathfrak{D}$ given by*

$$\square \mapsto \begin{array}{|c|} \hline \text{---} \\ \hline \end{array}, \quad \begin{array}{|c|} \hline | \\ \hline \end{array} \mapsto \begin{array}{|c|} \hline \diagdown \\ \hline \end{array}, \quad \begin{array}{|c|} \hline \text{└} \\ \hline \end{array} \mapsto \begin{array}{|c|} \hline \diagdown \\ \hline \end{array}, \quad \begin{array}{|c|} \hline \text{├} \\ \hline \end{array} \mapsto \begin{array}{|c|} \hline | \\ \hline \end{array}, \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \mapsto \square, \quad \begin{array}{|c|} \hline \text{└} \\ \hline \end{array} \mapsto \begin{array}{|c|} \hline \diagdown \\ \hline \end{array}.$$

We note that the map in Proposition 4.1 is defined by interchanging $0 \leftrightarrow 1$ on the horizontal (auxiliary space) component between the models.

Example 4.2. We apply the bijection defined in Proposition 4.1 to Figure 3 (right):



Now we want to reintroduce the vexillary permutation w into these uncolored models. If we restrict our model to Λ_w (as opposed to an $n \times n$ grid), we can use [25, Lemma 7.2] (see also [11, Cor. 3.3]) to see that there are no vertices of type a_2 nor a_2^\dagger in \mathfrak{G}_w . Therefore we have an uncolored five-vertex model whose partition function is $\beta^{\ell(w)} G_w(\mathbf{x}, \mathbf{y}; \beta)$. Equivalently for the corresponding semidual model on Λ_w there are no vertical lines, and the states can be thought of as a family of nonintersecting lattice paths. (These are also the tiles in [26, Eq. (7)] rotated 180 degrees and reflected across the vertical axis with $z \mapsto 1 - (x \oplus y)$ at $\beta = -1$.) In terms of the partition function, all this does is remove the $\beta^{\ell(w)}$ as only the weight of vertex \bar{b}_2 changes by removing the β , of which there are precisely $\ell(w)$ such vertices.

Proposition 4.3. *The model \mathfrak{D}_w is integrable with $\beta^{\ell(w)} Z(\mathfrak{D}_w; \mathbf{x}, \mathbf{y}; \beta) = Z(\mathfrak{G}_w; \mathbf{x}, \mathbf{y}; \beta)$.*

Let $F_w := (F_i)_{i=1}^k$ denote the heights of the right endpoints of the paths in Λ_w that end on the right boundary. So the tile immediately to the right of such a boundary point is a  not in Λ_w . Any other path that does not have such a right endpoint must simply move diagonally. The sequence F_w is exactly the flagging associated to w .


Now if we rotate the semidual model \mathfrak{D}_w by 180 degrees and extend from the endpoints in the (rotated) shape Λ_w by diagonal lines (which go to the northwest), we obtain precisely a state in the uncolored model \mathfrak{S}_{λ_w} (see also [22]). Thus we can apply the natural bijection Θ between marked states, where we allow only tiles  to be marked, and set-valued tableaux via marked Gelfand–Tsetlin patterns from [8, Sec. 2.3]. However, the flagging F_w imposes a restriction on the possible states and set-valued tableaux. It is straightforward to see that this precisely corresponds to restricting to the set of flagged set-valued tableaux $\text{SVT}_{\lambda_w, F_w}$. Hence, we have a new proof of Theorem 2.1.

As in [11, Cor. 5.9], this gives an integrable systems proof of [6, Thm 3.1]. The bijection Θ is also the same as the bijection from pipe dreams to $\text{SVT}_{\lambda_w, F_w}$ given in [11, Prop. 5.3], which is different from the formula in [9] (see, e.g., [11, Ex. 5.10]).

We give new a proof of [20, Thm. 8.7] and [22, 26] by taking the stable limit of the semidual model $\mathfrak{D}_{1^k \times w}$ as $k \rightarrow \infty$ and restricting to a finite number of variables.

Theorem 4.4 ([22, 26]). *The partition function of the stable limit model equals $G_{\lambda_w}(\mathbf{x}|\mathbf{y}; \beta)$.*

Example 4.5. We consider $\lambda = (1)$ and $n = 2$. We consider $w = [2, 1]$ and $1^2 \times w$:




where we have drawn the \mathfrak{S}_w states on the left and the semidual states on the right.

We also note that the symmetry used in [25, Lemma 8.1] is precisely the natural symmetry from reflecting the semidual model along the $y = x$ line. Furthermore, the bijection in [25, Lemma 8.2] comes from the reflected elementary excitations/emissions, where we instead consider the paths in Λ_w (which in this case equals λ_w) as being fixed.

Another benefit of the semidual model is being able to interpret a state as a family of nonintersecting lattice paths (NILPs). The *Lindström–Gessel–Viennot (LGV) lemma* posits that the sum of the weights over all NILPs in a edge-weighted directed graph can be given as a determinant of the matrix $[p_{ab}]_{a,b=1}^n$, where $p_{a,b}$ is the sum over each path from the a -th starting point to the b -th ending point and taking the product of the edge weights. Thus, we want to construct a weight preserving bijection from a marked state of the semidual model into a NILP to express the partition function as a determinant.

We use the following local translation from tiles to a directed graph:



with a non-trivial edge weight on only the second horizontal step in each tile that depends on the position of the tile. We call the blue and gold edges *Schubert edges* and the red edges *K-theory edges*. All edges have weight 1 except for the Schubert edges in gold, which have weight $w(i, j)$ that we will give later. We can restrict to the paths that are not simply diagonals (*i.e.*, have at least one horizontal step).

We first consider what happens when we apply the LGV lemma naïvely using the tiles. In this case, we see that we must have $\beta = 0$, and so we take $w(i, j) = x_i \oplus y_j$, remove the K-theory edges, and only consider unmarked states. Let h_b denote the height of the b -th endpoint and $\lambda = \lambda_w$. One can see that $p_{ab} = G_{(\lambda_b + a - b)}(x_a, \dots, x_{h_b} | \mathbf{y}; 0)$. Thus, we have the following expression for double Schubert polynomials by the LGV lemma.

Theorem 4.6. *If w is vexillary, then $Z(\mathfrak{D}_w; \mathbf{x}, \mathbf{y}; 0) = \det [p_{ab}]_{a,b=1}^n = G_w(\mathbf{x}, \mathbf{y}; 0)$.*

Our formula at $\mathbf{y} = 0$ likely differs from [24] by some sequence of row operations.

Example 4.7. Consider the permutation $w = s_2 s_3 s_2$ (for an example of a state in \mathfrak{D}_w , see Example 4.2). We compute $\lambda_w = (2, 1)$ and $F_w = (2, 3)$. Applying Theorem 4.6,

$$\begin{aligned} Z(\mathfrak{D}_w; \mathbf{x}, \mathbf{y}; 0) &= \det \begin{bmatrix} G_{(2)}(x_1, x_2 | \mathbf{y}; 0) & 1 & 0 & 0 \\ G_{(3)}(x_2 | \mathbf{y}; 0) & G_{(1)}(x_2, x_3 | \mathbf{y}; 0) & 0 & 0 \\ 0 & G_{(2)}(x_3 | \mathbf{y}; 0) & 1 & 0 \\ 0 & 0 & G_{(1)}(x_4 | \mathbf{y}; 0) & 1 \end{bmatrix} \\ &= \det \begin{bmatrix} G_{(2)}(x_1, x_2 | \mathbf{y}; 0) & 1 \\ \prod_{j=1}^3 x_2 \oplus y_j & x_2 \oplus y_1 + x_3 \oplus y_2 \end{bmatrix}, \end{aligned}$$

where $G_{(2)}(x_1, x_2 | \mathbf{y}; 0) = (x_1 \oplus y_1)(x_1 \oplus y_2) + (x_1 \oplus y_1)(x_2 \oplus y_3) + (x_2 \oplus y_2)(x_2 \oplus y_3)$. Note that going from the 4×4 determinant to the 2×2 determinant comes from the fact that two of the paths contain no horizontal steps. Hence, we have

$$\begin{aligned} Z(\mathfrak{D}_w; \mathbf{x}, \mathbf{y}; 0) &= (x_1 \oplus y_1)(x_1 \oplus y_2)(x_2 \oplus y_1) + (x_1 \oplus y_1)(x_2 \oplus y_3)(x_2 \oplus y_1) \\ &\quad + (x_1 \oplus y_1)(x_1 \oplus y_2)(x_3 \oplus y_2) + (x_1 \oplus y_1)(x_2 \oplus y_3)(x_3 \oplus y_2) \\ &\quad + (x_2 \oplus y_2)(x_2 \oplus y_3)(x_3 \oplus y_2) = G_{\lambda_w, F_w}(x_1, x_2 | \mathbf{y}; 0), \end{aligned}$$

with corresponding flagged set-valued tableaux ($\beta = 0$ gives semistandard tableaux)

$$\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline 2 & 2 \\ \hline 3 & \\ \hline \end{array}.$$

If we compare this with the formula from [24, Thm. 1.3], we have

$$\begin{aligned} G_{\lambda_w, F_w}(\mathbf{x}|0;0) &= \det \begin{bmatrix} x_1^2 + x_1x_2 + x_2^2 & 1 \\ x_1^3 + x_1^2x_2 + x_1x_2^2 + x_2^3 & x_1 + x_2 + x_3 \end{bmatrix}, \\ &= x_1^2x_2 + x_1x_2^2 + x_1^2x_3 + x_1x_2x_3 + x_2^2x_3. \end{aligned}$$

Note that subtracting x_1 times the first row from the second in the above matrix is precisely the 2×2 determinant used to compute $Z(\mathfrak{D}_w; \mathbf{x}, 0; 0)$.

If instead we want generic β , then we take $w(i, j) = \beta(x_i \oplus y_j)$ to get a relation with the L -matrix from Figure 4. However, it is impossible for the NILPs to agree if marking a  tile corresponds to traveling along the K-theory edge as we can now also use .

Acknowledgments

The authors thank A. Weigandt, K. Motegi, K. Sakai, and P. Zinn-Justin for useful discussions and the authors of [4] for sharing their preprint.

References

- [1] D. Anderson. “K-theoretic Chern class formulas for vexillary degeneracy loci”. *Adv. Math.* **350** (2019), pp. 440–485. [DOI](#).
- [2] A. Borodin and M. Wheeler. “Nonsymmetric Macdonald polynomials via integrable vertex models”. 2019. [arXiv:1904.06804](#).
- [3] B. Brubaker, V. Buciumas, D. Bump, and H. P. A. Gustafsson. “Colored vertex models and Iwahori Whittaker functions”. 2019. [arXiv:1906.04140](#).
- [4] B. Brubaker, C. Frechette, A. Hardt, E. Tibor, and K. Weber. “Frozen pipes: Lattice models for Grothendieck polynomials”. 2020. [arXiv:2007.04310](#).
- [5] B. Brubaker, V. Buciumas, D. Bump, and H. P. A. Gustafsson. “Colored five-vertex models and Demazure atoms”. *J. Combin. Theory Ser. A* **178** (2021), p. 105354. [DOI](#).
- [6] A. S. Buch. “A Littlewood-Richardson rule for the K-theory of Grassmannians”. *Acta Math.* **189.1** (2002), pp. 37–78. [DOI](#).
- [7] V. Buciumas and T. Scrimshaw. “Double Grothendieck polynomials and colored lattice models”. *Int. Math. Res. Not. IMRN* (2020). [DOI](#).
- [8] V. Buciumas, T. Scrimshaw, and K. Weber. “Colored five-vertex models and Lascoux polynomials and atoms”. *J. Lond. Math. Soc.* **102.3** (2020), pp. 1047–1066. [DOI](#).
- [9] S. Fomin and A. N. Kirillov. “Grothendieck polynomials and the Yang-Baxter equation”. *Formal power series and algebraic combinatorics/Séries formelles et combinatoire algébrique*. DIMACS, Piscataway, NJ, 1994, pp. 183–189.

- [10] A. Knutson and E. Miller. “Subword complexes in Coxeter groups”. *Adv. Math.* **184.1** (2004), pp. 161–176. [DOI](#).
- [11] A. Knutson, E. Miller, and A. Yong. “Gröbner geometry of vertex decompositions and of flagged tableaux”. *J. Reine Angew. Math.* **630** (2009), pp. 1–31. [DOI](#).
- [12] C. Krattenthaler. “On multiplicities of points on Schubert varieties in Grassmannians. II”. *J. Algebraic Combin.* **22.3** (2005), pp. 273–288. [DOI](#).
- [13] V. Kreiman. “Schubert classes in the equivariant K -theory and equivariant cohomology of the Grassmannian”. 2005. [arXiv:math/0512204](#).
- [14] V. Kreiman and V. Lakshmibai. “Multiplicities of singular points in Schubert varieties of Grassmannians”. *Algebra, arithmetic and geometry with applications (West Lafayette, IN, 2000)*. Springer, Berlin, 2004, pp. 553–563.
- [15] G. Kuperberg. “Another proof of the alternating-sign matrix conjecture”. *Int. Math. Res. Not. IMRN* **3** (1996), pp. 139–150. [DOI](#).
- [16] T. Lam, S. J. Lee, and M. Shimozono. “Back stable Schubert calculus”. *Compos. Math.* (2020). To appear. [arXiv:1806.11233](#).
- [17] A. Lascoux. “Chern and Yang through ice”. Preprint. 2002.
- [18] A. Lascoux and M.-P. Schützenberger. “Structure de Hopf de l’A.au de cohomologie et de l’A.au de Grothendieck d’une variété de drapeaux”. *C. R. Acad. Sci. Paris Sér. I Math.* **295.11** (1982), pp. 629–633.
- [19] T. Matsumura and S. Sugimoto. “Factorial flagged Grothendieck polynomials”. *Proceedings of the conference “An international festival in Schubert Calculus” held in Guangzhou in November 2017*. Springer Proceedings in Mathematics & Statistics. To appear. Springer, New York, 2020. [arXiv:1903.02169](#).
- [20] P. J. McNamara. “Factorial Grothendieck polynomials”. *Electron. J. Combin.* **13.1** (2006), Research Paper 71, 40. [Link](#).
- [21] C. Monical, O. Pechenik, and T. Scrimshaw. “Crystal structures for symmetric Grothendieck polynomials”. *Transform. Groups* (2020). To appear. [DOI](#).
- [22] K. Motegi and K. Sakai. “Vertex models, TASEP and Grothendieck polynomials”. *J. Phys. A* **46.35** (2013), pp. 355201, 26. [DOI](#).
- [23] *Sage Mathematics Software (Version 9.1)*. <https://www.sagemath.org>. The Sage Developers. 2020.
- [24] M. L. Wachs. “Flagged Schur functions, Schubert polynomials, and symmetrizing operators”. *J. Combin. Theory Ser. A* **40.2** (1985), pp. 276–289. [DOI](#).
- [25] A. Weigandt. “Bumpless pipe dreams and alternating sign matrices”. 2020. [arXiv:2003.07342](#).
- [26] M. Wheeler and P. Zinn-Justin. “Littlewood-Richardson coefficients for Grothendieck polynomials from integrability”. *J. Reine Angew. Math.* **757** (2019), pp. 159–195. [DOI](#).