

# Integrality in the Matching-Jack Conjecture and the Farahat–Higman Algebra

Houcine Ben Dali\*<sup>1,2</sup>

<sup>1</sup>Université de Lorraine, CNRS, IECL, F-54000 Nancy, France

<sup>2</sup>Université de Paris, CNRS, IRIF, F-75006, Paris, France

**Abstract.** Using Jack polynomials, Goulden and Jackson have introduced a one parameter deformation  $\tau_b$  of the generating series of bipartite maps. The Matching-Jack conjecture suggests that the coefficients  $c_{\mu,\nu}^\lambda$  of the function  $\tau_b$  in the power-sum basis are non-negative integer polynomials in the deformation parameter  $b$ . Dołęga and Féray have proved in 2016 the “polynomiality” part in the Matching-Jack conjecture. In this paper, we prove the “integrality” part.

The proof is based on a recent work of the author that deduces the Matching-Jack conjecture for marginal sums  $c_{\mu,l}^\lambda$  from an analog result for the  $b$ -conjecture, established in 2020 by Chapuy and Dołęga. A key step in the proof involves a new connection with the graded Farahat–Higman algebra.

**Keywords:** Matching-Jack conjecture, Jack polynomials, Farahat–Higman algebra, top coefficients

## 1 Introduction

### 1.1 Coefficients $c_{\mu,\nu}^\lambda$ and the Matching-Jack conjecture

Jack polynomials  $J_\theta^{(\alpha)}$  are symmetric functions that depend on a deformation parameter  $\alpha$  (see [14, 21, 19]). We consider the power-sum functions  $\mathbf{p} := (p_1, p_2, \dots)$ ,  $\mathbf{q} := (q_1, q_2, \dots)$  and  $\mathbf{r} := (r_1, r_2, \dots)$  associated to three different alphabets, and we denote by  $J_\theta^{(\alpha)}$  the Jack polynomial of parameter  $\alpha$  expressed in the power-sum basis. In [11], Goulden and Jackson have introduced the function  $\tau_b$  that depends on the parameter  $b := \alpha - 1$  and defined by:

$$\tau_b(t, \mathbf{p}, \mathbf{q}, \mathbf{r}) := \sum_{n \geq 0} t^n \sum_{\theta \vdash n} \frac{1}{j_\theta^{(\alpha)}} J_\theta^{(\alpha)}(\mathbf{p}) J_\theta^{(\alpha)}(\mathbf{q}) J_\theta^{(\alpha)}(\mathbf{r}),$$

where  $j_\theta^{(\alpha)}$  is the square norm of the Jack polynomial  $J_\theta^{(\alpha)}$  with the respect to an  $\alpha$ -deformation of the Hall scalar product.

---

\*[houcine.ben-dali@univ-lorraine.fr](mailto:houcine.ben-dali@univ-lorraine.fr)

The main motivation of studying this function comes from the fact that it gives for  $b = 0$  and  $b = 1$  respectively, the generating series of bipartite maps on oriented and general surfaces (see [12]). Maps are graphs embedded into surfaces, orientable or not. The enumeration of maps involves various methods such as matrix integrals, representation theory tools and bijective methods [4, 8, 18].

We consider the coefficients  $c_{\mu,\nu}^\lambda(b)$  defined by

$$\tau_b(t, \mathbf{p}, \mathbf{q}, \mathbf{r}) = \sum_{n \geq 0} t^n \sum_{\lambda, \mu, \nu \vdash n} \frac{c_{\mu,\nu}^\lambda(b)}{z_\lambda (1+b)^{\ell(\lambda)}} p_\lambda q_\mu r_\nu. \quad (1.1)$$

These coefficients are the main objects of the Matching-Jack conjecture, formulated by Goulden and Jackson [11, Conjecture 3.5].

**The Matching-Jack conjecture** (Goulden and Jackson). *For every partitions  $\lambda, \mu, \nu$  of size  $n \geq 1$ , the coefficient  $c_{\mu,\nu}^\lambda$  is a polynomial in  $b$  with non-negative integer coefficients.*

The Matching-Jack conjecture is equivalent to saying that  $\tau_b$  is a generating series of matchings with some particular weights (see [11, Conjecture 4.2]). There exists a connected version of this conjecture called the  $b$ -conjecture. The  $b$ -conjecture suggests that the function  $(1+b) \frac{\partial}{\partial t} \log(\tau_b)$  has also a positivity property and that it enumerates bipartite maps with some weights [11, Conjectures 6.2 and 6.3]. Since bipartite maps can be encoded with matchings (see, e.g., [2, 7, 12]), the Matching-Jack conjecture and the  $b$ -conjecture are related. However, no implication between them has been proved and both of them are still open.

## 1.2 Former results and main theorem

In addition to the special cases  $b = 0$  and  $b = 1$  that follow from connections with representation theory (see [18, 11]), several partial results related to the Matching-Jack conjecture have been established, we cite here some of them.

The first type of results is related to coefficients  $c_{\mu,\nu}^\lambda$  for general partitions  $\lambda, \mu, \nu$ . It follows from the Jack polynomials theory that these coefficients are rational functions in  $b$ , the polynomiality has been proved by Féray and Dołęga.

**Theorem 1.1** ([6, Corollary 4.2]). *For all partitions  $\lambda, \mu, \nu \vdash n \geq 1$ , the coefficient  $c_{\mu,\nu}^\lambda$  is polynomial in  $b$  with rational coefficients.*

Other properties about the structure of these polynomials have been proved in [17].

The second type of results gives the Matching-Jack conjecture for some family of partitions  $(\lambda, \mu, \nu)$  and specializations of the function  $\tau_b$ . The following has been established in [2], it has been deduced from an analogous result for the  $b$ -conjecture proved by Chapuy and Dołęga in [5].

**Theorem 1.2** ([2]). *For every  $n, l \geq 1$  and for every partitions  $\lambda, \mu \vdash n$ , the marginal coefficient  $c_{\mu, l}^\lambda$  defined by*

$$c_{\mu, l}^\lambda := \sum_{\ell(v)=l} c_{\mu, v}^\lambda,$$

*is a polynomial in  $b$  with non-negative integer coefficients.*

This previous theorem covers other partial results in this direction (see [16, 15]). The main result of this paper gives integrality in the Matching-Jack conjecture.

**Theorem 1.3.** *For every  $\lambda, \mu, \nu \vdash n \geq 1$ , the coefficient  $c_{\mu, \nu}^\lambda$  is a polynomial in  $b$  with integer coefficients.*

The proof of this theorem is based on [Theorem 1.2](#). Since the approach used here is independent from the one considered in [6], it gives a new proof of [Theorem 1.1](#).

### 1.3 The Farahat–Higman algebra

The Farahat–Higman algebra was introduced in [9] in order to study the structure coefficients of the conjugacy classes  $C_\mu(n)$  in the center of the symmetric group algebra  $Z(\mathbb{Z}\mathfrak{S}_n)$ . It has been shown that the Farahat–Higman algebra is isomorphic to the algebra of integral symmetric functions (see [10, 22]), it is also related to the algebra of partial permutations introduced by Ivanov and Kerov in [13].

In [Section 4](#), we will consider a graded version of the Farahat–Higman algebra that we denote  $\mathcal{Z}_\infty$  and that has been introduced in [19, Example 24, page 131]. In [Theorem 4.4](#), we exhibit a new basis of  $\mathcal{Z}_\infty$ , which is useful in the proof of the main theorem.

### 1.4 Steps of the proof

A key tool of the proof of [Theorem 1.3](#) is the following multiplicativity property that can be obtained using the orthogonality of the Jack polynomials (see [3] for more details).

**Proposition 1.4.** *For every  $\lambda, \mu, \nu \vdash n \geq 1$  and  $l \geq 1$ , we have*

$$\sum_{\kappa \vdash n} c_{\mu, \kappa}^\lambda c_{\nu, l}^\kappa = \sum_{\theta \vdash n} c_{\theta, l}^\lambda c_{\mu, \nu}^\theta. \quad (1.2)$$

This property will be considered as a system of linear equations that allows to recover  $c_{\mu, \nu}^\lambda$  from the coefficients  $c_{\mu, l}^\lambda$  (see [Remark 1](#)). We now give the key steps of the proof of [Theorem 1.3](#).

- We prove that for a particular choice of set of parameters  $\lambda, \mu, \nu, l$ , [Equation \(1.2\)](#) gives a square linear system with some triangularity property that allows to obtain information on  $c_{\mu, \nu}^\lambda$  by induction on the length of  $\nu$  (see [Lemma 3.1](#)).

- We prove that the diagonal blocks of the matrix encoding this system, denoted  $Q^{(r)}$ , contain some coefficients  $t_\pi^\rho$ , that are independent from  $b$  (see [Section 2](#) and [Proposition 3.2](#)).
- We prove that the matrix  $Q^{(r)}$  is invertible in  $\mathbb{Z}$  by proving that it is a change-of-basis matrix in the graded Farahat–Higman algebra  $\mathcal{Z}_\infty$  (see [Proposition 4.3](#) and [Theorem 4.4](#)).

Using [Remark 1](#) below and the combinatorial interpretation of the coefficients  $c_{\mu,l}^\lambda$  given in [\[2\]](#), it is possible to give a new proof for the Matching-Jack conjecture for  $b = 0$  and  $b = 1$  that does not use representation theory (such a proof follows in a more intricate way from [\[5\]](#), private communication). Unfortunately, the non-negativity of the coefficients of  $c_{\mu,\nu}^\lambda$  as polynomials in  $b$  seems to be out of reach with our approach. We were also not able to use the same arguments to obtain integrality in the  $b$ -conjecture. However, it is possible to obtain the integrality for the cumulants of  $c_{\mu,\nu}^\lambda$ , which up to rescaling by a factor of the form  $\frac{n}{z_\lambda(1+b)^{\ell(\lambda)-1}}$ , give the coefficients of the  $b$ -conjecture.

## 1.5 Some notations

An *integer partition*  $\lambda = [\lambda_1, \lambda_2, \dots, \lambda_l]$  is a sequence of weakly decreasing positive integers  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_l > 0$ . The integer  $l$  is the *length* of the partition  $\lambda$ , *i.e.* its number of parts. It is denoted by  $\ell(\lambda)$ . We define the *rank* of a partition  $\lambda$  by  $\text{rk}(\lambda) := n - \ell(\lambda)$ . We denote by  $\lambda - \mathbf{1}$  the partition  $[\lambda_1 - 1, \lambda_2 - 1, \dots, \lambda_l - 1]$ , and if  $r \geq l$ , we denote by  $\lambda \oplus 1^r$  the partition  $[\lambda_1 + 1, \dots, \lambda_l + 1, \underbrace{1, 1, \dots, 1}_{r-l \text{ times}}]$ .

## 2 Top coefficients $t_\pi^\rho$

As announced in the introduction, the proof of [Theorem 1.3](#) involves the resolution of a linear system satisfied by the coefficients  $c_{\mu,\nu}^\lambda$ . In this section, we introduce the matrices  $Q^{(r)}$  that encode this system. We start by the following lemma, which is a consequence of [Theorem 1.2](#) and some basic properties of the coefficients  $c_{\mu,l}^\lambda$  (see [\[3\]](#) for the proof).

**Lemma 2.1.** *For every partitions  $\lambda, \mu, \vdash n \geq 1$  and  $l \geq 1$ , we have the following bound on the degree of  $c_{\mu,l}^\lambda$  as a polynomial in  $b$ :*

$$\deg_b(c_{\mu,l}^\lambda) \leq n - l + \ell(\lambda) - \ell(\mu).$$

We consider some stability properties of the coefficients  $c_{\mu,l}^\lambda$  for which the previous bound is zero.

**Proposition 2.2.** *Let  $\kappa, \nu, \mu \vdash n \geq 1$ , such that  $\text{rk}(\kappa) = \text{rk}(\nu) + \text{rk}(\mu)$ . We have*

$$c_{\nu, \mu}^{\kappa} = c_{\nu \cup 1^m, \mu \cup 1^m}^{\kappa \cup 1^m}, \quad \text{for every } m \geq 1.$$

*In particular, for every  $\kappa, \nu, \mu \vdash n \geq 1$ , and  $l \geq 1$  such that  $\text{rk}(\kappa) = \text{rk}(\nu) + n - l$ , we have*

$$c_{\nu, l}^{\kappa} = c_{\nu \cup 1^m, l+m}^{\kappa \cup 1^m}, \quad \text{for every } m \geq 1.$$

*Proof.* This is a consequence of [Proposition 4.1](#) and [Equation \(4.2\)](#) below. □

We introduce the following definition.

**Definition 2.1.** Let  $\rho$  and  $\pi$  be two partitions of size  $r \geq 1$ . We consider two partitions  $\kappa$  and  $\nu$  of the same size  $n \geq r + \ell(\rho)$ , such that  $\rho = \kappa - \mathbf{1}$  (or equivalently  $\kappa = \rho \oplus 1^{n-r}$ ),  $\nu = \pi \cup 1^{n-r}$  and  $l$  is such that  $n - l + \text{rk}(\nu) = r$ . We define the *top coefficient*<sup>1</sup>  $t_{\pi}^{\rho} := c_{\nu, l}^{\kappa}$ . Note that given [Proposition 2.2](#) this definition does not depend on  $n$ .

We consider the matrix of top coefficients  $Q^{(r)} := (t_{\pi}^{\rho})_{\pi, \rho \vdash r}$ . We give here  $Q^{(r)}$  for  $r = 3$ ,

$\pi \setminus \rho$	[3]	[2, 1]	[1 <sup>3</sup> ]
[3]	4	1	0
[2, 1]	6	4	3
[1 <sup>3</sup> ]	1	1	1

The following theorem will be proved in [Section 4](#) (see [Theorem 4.4](#)).

**Theorem 2.3.** *The matrix  $Q^{(r)} = (t_{\mu}^{\rho})_{\rho, \mu \vdash r}$  is invertible in  $\mathbb{Z}$  for every  $r \geq 1$ .*

There exists an explicit expression of the top coefficients  $t_{\pi}^{\rho}$ , see [\[1\]](#). However, for the proof of [Theorem 2.3](#), it will be more natural to consider the algebraic definition of these coefficients and see the matrix  $Q^{(r)}$  as a change-of-basis matrix (see [Proposition 4.3](#)).

### 3 Proof of [Theorem 1.3](#)

The main purpose of this section is to prove that [Theorem 2.3](#) implies [Theorem 1.3](#).

We fix  $n > 0$ . For  $1 \leq r < n$ , we introduce the assertion  $\mathcal{A}_n^{(r)}$ :

$\mathcal{A}_n^{(r)}$  : for every  $\lambda, \mu, \kappa \vdash n$  such that  $\text{rk}(\kappa) = r$ ,  $c_{\mu, \kappa}^{\lambda}$  is an integer polynomial in  $b$ .

We will prove  $\mathcal{A}_n^{(r)}$  by induction on  $r$ .

---

<sup>1</sup>This terminology will be justified in [Section 4](#)

**Lemma 3.1.** *We fix  $1 \leq r < n$ , and we assume that the assertions  $\mathcal{A}_n^{(i)}$  hold for  $i < r$ . Let  $\lambda, \mu \vdash n$  and let  $(\nu, l)$  be a pair satisfying the condition*

$$\nu \vdash n, \text{rk}(\nu) < r \quad \text{and} \quad n - l + \text{rk}(\nu) = r. \quad (3.1)$$

*Then we have that*

$$P_{\lambda, \mu, \nu, l}^{(r)} := \sum_{\text{rk}(\kappa)=r} c_{\mu, \kappa}^{\lambda} c_{\nu, l}^{\kappa} \quad (3.2)$$

*is an integer polynomial in  $b$ .*

*Proof.* Note that with the conditions of the proposition, the right hand-side  $\sum_{\theta \vdash n} c_{\theta, l}^{\lambda} c_{\mu, \nu}^{\theta}$  in Equation (1.2) is an integer polynomial in  $b$  (we use the induction hypothesis and Theorem 1.2). This implies that the left hand-side  $\sum_{\kappa \vdash n} c_{\mu, \kappa}^{\lambda} c_{\nu, l}^{\kappa}$  in Equation (1.2) is an integer polynomial in  $b$ . We conclude using the two following facts:

- if  $\text{rk}(\kappa) > r$  then  $c_{\nu, l}^{\kappa} = 0$ . Indeed, from Lemma 2.1, we get that

$$\deg_b(c_{\nu, l}^{\kappa}) \leq n - l + \text{rk}(\nu) - \text{rk}(\kappa) = r - \text{rk}(\kappa) < 0.$$

- if  $\text{rk}(\kappa) < r$  then we know that  $c_{\mu, \kappa}^{\lambda}$  is an integer polynomial from  $\mathcal{A}_n^{(\text{rk}(\kappa))}$ .  $\square$

For fixed  $\lambda, \mu \vdash n$  and  $r < n$ , one can note that we get from the previous lemma more equations of type (3.2) than variables  $c_{\mu, \kappa}^{\lambda}$ , where  $\text{rk}(\kappa) = r$ . In order to obtain a square system, we consider equations (3.2) indexed by partitions  $(\nu, l)$  satisfying the following condition that refines condition (3.1):

$$(\nu, l) = (\pi \cup 1^{n-r}, l), \text{ where } \pi \vdash r \text{ and } n - l + \text{rk}(\pi) = r. \quad (3.3)$$

We denote by  $\mathcal{S}_{\lambda, \mu}^{(r)}$  the linear system obtained by taking the equations (3.2) for  $(\nu, l)$  satisfying condition (3.3), and we denote by  $\mathcal{Q}_n^{(r)}$  the matrix associated to this system. In other terms  $\mathcal{Q}_n^{(r)} = (c_{\nu, l}^{\kappa})$  where indices  $\kappa$  of columns are partitions of  $n$  of rank  $r$ , and indices of rows are pairs  $(\nu, l)$  satisfying the condition (3.3). Note that this matrix is independent from  $\lambda$  and  $\mu$ . In the case  $2r \leq n$ , one can see that the system  $\mathcal{S}_{\lambda, \mu}^{(r)}$  is a square system and that  $\mathcal{Q}_n^{(r)}$  is the matrix  $\mathcal{Q}^{(r)}$  defined in Section 2. The situation in the case  $2r > n$  is more intricate. In general, we have the following proposition:

**Proposition 3.2.** *For every  $1 \leq r < n$ , the matrix  $\mathcal{Q}_n^{(r)}$  is a submatrix of  $\mathcal{Q}^{(r)}$  defined in Section 2, obtained by erasing only columns. More precisely,  $\mathcal{Q}_n^{(r)} = (t_{\pi}^{\rho})$ , where the rows index  $\pi$  is a partition of  $r$  and the columns index  $\rho$  is a partition of  $r$  such that  $\ell(\rho) \leq n - r$ .*

*Proof.* We have the following bijection

$$\begin{aligned} \{\kappa \text{ such that } \kappa \vdash n \text{ and } \text{rk}(\kappa) = r\} &\xrightarrow{\sim} \{\rho \text{ such that } \rho \vdash r \text{ and } \ell(\rho) \leq n - r\}, \\ \kappa &\longmapsto \kappa - \mathbf{1}, \\ \rho \oplus 1^{n-r} &\longleftarrow \rho, \end{aligned}$$

Recall that  $t_\pi^\rho = c_{\pi \cup 1^{n-r}, n-r+\text{rk}(\pi)}^\kappa$ , where  $\kappa = \rho \oplus 1^{n-r}$  (see [Definition 2.1](#)). This concludes the proof.  $\square$

We now prove [Theorem 1.3](#).

*Proof of Theorem 1.3.* We prove  $\mathcal{A}_n^{(r)}$  by induction on  $r$ . For  $r = 0$ , the only partition of rank 0 is  $\kappa = [1^n]$  and we know that  $c_{\mu, [1^n]}^\lambda = \delta_{\lambda, \mu}$  for all partitions  $\lambda, \mu$ , where  $\delta_{\lambda, \mu}$  is the Kronecker delta (see [\[11, Lemma 3.3\]](#)).

Now we fix  $r > 0$  and we assume that  $\mathcal{A}_n^{(j)}$  holds for each  $j \leq r - 1$ . We fix two partitions  $\lambda, \mu \vdash n \geq 1$ , and we consider the system  $\mathcal{S}_{\lambda, \mu}^{(r)}$ . It can be written as follows:  $\mathcal{Q}_n^{(r)} X_{\lambda, \mu}^{(r)} = Y_{\lambda, \mu}^{(r)}$ , where  $Y_{\lambda, \mu}^{(r)}$  is the column vector containing the polynomials  $P_{\lambda, \mu, \nu, l}^{(r)}$  for  $(\nu, l)$  satisfying [\(3.3\)](#), and  $X_{\lambda, \mu}^{(r)}$  is the column vector containing  $c_{\mu, \kappa}^\lambda$  for  $\kappa \vdash n$  of rank  $r$ . We denote the column vector  $\tilde{X}_{\lambda, \mu}^{(r)} := (x_{\mu, \rho}^\lambda)$  for  $\rho \vdash r$ , where

$$x_{\mu, \rho}^\lambda := \begin{cases} c_{\mu, \rho \oplus 1^{n-r}}^\lambda & \text{if } \ell(\rho) \leq n - r, \\ 0 & \text{otherwise.} \end{cases}$$

The system  $\mathcal{S}_{\lambda, \mu}^{(r)}$  can be rewritten as follows:  $\mathcal{Q}^{(r)} \tilde{X}_{\lambda, \mu}^{(r)} = Y_{\lambda, \mu}^{(r)}$ . But we know from [Theorem 2.3](#) that  $\mathcal{Q}^{(r)}$  is invertible in  $\mathbb{Z}$ , and since the entries of  $Y_{\lambda, \mu}^{(r)}$  are integer polynomials in  $b$  (see [Lemma 3.1](#)), we deduce that this is also the case for the entries of  $\tilde{X}_{\lambda, \mu}^{(r)}$ . Hence, the coefficients  $c_{\mu, \kappa}^\lambda$  are integer polynomials in  $b$ , when the partition  $\kappa$  has rank  $r$ . This gives the assertion  $\mathcal{A}_n^{(r)}$ .  $\square$

*Remark 1.* Note that the previous proof implies that [Equation \(1.2\)](#) allows to recover the coefficients  $c_{\mu, \nu}^\lambda$  from their marginal sums. More precisely, if we have a family  $(y_{\mu, \nu}^\lambda)_{\lambda, \mu, \nu \vdash n}$  of rational functions in  $b$  indexed by partitions of size  $n$ , satisfying

$$y_{\mu, [1^n]}^\lambda = \delta_{\lambda, \mu} \quad \text{and} \quad \sum_{\kappa \vdash n} y_{\mu, \kappa}^\lambda c_{\nu, l}^\kappa = \sum_{\theta \vdash n} c_{\theta, l}^\lambda y_{\mu, \nu}^\theta$$

then we have that  $y_{\mu, \nu}^\lambda = c_{\mu, \nu}^\lambda$  for every partitions  $\lambda, \mu, \nu \vdash n$ .

## 4 Graded Farahat–Higman Algebra

In this section we explain the main steps of the proof of [Theorem 2.3](#). The main idea is to see the matrix  $Q^{(r)}$  as a change-of-basis matrix in the Farahat–Higman algebra.

If  $\sigma$  is a permutation of cyclic type  $\lambda$ , we define its *reduced cyclic type* as the partition  $\lambda - \mathbf{1}$  (see [\[20\]](#)). Hence if  $\sigma \in \mathfrak{S}_n \subset \mathfrak{S}_{n+1} \cdots$ , the reduced cyclic type of  $\sigma$  does not depend on  $n$ . For every partition  $\lambda$ , we define  $C_\lambda(n) \in \mathbb{Z}\mathfrak{S}_n$  as the sum of all permutations in  $\mathfrak{S}_n$  of reduced cyclic type  $\lambda$ . Note that  $C_\lambda(n) = 0$  if  $|\lambda| + \ell(\lambda) > n$ . Hence, for every  $n \geq 0$ , the family  $(C_\lambda(n))_{|\lambda| + \ell(\lambda) \leq n}$  form a basis of the center of the group algebra of  $\mathfrak{S}_n$ :

$$Z(\mathbb{Z}\mathfrak{S}_n) = \text{Span}_{\mathbb{Z}}\{C_\lambda(n), |\lambda| + \ell(\lambda) \leq n\}.$$

The multiplication in this algebra is given by

$$C_\lambda(n)C_\mu(n) = \sum_{|\kappa| + \ell(\kappa) \leq n} \rho_{\lambda, \mu}^\kappa(n) C_\kappa(n), \quad (4.1)$$

for some structure coefficients  $\rho_{\lambda, \mu}^\kappa(n)$ . The latter are linked to the coefficients  $c_{\mu, \nu}^\lambda$  as follows (see [\[11, Proposition 3.1\]](#)):

$$\rho_{\lambda, \mu}^\kappa(n) = \begin{cases} c_{\lambda \oplus \mathbf{1}^{n-|\kappa|}, \mu \oplus \mathbf{1}^{n-|\mu|}}^{\kappa \oplus \mathbf{1}^{n-|\kappa|}}(0) & \text{if } \max(|\lambda| + \ell(\lambda), |\mu| + \ell(\mu), |\kappa| + \ell(\kappa)) \leq n, \\ 0 & \text{otherwise.} \end{cases} \quad (4.2)$$

The following proposition is due to Farahat and Higman.

**Proposition 4.1** ([\[9\]](#)). *The structure coefficients  $\rho_{\lambda, \mu}^\kappa$  satisfy the following properties:*

1.  $\rho_{\lambda, \mu}^\kappa = 0$  if  $|\kappa| > |\lambda| + |\mu|$ .
2.  $\rho_{\lambda, \mu}^\kappa$  is independent from  $n$  if  $|\kappa| = |\lambda| + |\mu|$ .
3.  $\rho_{\lambda, \mu}^\kappa$  is a polynomial in  $n$  if  $|\kappa| < |\lambda| + |\mu|$ .

We are here interested in the structure coefficients  $\rho_{\lambda, \mu}^\kappa$  in the case  $|\kappa| = |\lambda| + |\mu|$ , these coefficients are called the *top connection coefficients* of the Farahat–Higman algebra. To study these coefficients, we consider the graded algebra  $\mathcal{Z}_n$  associated to  $Z(\mathbb{Z}\mathfrak{S}_n)$  with respect to the filtration  $\deg(C_\lambda(n)) = |\lambda|$ . We denote by  $\mathbf{c}_\lambda(n)$  the image of  $C_\lambda(n)$  in  $\mathcal{Z}_n$ . Concretely,  $\mathcal{Z}_n = \bigoplus_{1 \leq r \leq n-1} \mathcal{Z}_n^{(r)}$ , where  $\mathcal{Z}_n^{(r)} := \text{Span}_{\mathbb{Z}}\{\mathbf{c}_\lambda(n) : \lambda \vdash r \text{ and } \ell(\lambda) \leq n - r\}$ , and the multiplication in  $\mathcal{Z}_n$  is defined by

$$\mathbf{c}_\lambda(n) \mathbf{c}_\mu(n) = \sum_{\kappa \vdash |\lambda| + |\mu|} \rho_{\lambda, \mu}^\kappa(n) \mathbf{c}_\kappa(n) = \sum_{\kappa \vdash |\lambda| + |\mu|} c_{\lambda \oplus \mathbf{1}^{n-|\kappa|}, \mu \oplus \mathbf{1}^{n-|\mu|}}^{\kappa \oplus \mathbf{1}^{n-|\kappa|}}(0) \mathbf{c}_\kappa(n).$$



Note that compared to [Equation \(4.1\)](#), we keep only the top degree terms. The graded algebra  $\mathcal{Z}_n$  comes with a linear isomorphism  $\phi_n: Z(\mathbb{Z}\mathfrak{S}_n) \xrightarrow{\sim} \mathcal{Z}_n$ , that sends  $C_\lambda(n)$  to  $c_\lambda(n)$  (which is obviously not an algebra morphism). Since the structure coefficients in  $\mathcal{Z}_n$  are independent from  $n$  (see [Proposition 4.1](#) item 2), we can define a family of  $\mathbb{Z}$ -algebra morphisms:

$$\begin{aligned} \psi_n: \mathcal{Z}_{n+1} &\longrightarrow \mathcal{Z}_n \\ c_\lambda(n+1) &\longmapsto \begin{cases} c_\lambda(n) & \text{if } |\lambda| + \ell(\lambda) \leq n, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Let  $\mathcal{Z}_\infty := \varprojlim \mathcal{Z}_n$  be the projective limit of the  $\mathcal{Z}_n$ 's, and let the limit  $c_\lambda := \varprojlim c_\lambda(n)$  in  $\mathcal{Z}_\infty$ . We also define  $\mathcal{Z}_\infty^{(r)} := \text{Span}_{\mathbb{Z}}\{c_\lambda : \lambda \vdash r\}$ , hence  $\mathcal{Z}_\infty = \bigoplus_{r \geq 1} \mathcal{Z}_\infty^{(r)}$  (see [[19](#), Example 24, page 131] for further details about the construction of the algebra  $\mathcal{Z}_\infty$ ).

For every  $r \geq 1$ , we set  $f_r(n) := \sum_{\lambda \vdash r} c_\lambda(n)$ , and for every partition  $\mu$ ,  $f_\mu(n) := \prod_i f_{\mu_i}(n)$ . We also define  $g_\nu(n) := c_{\nu-1}(n) f_{\ell(\nu)}(n)$ , for every partition  $\nu$ . Note that

$$\deg(f_\nu(n)) = \deg(g_\nu(n)) = |\nu|.$$

Finally we define the limits  $f_\mu := \varprojlim f_\mu(n)$  and  $g_\nu := \varprojlim g_\nu(n)$  in  $\mathcal{Z}_\infty$ . We have the following theorem due to Farahat and Higman.

**Theorem 4.2** ([[9](#)]). *For every  $r \geq 0$ ,  $(f_\mu)_{\mu \vdash r}$  is a  $\mathbb{Z}$ -basis of  $\mathcal{Z}_\infty^{(r)}$ .*


The following proposition relates the matrix  $Q^{(r)}$  defined in [Section 2](#) to the graded Farahat–Higman algebra  $\mathcal{Z}_\infty$ .

**Proposition 4.3.** *For every partitions  $\lambda, \mu \vdash r \geq 0$ , we have  $t_\mu^\lambda = [c_\lambda] g_\mu$ . In other terms, for every  $r > 0$ , the matrix  $(Q^{(r)})^T$  is the matrix of  $(g_\mu)_{\mu \vdash r}$  in the basis  $(c_\lambda)_{\lambda \vdash r}$ .*

Hence, our goal is to prove the following theorem that implies [Theorem 2.3](#):

**Theorem 4.4.** *For every  $r \geq 0$ , the family  $(g_\mu)_{\mu \vdash r}$  is a  $\mathbb{Z}$ -basis for  $\mathcal{Z}_\infty^{(r)}$ .*

The proof of [Theorem 4.4](#) involves studying triangularity properties of change-of-basis matrices in the algebra  $\mathcal{Z}_\infty$ . The indices of the different bases should be considered with some particular orders on partitions that are not detailed here. We explain here the different steps of the proof on examples for small values of  $r$ , the full proof is given in [[3](#)]. The following diagram illustrates the different basis of  $\mathcal{Z}_\infty^{(r)}$  involved in the proof of [Theorem 4.4](#), and the associated transition matrices.



In order to prove that  $(\mathbf{g}_\lambda)$  is a  $\mathbb{Z}$  basis of  $\mathcal{Z}_\infty$ , we consider its matrix  $\mathcal{N}$  in the basis  $(\mathbf{f}_\lambda)$ . With elementary manipulations of the definitions, we can see that this matrix is block upper triangular, when the partitions indexing the bases are considered with some particular orders.

*Example 4.1.* We give here the matrix  $\mathcal{N}^{(r)}$  for  $r = 5$  (the diagonal blocks are colored in gray).

$\mathbf{f} \setminus \mathbf{g}$	$[1^5]$	$[2, 1^3]$	$[2^2, 1]$	$[3, 1^2]$	$[3, 2]$	$[4, 1]$	$[5]$
$[5]$	1	0	0	0	0	0	0
$[4, 1]$		1	0	0	0	0	-4
$[3, 2]$			3	-1	-12	3	0
$[3, 1^2]$			-2	1	0	0	4
$[2^2, 1]$					10	-3	2
$[2, 1^3]$					-3	1	-4
$[1^5]$							1

We need to prove that each one of the diagonal blocks of this matrix has determinant  $\pm 1$ . To this purpose, we prove that these diagonal blocks are South-East blocks of the matrix  $\mathcal{M}^{(i)}$ , the matrix of  $(\mathbf{c}_\lambda)_{\lambda+i}$  in  $(\mathbf{f}_\lambda)_{\lambda+i}$ , where  $i \leq r$ .

*Example 4.2.* For  $r = 3$ ,

$$\mathcal{M}^{(3)} \text{ is given by } \begin{array}{|c|c|c|c|} \hline \mathbf{f} \setminus \mathbf{c} & [1^3] & [2, 1] & [3] \\ \hline [3] & 10 & -12 & 3 \\ \hline [2, 1] & -7 & 10 & -3 \\ \hline [1^3] & 2 & -3 & 1 \\ \hline \end{array} \text{ and } \mathcal{M}^{(2)} \text{ by } \begin{array}{|c|c|c|} \hline \mathbf{f} \setminus \mathbf{c} & [1^2] & [2] \\ \hline [2] & 3 & -1 \\ \hline [1^2] & -2 & 1 \\ \hline \end{array}.$$

Note that the third diagonal block of the matrix  $\mathcal{N}^{(3)}$  given in [Example 4.1](#) is equal to the matrix  $\mathcal{M}^{(2)}$  and its fourth diagonal block is a South-East diagonal block of  $\mathcal{M}^{(3)}$ .

We need to show that the East-South blocks of  $\mathcal{M}^{(r)}$  have determinant 1. We show that the matrix  $\mathcal{M}^{(r)}$  has a decomposition  $\mathcal{M}^{(r)} = \mathcal{U}^{(r)}\mathcal{L}^{(r)}$  where  $\mathcal{U}^{(r)}$  (respectively  $\mathcal{L}^{(r)}$ ) is an upper (respectively lower) triangular matrix, with coefficients equal to 1 on the diagonal. This decomposition is obtained by considering an intermediate basis  $(\mathbf{m}_\lambda)$ , defined as the evaluation of the monomial symmetric functions in the Jucys–Murphy

elements. Indeed, this basis has the property that its transition matrices to both bases  $(c_\lambda)$  and  $(f_\lambda)$  are triangular (we use a result of Matsumoto and Novak [20] and some basic properties on symmetric functions). Finally, we note that this decomposition of the matrix  $\mathcal{M}^{(r)}$  induces a similar decomposition for every one of its South-East block, and hence these blocks have determinant 1.

*Example 4.3.* For  $r = 3$ , we have the decomposition  $\mathcal{M}^{(3)} = \mathcal{U}^{(3)} \mathcal{L}^{(3)}$ , where

$$\mathcal{U}^{(3)} \text{ is given by } \begin{array}{c|ccc} \mathfrak{f} \setminus \mathfrak{m} & [1^3] & [2, 1] & [3] \\ \hline [3] & 1 & -3 & 3 \\ [2, 1] & 0 & 1 & -3 \\ [1^3] & 0 & 0 & 1 \end{array} \text{ and } \mathcal{L}^{(3)} \text{ by } \begin{array}{c|ccc} \mathfrak{m} \setminus \mathfrak{c} & [1^3] & [2, 1] & [3] \\ \hline [3] & 1 & 0 & 0 \\ [2, 1] & -1 & 1 & 0 \\ [1^3] & 2 & -3 & 1 \end{array}.$$

## Acknowledgements

The author would like to thank his advisors Valentin Féray and Guillaume Chapuy for many interesting discussions and for reviewing this paper several times. He also thanks Guillaume Chapuy and Maciej Dołęga for suggesting to consider the multiplicativity property in order to deduce properties of the coefficients  $c_{\mu, \nu}^\lambda$ .

This is an extended abstract of [3].

## References

- [1] F. Bédard and A. Goupil. “The poset of conjugacy classes and decomposition of products in the symmetric group”. *Canad. Math. Bull.* **35.2** (1992), 152–160.
- [2] H. Ben Dali. “Generating series of non-orientable constellations, and marginal sums in the Matching-Jack conjecture” (2021). [arXiv:2106.15414](https://arxiv.org/abs/2106.15414).
- [3] H. Ben Dali. “Integrality in the Matching-Jack conjecture and the Farahat-Higman algebra” (2022). [arXiv:2203.14879](https://arxiv.org/abs/2203.14879).
- [4] G. Chapuy. “A new combinatorial identity for unicellular maps, via a direct bijective approach”. *Adv. in Appl. Math* **47.4** (2011), pp. 874–893.
- [5] G. Chapuy and M. Dołęga. “Non orientable branched coverings,  $b$ -Hurwitz numbers, and positivity for multiparametric Jack expansions”. 2020. [arXiv:2004.07824](https://arxiv.org/abs/2004.07824).
- [6] M. Dołęga and V. Féray. “Gaussian fluctuations of Young diagrams and structure constants of Jack characters”. *Duke Math. J.* **165.7** (2016), pp. 1193–1282.
- [7] M. Dołęga, V. Féray, and P. Sniady. “Jack polynomials and orientability generating series of maps”. *Sém. Lothar. Combin.* **70** (2014), Article B70j.
- [8] B. Eynard. *Counting Surfaces*. Vol. 70. Progress in Mathematical Physics. Springer, 2016.

- [9] H. K. Farahat and G. Higman. “The centres of symmetric group rings”. *Proc. Roy. Soc. Edinburgh Sect. A* **250.1261** (1959), pp. 212–221.
- [10] I. P. Goulden and D. M. Jackson. “Symmetrical functions and Macdonald’s result for top connexion coefficients in the symmetrical group”. *J. Algebra* **166.2** (1994), pp. 364–378.
- [11] I. P. Goulden and D. M. Jackson. “Connection coefficients, matchings, maps and combinatorial conjectures for Jack symmetric functions”. *Trans. Amer. Math. Soc.* **348.3** (1996), pp. 873–892.
- [12] I. P. Goulden and D. M. Jackson. “Maps in locally orientable surfaces, the double coset algebra, and zonal polynomials”. *Canadian J. Math.* **48.3** (1996), pp. 569–584.
- [13] V. N. Ivanov and S. V. Kerov. “The algebra of conjugacy classes in symmetric groups, and partial permutations”. *Zapiski Nauchnykh Seminarov POMI* **256** (1999), pp. 95–120.
- [14] H. Jack. “A class of symmetric polynomials with a parameter”. *Proc. Roy. Soc. Edinburgh Sect. A* **69** (1971), pp. 1–18.
- [15] A. L. Kanunnikov, V. V. Promyslov, and E. A. Vassilieva. “A labelled variant of the matchings-Jack and hypermap-Jack conjectures”. *Sém. Lothar. Combin.* **80B.1** (2018), Art. 45, 12.
- [16] A. L. Kanunnikov and E. A. Vassilieva. “On the Matchings-Jack conjecture for Jack connection coefficients indexed by two single part partitions”. *Electron. J. Combin.* **23.1** (2016), Paper 1.53, 30. MR 3484758.
- [17] M. A. La Croix. “The combinatorics of the Jack parameter and the genus series for topological maps”. PhD thesis. University of Waterloo, 2009.
- [18] S. K. Lando and A. K. Zvonkin. *Graphs on Surfaces and Their Applications*. Vol. 141. Encyclopaedia of Mathematical Sciences, Springer-Verlag, Berlin, 2004.
- [19] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Second edition. Oxford Mathematical Monographs, Clarendon Press, Oxford, 1995.
- [20] S. Matsumoto and J. Novak. “Jucys–Murphy Elements and Unitary Matrix Integrals”. *Int. Math. Res. Not. IMRN* **2013.2** (2013), pp. 362–397.
- [21] R. Stanley. “Some combinatorial properties of Jack symmetric functions”. *Adv. Math.* **77** (1989), pp. 76–115.
- [22] C. Sylvie, A. Goupil, and G. Schaeffer. “Content evaluation and class symmetric functions”. *Adv. Math.* **188.2** (2004), pp. 315–336.