

GAMMA-POSITIVITY IN COMBINATORICS AND GEOMETRY

CHRISTOS A. ATHANASIADIS

ABSTRACT. Gamma-positivity is an elementary property that polynomials with symmetric coefficients may have, which directly implies their unimodality. The idea behind it stems from work of Foata, Schützenberger and Strehl on the Eulerian polynomials; it was revived independently by Brändén and Gal in the course of their study of poset Eulerian polynomials and face enumeration of flag simplicial spheres, respectively, and has found numerous applications since then. This paper surveys some of the main results and open problems on gamma-positivity, appearing in various combinatorial or geometric contexts, as well as some of the diverse methods that have been used to prove it.

CONTENTS

1. Introduction	2
2. Gamma-positivity in combinatorics	3
2.1. Variations of Eulerian polynomials	3
2.2. Coxeter–Narayana polynomials	24
2.3. Polynomials arising from enriched poset partitions	27
2.4. Polynomials arising from poset homology	28
2.5. Polynomials arising from symmetric functions	30
2.6. Polynomials arising from trees	33
3. Gamma-positivity in geometry	34
3.1. Flag triangulations of spheres	35
3.2. Flag triangulations of simplices	39
3.3. Examples	42
4. Methods	46
4.1. Valley hopping	47
4.2. Methods of geometric combinatorics	49
4.3. Symmetric/quasisymmetric function methods	52
5. Generalizations	54
5.1. Nonsymmetric γ -positivity	54
5.2. Equivariant γ -positivity	55
5.3. q - γ -positivity	58
References	59

2010 *Mathematics Subject Classification*. Primary 05A15, 05E45; Secondary 05A05, 05E05, 06A11, 20F55.

Key words and phrases. Gamma-positivity, unimodality, Eulerian polynomial, flag triangulation, h -polynomial, local h -polynomial, quasisymmetric function.

1. INTRODUCTION

The unimodality of polynomials is a major theme which has occupied mathematicians for the past few decades. At least three surveys [27, 32, 122] about unimodal, log-concave and real-rooted polynomials, showing the enormous variety of methods which are available to prove these properties, have been written. This survey article focuses on a related elementary property, that of γ -positivity, which directly implies symmetry and unimodality and has provided a new exciting approach to this topic. Gamma-positivity appears surprisingly often in combinatorial and geometric contexts; this article aims to discuss some of the main examples, results, methods and open problems around it.

This introductory section provides basic definitions and related comments, as well as an outline of the remainder of the article. We recall that a polynomial $f(x) = \sum_i a_i x^i \in \mathbb{R}[x]$ is called

- *symmetric*, with center of symmetry $n/2$, if $a_i = a_{n-i}$ for all $i \in \mathbb{Z}$ (where $a_i = 0$ for negative values of i),
- *unimodal*, if $0 \leq a_0 \leq a_1 \leq \dots \leq a_k \geq a_{k+1} \geq \dots$ for some $k \in \mathbb{N}$,
- *γ -positive*, if

$$(1) \quad f(x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \gamma_i x^i (1+x)^{n-2i}$$

for some $n \in \mathbb{N}$ and nonnegative reals $\gamma_0, \gamma_1, \dots, \gamma_{\lfloor n/2 \rfloor}$, and

- *real-rooted*, if every root of $f(x)$ is real, or else $f(x) = 0$.

The notion of γ -positivity appeared first in the work of D. Foata and M. Schützenberger [56] and subsequently of D. Foata and V. Strehl [57, 58] on the classical Eulerian polynomials, discussed in detail in Sections 2.1.1 and 4.1. After having implicitly reappeared in the theory of enriched poset partitions of J. Stembridge [133] (see also Section 2.3), it was brought again to light independently by P. Brändén [24, 26] and Š. Gal [61] in the course of their study of poset Eulerian polynomials and face enumeration of flag triangulations of spheres, respectively (see Sections 2.1.2 and 3.1). These works made it clear that γ -positivity is a concept of independent interest which provides a powerful approach to the problem of unimodality for symmetric polynomials. Symmetry and real-rootedness of a polynomial $f(x) \in \mathbb{R}_{\geq 0}[x]$ implies its γ -positivity [24, Lemma 4.1] [61, Remark 3.1.1]. On the other hand, every γ -positive polynomial (whether real-rooted or not) is symmetric and unimodal, as a sum of symmetric and unimodal polynomials with a common center of symmetry (a fact already implicit in [62, p. 136]). Thus, γ -positivity can be applied to general situations and may lead to a more elementary proof of the unimodality of $f(x)$, even when the latter is real-rooted. Moreover, an explicit expression of the form (1) gives additional information about $f(x)$ (for instance, it implies a formula for $f(-1)$ and predicts its sign) and the problem to interpret algebraically or combinatorially the coefficients γ_i is often of independent interest.

The present article is an expanded version of the author's lectures at the *77th Séminaire Lotharingien de Combinatoire* (September 2016, Strobl, Austria). Section 2 discusses the

variety of examples of γ -positive polynomials in combinatorics, beginning with the prototypical example of Eulerian polynomials, together with some general results. Many of these examples come up again in the geometric contexts of Section 3. That section centers around Gal’s conjecture, claiming that h -polynomials of flag simplicial homology spheres are γ -positive. This conjecture and its relatives provide a general framework under which a lot of the seemingly unrelated γ -positivity phenomena of Section 2 can be considered, partially explains their abundance and allows for tools from geometric combinatorics to be applied to their study. Section 4 discusses the plethora of methods used in the literature to prove γ -positivity. Section 5 is devoted to generalizations and variations of the concept of γ -positivity, giving an emphasis to equivariant and symmetric function generalizations. The choice of topics is strongly affected by the author’s knowledge and personal taste; in particular, possible probabilistic aspects of γ -positivity are not treated. Some previously unpublished statements and open problems are included. Other expositions on γ -positivity can be found in [27, Section 3] [97, Chapter 4].

Notation. For integers $a \leq b$ we set $[a, b] := \{a, a + 1, \dots, b\}$ and use the abbreviation $[n] := [1, n]$ for $n \in \mathbb{N}$. We will denote by $|\Omega|$ the cardinality, and by 2^Ω the set of all subsets, of a finite set Ω . For $\Omega \subseteq \mathbb{Z}$, we will also denote by $\text{Stab}(\Omega)$ the set of all subsets of Ω which do not contain two consecutive integers.

2. GAMMA-POSITIVITY IN COMBINATORICS

This section describes instances of γ -positivity in combinatorics. Much of the motivation comes from the study of Eulerian polynomials which, along with generalizations and variations, are discussed in detail. For any undefined notation or terminology, we refer to Stanley’s textbooks [125, 127].

2.1. Variations of Eulerian polynomials.

2.1.1. *Eulerian polynomials.* One of the most important polynomials in combinatorics is the *Eulerian polynomial*, defined by any of the equivalent formulas

$$(2) \quad A_n(x) := \sum_{w \in \mathfrak{S}_n} x^{\text{asc}(w)} = \sum_{w \in \mathfrak{S}_n} x^{\text{des}(w)} = \sum_{w \in \mathfrak{S}_n} x^{\text{exc}(w)} = \sum_{w \in \mathfrak{S}_n} x^{\text{wexc}(w)-1}$$

for every positive integer n . Here $\text{asc}(w)$, $\text{des}(w)$, $\text{exc}(w)$ and $\text{wexc}(w)$ denotes the cardinality of the set $\text{Asc}(w)$ of *ascents*, $\text{Des}(w)$ of *descents*, $\text{Exc}(w)$ of *excedances* and $\text{Wexc}(w)$ of *weak excedances*, respectively, of the permutation $w \in \mathfrak{S}_n$, defined by

- $\text{Asc}(w) := \{i \in [n - 1] : w(i) < w(i + 1)\}$,
- $\text{Des}(w) := \{i \in [n - 1] : w(i) > w(i + 1)\}$,
- $\text{Exc}(w) := \{i \in [n - 1] : w(i) > i\}$,
- $\text{Wexc}(w) := \{i \in [n] : w(i) \geq i\}$.

For the first few values of n , we have

$$A_n(x) = \begin{cases} 1, & \text{if } n = 1, \\ 1 + x, & \text{if } n = 2, \\ 1 + 4x + x^2, & \text{if } n = 3, \\ 1 + 11x + 11x^2 + x^3, & \text{if } n = 4, \\ 1 + 26x + 66x^2 + 26x^3 + x^4, & \text{if } n = 5, \\ 1 + 57x + 302x^2 + 302x^3 + 57x^4 + x^5, & \text{if } n = 6, \\ 1 + 120x + 1191x^2 + 2416x^3 + 1191x^4 + 120x^5 + x^6, & \text{if } n = 7. \end{cases}$$

The Eulerian polynomial $A_n(x)$, which is clearly symmetric, provides the prototypical example of a γ -positive polynomial in combinatorics. The corresponding γ -coefficients for the first few values of n are determined from the expressions

$$A_n(x) = \begin{cases} 1 + x, & \text{if } n = 2, \\ (1 + x)^2 + 2x, & \text{if } n = 3, \\ (1 + x)^3 + 8x(1 + x), & \text{if } n = 4, \\ (1 + x)^4 + 22x(1 + x)^2 + 16x^2, & \text{if } n = 5, \\ (1 + x)^5 + 52x(1 + x)^3 + 136x^2(1 + x), & \text{if } n = 6, \\ (1 + x)^6 + 114x(1 + x)^4 + 720x^2(1 + x)^2 + 272x^3, & \text{if } n = 7. \end{cases}$$

The γ -positivity of $A_n(x)$ (which, of course, follows from the well known fact that $A_n(x)$ is real-rooted for all n) was first shown combinatorially by Foata and Schützenberger [56, Theorem 5.6]. An explicit combinatorial interpretation of the corresponding γ -coefficients follows from the results of [58]. Recall that a *double excedance* of $w \in \mathfrak{S}_n$ is any index $1 \leq i \leq n$ such that $w(i) > i > w^{-1}(i)$ and that $w \in \mathfrak{S}_n$ is called an *up-down* permutation, if $\text{Asc}(w) = \{1, 3, 5, \dots\} \cap [n-1]$. The first four, as well as the last two, interpretations of $\gamma_{n,i}$ in the following fundamental result are easily shown to be equivalent to one another; the fifth one follows from the fourth and the bijection (one of the fundamental transformations of Foata and Schützenberger [56]) of [127, Proposition 1.3.1]. An additional interpretation, in terms of increasing binary trees, is discussed in [62, p. 136].

Theorem 2.1 (cf. FOATA–STREHL [58]). *For all $n \geq 1$,*

$$(3) \quad A_n(x) = \sum_{i=0}^{\lfloor (n-1)/2 \rfloor} \gamma_{n,i} x^i (1+x)^{n-1-2i},$$

where $\gamma_{n,i}$ is equal to each of the following:

- the number of $w \in \mathfrak{S}_n$ for which $\text{Asc}(w) \in \text{Stab}([n-2])$ has i elements,
- the number of $w \in \mathfrak{S}_n$ for which $\text{Asc}(w) \in \text{Stab}([2, n-1])$ has i elements,
- the number of $w \in \mathfrak{S}_n$ for which $\text{Des}(w) \in \text{Stab}([n-2])$ has i elements,
- the number of $w \in \mathfrak{S}_n$ for which $\text{Des}(w) \in \text{Stab}([2, n-1])$ has i elements,
- the number of $w \in \mathfrak{S}_n$ with i excedances and no double excedance, for which the smallest of the maximum elements of the cycles of w is a fixed point,

- the number of $w \in \mathfrak{S}_n$ with i excedances and no double excedance, for which the largest of the minimum elements of the cycles of w is a fixed point.

In particular, $A_n(x)$ is γ -positive and

$$(4) \quad A_n(-1) = \begin{cases} 0, & \text{if } n \text{ is even,} \\ (-1)^{(n-1)/2} \gamma_{n,(n-1)/2}, & \text{if } n \text{ is odd,} \end{cases}$$

for all n , where $\gamma_{n,(n-1)/2}$ is the number of up-down permutations in \mathfrak{S}_n .

We will now describe some refinements and variations of this theorem (more related results appear in the sequel). The following two theorems refine the third interpretation of the γ -positivity of $A_n(x)$, given in Theorem 2.1. For $w \in \mathfrak{S}_n$, we denote by $\text{inv}(w)$ the number of inversions (pairs $(i, j) \in [n] \times [n]$ such that $i < j$ and $w(i) > w(j)$) of w , and let

- $(2-13)w$ be the number of pairs $(i, j) \in [n-1] \times [n-1]$ such that $i < j$ and $w(j) < w(i) < w(j+1)$,
- $(31-2)w$ be the number of pairs $(i, j) \in [n] \times [n]$ such that $i+1 < j$ and $w(i+1) < w(j) < w(i)$,
- $\text{cros}(w)$ be the number of pairs $(i, j) \in [n] \times [n]$ such that $i < j \leq w(i) < w(j)$, or $w(i) < w(j) < i < j$,
- $\text{nest}(w)$ be the number of pairs $(i, j) \in [n] \times [n]$ such that $i < j \leq w(j) < w(i)$, or $w(j) < w(i) < i < j$.

Equations (5) and (6) appear as the specialization $u = v = w = 1$ of [115, Theorem 2] and as [115, Corollary 6]; the expansion (6) for the left-hand side of (5) was originally shown by Brändén [26, Section 5]. Equation (7) is the main statement of [116, Theorem 1]; the positivity of the coefficient of $x^i(1+x)^{n-1-2i}$, as a polynomial in q , was conjectured in a preprint version of [22]. A related result appears in [22, Remark 3.4].

Theorem 2.2 (BRÄNDÉN [26], SHIN–ZENG [115, 116]). *For all $n \geq 1$,*

$$(5) \quad \sum_{w \in \mathfrak{S}_n} p^{(2-13)w} q^{(31-2)w} x^{\text{des}(w)} = \sum_{w \in \mathfrak{S}_n} p^{\text{nest}(w)} q^{\text{cros}(w)} x^{\text{wexc}(w)-1}$$

$$(6) \quad = \sum_{i=0}^{\lfloor (n-1)/2 \rfloor} a_{n,i}(p, q) x^i (1+x)^{n-1-2i}$$

and

$$(7) \quad \sum_{w \in \mathfrak{S}_n} q^{\text{inv}(w) - \text{exc}(w)} x^{\text{exc}(w)} = \sum_{i=0}^{\lfloor (n-1)/2 \rfloor} a_{n,i}(q^2, q) x^i (1+x)^{n-1-2i},$$

where

$$(8) \quad a_{n,i}(p, q) = \sum_w p^{(2-13)w} q^{(31-2)w},$$

and the sum runs through all permutations $w \in \mathfrak{S}_n$ for which $\text{Des}(w) \in \text{Stab}([n-2])$ has i elements.

The following theorem of J. Shareshian and M. Wachs, the proof of which is sketched in Section 4.3, follows from the methods used to prove the specialization $p = 1$ in [114, Theorem 4.4]. This special case was stated in [111, Remark 5.5] without giving an explicit interpretation for the coefficients $\gamma_i(1, q)$; it is also implicit in [84] (see Equations (1.4) and (6.1) there) and is reproven by different methods in [83]. For $w \in \mathfrak{S}_n$, we set

$$\text{des}^*(w) := \begin{cases} \text{des}(w), & \text{if } w(1) = 1, \\ \text{des}(w) - 1, & \text{if } w(1) > 1, \end{cases}$$

and denote by $\text{maj}(w)$ the major index (sum of the elements of $\text{Des}(w)$) of w .

Theorem 2.3 (cf. [114, Section 4]). *For all $n \geq 1$,*

$$(9) \quad \sum_{w \in \mathfrak{S}_n} p^{\text{des}^*(w)} q^{\text{maj}(w) - \text{exc}(w)} x^{\text{exc}(w)} = \sum_{i=0}^{\lfloor (n-1)/2 \rfloor} \gamma_{n,i}(p, q) x^i (1+x)^{n-1-2i},$$

where

$$(10) \quad \gamma_{n,i}(p, q) = \sum_w p^{\text{des}(w^{-1})} q^{\text{maj}(w^{-1})},$$

and the sum runs through all permutations $w \in \mathfrak{S}_n$ for which $\text{Des}(w) \in \text{Stab}([n-2])$ has i elements.

For further generalizations of the γ -positivity of $A_n(x)$, see, for instance, [26, Section 4] and [92, Section 4].

We close this section by discussing an interesting variant of $A_n(x)$, defined as

$$(11) \quad \tilde{A}_n(x) := 1 + x \sum_{k=1}^n \binom{n}{k} A_k(x)$$

and called a *binomial Eulerian polynomial*. For the first few values of n , we have

$$\tilde{A}_n(x) = \begin{cases} 1 + x, & \text{if } n = 1, \\ 1 + 3x + x^2, & \text{if } n = 2, \\ 1 + 7x + 7x^2 + x^3, & \text{if } n = 3, \\ 1 + 15x + 33x^2 + 15x^3 + x^4, & \text{if } n = 4, \\ 1 + 31x + 131x^2 + 131x^3 + 31x^4 + x^5, & \text{if } n = 5, \\ 1 + 63x + 473x^2 + 883x^3 + 473x^4 + 63x^5 + x^6, & \text{if } n = 6. \end{cases}$$

This polynomial first appeared in an enumerative-geometric context in [102, Section 10.4], where it was shown to equal the h -polynomial of the n -dimensional stellohedron (see Section 3 for an explanation of these terms); its symmetry follows from this interpretation and was rediscovered in [42].

The γ -positivity of $\tilde{A}_n(x)$ follows from a more general theorem of A. Postnikov, V. Reiner and L. Williams [102, Theorem 11.6] (see also Theorem 3.11 in the sequel); for the

first few values of n , we have

$$\tilde{A}_n(x) = \begin{cases} 1 + x, & \text{if } n = 1, \\ (1 + x)^2 + x, & \text{if } n = 2, \\ (1 + x)^3 + 4x(1 + x), & \text{if } n = 3, \\ (1 + x)^4 + 11x(1 + x)^2 + 5x^2, & \text{if } n = 4, \\ (1 + x)^5 + 26x(1 + x)^3 + 43x^2(1 + x), & \text{if } n = 5, \\ (1 + x)^6 + 57x(1 + x)^4 + 230x^2(1 + x)^2 + 61x^3, & \text{if } n = 6. \end{cases}$$

We prefer to state a version due to Shareshian and Wachs, which is similar to Theorem 2.1 and affords a q -analog, similar to that of Theorem 2.3 for $A_n(x)$.

Theorem 2.4 (SHARESHIAN–WACHS [114, Theorem 4.5]). *For all $n \geq 1$,*

$$(12) \quad 1 + x \sum_{k=1}^n \binom{n}{k}_q \sum_{w \in \mathfrak{S}_k} q^{\text{maj}(w) - \text{exc}(w)} x^{\text{exc}(w)} = \sum_{i=0}^{\lfloor n/2 \rfloor} \tilde{\gamma}_{n,i}(q) x^i (1 + x)^{n-2i},$$

where $\binom{n}{k}_q$ is a q -binomial coefficient,

$$(13) \quad \tilde{\gamma}_{n,i}(q) = \sum_w q^{\text{maj}(w^{-1})} = \sum_w q^{\text{inv}(w)},$$

and the sums run through all permutations $w \in \mathfrak{S}_n$ for which $\text{Des}(w) \in \text{Stab}([n-1])$ has i elements. In particular,

$$(14) \quad \tilde{A}_n(x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \tilde{\gamma}_{n,i} x^i (1 + x)^{n-2i},$$

where $\tilde{\gamma}_{n,i}$ is equal to each of the following:

- the number of $w \in \mathfrak{S}_n$ for which $\text{Asc}(w) \in \text{Stab}([n-1])$ has i elements,
- the number of $w \in \mathfrak{S}_n$ for which $\text{Des}(w) \in \text{Stab}([n-1])$ has i elements,
- the number of $w \in \mathfrak{S}_n$ with i excedances and no double excedance.

Moreover,

$$(15) \quad \tilde{A}_n(-1) = \begin{cases} 0, & \text{if } n \text{ is odd,} \\ (-1)^{n/2} \tilde{\gamma}_{n,n/2}, & \text{if } n \text{ is even,} \end{cases}$$

where $\tilde{\gamma}_{n,n/2}$ is the number of up-down permutations in \mathfrak{S}_n .

For an alternative approach to the γ -positivity of $\tilde{A}_n(x)$, see Remark 2.18.

Problem 2.5. *Find a p -analog of Theorem 2.4, similar to Theorem 2.3.*




FIGURE 1. A labeled poset with four elements

2.1.2. *Poset Eulerian polynomials.* Given a partially ordered set (poset, for short) \mathcal{P} with n elements, any bijective map $\omega : \mathcal{P} \rightarrow [n]$ is called a *labeling*. Let us write permutations $w \in \mathfrak{S}_n$ in one-line notation $(w(1), w(2), \dots, w(n))$.

Definition 2.6 (STANLEY [118] [127, Section 3.15.2]). *Let $\omega : \mathcal{P} \rightarrow [n]$ be a labeling of a poset \mathcal{P} . The (\mathcal{P}, ω) -Eulerian polynomial is defined as*

$$A_{\mathcal{P}, \omega}(x) = \sum_{w \in \mathcal{L}(\mathcal{P}, \omega)} x^{\text{des}(w)},$$

where $\mathcal{L}(\mathcal{P}, \omega)$ is the set which consists of all permutations $(a_1, a_2, \dots, a_n) \in \mathfrak{S}_n$ such that $\omega^{-1}(a_i) <_{\mathcal{P}} \omega^{-1}(a_j)$ implies $i < j$.

The polynomial $A_{\mathcal{P}, \omega}(x)$ plays a major role in Stanley's theory of (\mathcal{P}, ω) -partitions [118] [127, Section 3.15]; it reduces to the Eulerian polynomial $A_n(x)$ when \mathcal{P} is an antichain on n elements. For the labeled poset of Figure 2.1.2 we have $\mathcal{L}(\mathcal{P}, \omega) = \{(1, 4, 2, 3), (1, 4, 3, 2), (4, 1, 2, 3), (4, 1, 3, 2), (1, 3, 4, 2)\}$ and $A_{\mathcal{P}, \omega}(x) = 3x + 2x^2$. Examples where $A_{\mathcal{P}, \omega}(x)$ is not real-rooted were given by P. Brändén [25] and J. Stembridge [134] (see also [27, Section 6]), thus disproving long-standing conjectures of J. Neggers [89] [122, Conjecture 1] and R. Stanley [29, Conjecture 1] [32, Conjecture 3.9].

The polynomial $A_{\mathcal{P}, \omega}(x)$ does not depend on ω , when the latter is assumed to be order preserving (such labelings are called *natural*), and is thus denoted simply by $A_{\mathcal{P}}(x)$. For example, the labeling obtained from the one of Figure 1 by swapping 2 and 4 is natural and shows that for this poset $A_{\mathcal{P}}(x) = 1 + 3x + x^2$. Moreover, as a consequence of the reciprocity theorem [127, Theorem 3.15.10] for (\mathcal{P}, ω) -partitions, $A_{\mathcal{P}}(x)$ is symmetric if \mathcal{P} is graded; its unimodality in this case was first shown by V. Reiner and V. Welker [104], whose proof relied on a deep result from algebraic geometry. Brändén [24] [26, Section 6] gave two beautiful combinatorial proofs of the γ -positivity of $A_{\mathcal{P}}(x)$ for the more general class of sign-graded posets.

Theorem 2.7 (BRÄNDÉN [24]). *The polynomial $A_{\mathcal{P}}(x)$ is γ -positive for every finite graded poset \mathcal{P} .*

A generalization to finite crystallographic root systems was given by Stembridge [135].

For a different γ -positivity result for posets, due to Stembridge [133], which generalizes the γ -positivity of $A_n(x)$, see Section 2.3.

2.1.3. *Coxeter Eulerian polynomials.* Let (W, S) be a Coxeter system, with Coxeter length function $\ell_S : W \rightarrow \mathbb{N}$ (see [20, Chapter 1] for definitions). Assuming that W is finite, the W -Eulerian polynomial is defined as

$$(16) \quad W(x) := \sum_{w \in W} x^{\text{des}(w)},$$

where $\text{des}(w)$ is the number of right descents (elements $s \in S$ such that $\ell_S(ws) < \ell_S(w)$) of $w \in W$. The polynomial $W(x)$ was first studied systematically by F. Brenti [31]. It has similar properties as $A_n(x)$, to which it reduces when W is the symmetric group \mathfrak{S}_n ; in particular, $W(x)$ is easily verified to be symmetric for all finite Coxeter groups W . The following statement combines Theorem 2.1 with results of C. Chow [39] and J. Stembridge [135] (the proof that the W -Eulerian polynomials are real-rooted was completed more recently; see [107] and references therein).

Theorem 2.8 ([39] [135, Theorem 1.2]). *The W -Eulerian polynomial is γ -positive for every finite Coxeter group W .*

Problem 2.9. *Find a proof which does not use the classification of finite Coxeter groups.*

A common generalization of Theorems 2.7 and 2.8 is provided by [135, Corollary 7.10], mentioned earlier.

Just as is the case for the symmetric groups and the classical Eulerian polynomial, the corresponding γ -coefficients admit interesting combinatorial interpretations for the other infinite families of finite Coxeter groups as well. We now describe such interpretations for the hyperoctahedral groups \mathcal{B}_n . Recall that \mathcal{B}_n consists of all permutations w of the set $\Omega_n := \{1, -1, 2, -2, \dots, n, -n\}$ satisfying $w(-a) = -w(a)$ for each $a \in \Omega_n$. These can be viewed as signed permutations of length n (see Section 2.1.7 for the more general notion of r -colored permutation). The total order

$$(17) \quad -1 <_r -2 <_r -3 <_r \dots <_r 0 <_r 1 <_r 2 <_r 3 <_r \dots$$

of \mathbb{Z} is convenient to use when \mathcal{B}_n is thought of as a colored permutation (rather than as a Coxeter) group. The \mathcal{B}_n -Eulerian polynomial is given by

$$(18) \quad B_n(x) = \sum_{w \in \mathcal{B}_n} x^{\text{des}^B(w)} = \sum_{w \in \mathcal{B}_n} x^{\text{des}_B(w)},$$

where

- $\text{des}^B(w)$ is the number of indices $i \in \{0, 1, \dots, n-1\}$ such that $w(i) > w(i+1)$,
- $\text{des}_B(w)$ is the number of indices $i \in \{0, 1, \dots, n-1\}$ such that $w(i) >_r w(i+1)$

for $w \in \mathcal{B}_n$, with $w(0) := 0$. For the first few values of n , we have

$$B_n(x) = \begin{cases} 1 + x, & \text{if } n = 1, \\ 1 + 6x + x^2, & \text{if } n = 2, \\ 1 + 23x + 23x^2 + x^3, & \text{if } n = 3, \\ 1 + 76x + 230x^2 + 76x^3 + x^4, & \text{if } n = 4, \\ 1 + 237x + 1682x^2 + 1682x^3 + 237x^4 + x^5, & \text{if } n = 5, \\ 1 + 722x + 10543x^2 + 23548x^3 + 10543x^4 + 722x^5 + x^6, & \text{if } n = 6. \end{cases}$$

A *descending run* of a permutation $w \in \mathfrak{S}_n$ is any maximal string $\{a, a+1, \dots, b\}$ of integers such that $w(a) > w(a+1) > \dots > w(b)$. A *left peak* of w is any index $i \in [n-1]$ such that $w(i-1) < w(i) > w(i+1)$, where $w(0) := 0$ (note that 1 can be a left peak, but n cannot). The following result combines [39, Theorem 4.7] with [94, Proposition 4.15] and provides a \mathcal{B}_n -analog to Theorem 2.1. The permutation obtained from $w \in \mathcal{B}_n$ by forgetting all signs is denoted by $|w|$.

Theorem 2.10 ([39, 94]). *For all $n \geq 1$,*

$$(19) \quad B_n(x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \gamma_{n,i}^B x^i (1+x)^{n-2i},$$

where $\gamma_{n,i}^B$ is equal to each of the following:

- the number of permutations $w \in \mathfrak{S}_n$ with i left peaks, multiplied by 4^i ,
- the number of signed permutations $w \in \mathcal{B}_n$ with $\text{des}^B(w) = i$, such that $|w| \in \mathfrak{S}_n$ has i descending runs of size at least two.

In particular,

$$(20) \quad B_n(-1) = \begin{cases} 0, & \text{if } n \text{ is odd,} \\ (-1)^{n/2} \gamma_{n,n/2}^B, & \text{if } n \text{ is even,} \end{cases}$$

for all n , where $\gamma_{n,n/2}^B$ is the number of up-down permutations in \mathfrak{S}_n , multiplied by $4^{n/2}$.

For the first few values of n , the numbers $\gamma_{n,i}^B$ are determined from the expressions

$$B_n(x) = \begin{cases} 1 + x, & \text{if } n = 1, \\ (1+x)^2 + 4x, & \text{if } n = 2, \\ (1+x)^3 + 20x(1+x), & \text{if } n = 3, \\ (1+x)^4 + 72x(1+x)^2 + 80x^2, & \text{if } n = 4, \\ (1+x)^5 + 232x(1+x)^3 + 976x^2(1+x), & \text{if } n = 5, \\ (1+x)^6 + 716x(1+x)^4 + 7664x^2(1+x)^2 + 3904x^3, & \text{if } n = 6. \end{cases}$$

An interesting extension of Theorem 2.8 to affine Weyl groups was found by K. Dilks, T.K. Petersen and J. Stembridge [48]. Let (W, S) be a Coxeter system with Coxeter length function $\ell_S : W \rightarrow \mathbb{N}$, as before, and assume W is finite, crystallographic and irreducible.

Then, W has a longest element s_0 and the set of *affine right descents* of $w \in W$ is defined as

$$\text{aDes}(w) := \begin{cases} \text{Des}(w), & \text{if } \ell_S(ws_0) < \ell_S(w), \\ \text{Des}(w) \cup \{s_0\}, & \text{if } \ell_S(ws_0) > \ell_S(w). \end{cases}$$

The *affine Eulerian polynomial* associated to W is defined as

$$(21) \quad W_a(x) := \sum_{w \in W} x^{\text{ades}(w)},$$

where $\text{ades}(w)$ is the number of affine right descents of $w \in W$.

Theorem 2.11 (DILKS–PETERSEN–STEMBRIDGE [48, Theorem 4.2]). *The affine Eulerian polynomial $W_a(x)$ is γ -positive for every finite irreducible crystallographic Coxeter group W .*

We close this section with an intriguing related problem, posed by I. Gessel in 2005 (see [26, Conjecture 10.2] [96, Conjecture 1] [97, Problem 4.12]). The *two-sided Eulerian polynomial* associated to a finite Coxeter group W is defined as

$$(22) \quad W(x, y) = \sum_{w \in W} x^{\text{des}(w)} y^{\text{des}(w^{-1})}.$$

The specialization $W(x, x)$ appeared also in work of A. Hultman [71, Example 5.9]. The following result was conjectured by Gessel (unpublished) for the symmetric groups and, more generally, by Petersen [98, Conjecture 1] for finite Coxeter groups.

Theorem 2.12 (LIN [80]). *Let W be a symmetric or hyperoctahedral group. Then, there exist nonnegative integers $\gamma_{i,j} = \gamma_{i,j}(W)$ such that*

$$(23) \quad W(x, y) = \sum_{2i+j \leq n} \gamma_{i,j}(xy)^i (x+y)^j (1+xy)^{n-2i-j},$$

where n is the rank of W .

It remains an interesting open problem to find a combinatorial interpretation for the numbers $\gamma_{i,j}(W)$, even in the symmetric group case.

2.1.4. *Derangements.* Counting derangements (permutations without fixed points) in \mathfrak{S}_n by the number of excedances leads to a well-behaved analog of the Eulerian polynomial $A_n(x)$, defined by

$$(24) \quad d_n(x) := \sum_{w \in \mathcal{D}_n} x^{\text{exc}(w)},$$

where \mathcal{D}_n denotes the set of all derangements in \mathfrak{S}_n . For the first few values of n , we have

$$d_n(x) = \begin{cases} 0, & \text{if } n = 1, \\ x, & \text{if } n = 2, \\ x + x^2, & \text{if } n = 3, \\ x + 7x^2 + x^3, & \text{if } n = 4, \\ x + 21x^2 + 21x^3 + x^4, & \text{if } n = 5, \\ x + 51x^2 + 161x^3 + 51x^4 + x^5, & \text{if } n = 6, \\ x + 113x^2 + 813x^3 + 813x^4 + 113x^5 + x^6, & \text{if } n = 7. \end{cases}$$

The polynomial $d_n(x)$ (often called the *n*th derangement polynomial) was first considered in a purely combinatorial context in [122, p. 530] by Stanley who, however, seems to have been motivated by a geometric interpretation [123, Proposition 2.4] of $d_n(x)$; see Section 3.3 for more explanation. While the symmetry of $d_n(x)$ is nearly obvious, its unimodality was derived by Brenti [30, Corollary 1] from a more general result in the theory of symmetric functions, although it also follows from deep results of Stanley [123] on local h -polynomials, discussed in Section 3.2. A more elementary combinatorial proof was later given by Stembridge [131, Section 2]. More recently, using methods discussed in Section 4.2, M. Juhnke-Kubitzke, S. Murai and R. Sieg [73, Corollary 4.2] found the recurrence

$$(25) \quad d_n(x) = \sum_{k=0}^{n-2} \binom{n}{k} d_k(x)(x + x^2 + \cdots + x^{n-1-k}),$$

which directly implies the unimodality of $d_n(x)$ by induction on n .

The question of γ -positivity of $d_n(x)$ arises naturally. Just as is the case with $A_n(x)$, the polynomial $d_n(x)$ turns out to be real-rooted for all n [142], so the interesting part of the question is to find a proof of γ -positivity which provides a combinatorial interpretation for the γ -coefficients. For the first few values of n , we have

$$d_n(x) = \begin{cases} x, & \text{if } n = 2, \\ x(1+x), & \text{if } n = 3, \\ x(1+x)^2 + 5x^2, & \text{if } n = 4, \\ x(1+x)^3 + 18x^2(1+x), & \text{if } n = 5, \\ x(1+x)^4 + 47x^2(1+x)^2 + 61x^3, & \text{if } n = 6, \\ x(1+x)^5 + 108x^2(1+x)^3 + 479x^3(1+x), & \text{if } n = 7. \end{cases}$$

For a permutation $w \in \mathfrak{S}_n$, a *double descent* is any index $2 \leq i \leq n-1$ such that $w(i-1) > w(i) > w(i+1)$; a *left to right maximum* is any index $1 \leq j \leq n$ such that $w(i) < w(j)$ for all $1 \leq i < j$.

Theorem 2.13 (cf. [10, Theorem 1.4]). *We have $d_n(x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \xi_{n,i} x^i (1+x)^{n-2i}$, where $\xi_{n,i}$ is equal to each of the following:*

- the number of derangements $w \in \mathcal{D}_n$ with i excedances and no double excedance,

- the number of $w \in \mathfrak{S}_n$ for which $\text{Asc}(w) \in \text{Stab}([2, n-2])$ has $i-1$ elements,
- the number of $w \in \mathfrak{S}_n$ for which $\text{Des}(w) \in \text{Stab}([2, n-2])$ has $i-1$ elements,
- the number of permutations $w \in \mathfrak{S}_n$ with i descents and no double descent, such that every left to right maximum of w is a descent.

In particular, $d_n(x)$ is γ -positive and

$$(26) \quad d_n(-1) = \begin{cases} 0, & \text{if } n \text{ is odd,} \\ (-1)^{n/2} \xi_{n,n/2}, & \text{if } n \text{ is even,} \end{cases}$$

for all n , where $\xi_{n,n/2}$ is the number of up-down permutations in \mathfrak{S}_n .

The first three interpretations appeared (implicitly or explicitly) in various contexts in independent works by several authors, roughly at the same time, who employed different methods; see [10, Equations (1.3) and (3.2)] [84, Section 4] [111, Section 5] [114, Section 6] [115] [137]. These references provide various interesting refinements, some of which are described in the sequel. A symmetric function generalization is discussed in Section 2.5.

Three refinements of the first interpretation given in Theorem 2.13 were found by H. Shin and J. Zeng [115, 116]. We denote by $c(w)$ the number of cycles of $w \in \mathfrak{S}_n$ and note that the meanings of $\text{inv}(w)$ and $\text{nest}(w)$ have been explained earlier, before the statement of Theorem 2.2. The result in the following theorem about $\text{nest}(w)$ is the specialization $q = 1$ of [115, Corollary 9], the one about $c(w)$ (which also follows from the proof of Theorem 2.13 given in [10, Section 4]) is a restatement of [115, Theorem 11] and the one about $\text{inv}(w)$ appears as [116, Theorem 2].

Theorem 2.14 (SHIN–ZENG [115, 116]). *For all positive integers n and for each of the statistics $\text{stat}(w) \in \{c(w), \text{inv}(w), \text{nest}(w)\}$,*

$$(27) \quad \sum_{w \in \mathcal{D}_n} q^{\text{stat}(w)} x^{\text{exc}(w)} = \sum_{i=0}^{\lfloor n/2 \rfloor} b_{n,i}(q) x^i (1+x)^{n-2i},$$

where

$$(28) \quad b_{n,i}(q) = \sum_{w \in \mathcal{D}_n(i)} q^{\text{stat}(w)}$$

and $\mathcal{D}_n(i)$ consists of all elements of \mathcal{D}_n with exactly i excedances and no double excedance.

The following theorem refines the third interpretation of the γ -positivity of $d_n(x)$, given in Theorem 2.13. This result was stated in [111, Remark 5.5] without giving an explicit interpretation for the coefficients $\xi_i(p, q)$. The combinatorial interpretation which appears here follows from the methods used to prove the specialization $p = 1$ in [114, Theorem 6.1]; the proof will be sketched in Section 4.3. The special case $p = 1$ is also implicit in [84] (see Equation (1.3) and Corollary 3.7 there) and is proven by different methods in [83].

Theorem 2.15 (SHARESHIAN–WACHS [111, Remark 5.5] [114, Section 6]). *For all $n \geq 1$,*

$$(29) \quad \sum_{w \in \mathcal{D}_n} p^{\text{des}(w)} q^{\text{maj}(w) - \text{exc}(w)} x^{\text{exc}(w)} = \sum_{i=0}^{\lfloor n/2 \rfloor} \xi_{n,i}(p, q) x^i (1+x)^{n-2i},$$

where

$$(30) \quad \xi_{n,i}(p, q) = p \cdot \sum_w p^{\text{des}(w^{-1})} q^{\text{maj}(w^{-1})},$$

and the sum runs through all permutations $w \in \mathfrak{S}_n$ for which $\text{Des}(w) \in \text{Stab}([2, n-2])$ has $i-1$ elements.

The following result, which is derived using the same methods in Section 4.3, generalizes the specialization $p = 1$ of Theorem 2.15. We denote by $\text{fix}(w)$ the number of fixed points of $w \in \mathfrak{S}_n$.

Theorem 2.16 (SHARESHIAN–WACHS [111, Corollary 4.6] [114, Theorem 6.1]). *For all $n \geq 1$ and $0 \leq k \leq n$,*

$$(31) \quad \sum_{w \in \mathfrak{S}_n: \text{fix}(w)=k} q^{\text{maj}(w) - \text{exc}(w)} x^{\text{exc}(w)} = \binom{n}{k}_q \sum_{i=0}^{\lfloor (n-k)/2 \rfloor} \xi_{n-k,i}(q) x^i (1+x)^{n-k-2i},$$

where $\binom{n}{k}_q$ is a q -binomial coefficient, $\xi_{n,i}(q) := \xi_{n,i}(1, q)$ and $\xi_{n,i}(p, q)$ is as in the statement of Theorem 2.15.

Problem 2.17. *Find a p -analog of Theorem 2.16 which reduces to Theorem 2.15 for $k = 0$.*

For a generalization of Theorem 2.13 to r -colored permutations and some related results, see Section 2.1.7. We now briefly describe an application to the γ -positivity of binomial Eulerian polynomials.

Remark 2.18. By the symmetry $\tilde{A}_n(x) = x^n \tilde{A}_n(1/x)$ of the binomial Eulerian polynomial $\tilde{A}_n(x)$, its defining equation (11) can be rewritten as

$$\tilde{A}_n(x) = \sum_{m=0}^n \binom{n}{m} x^{n-m} A_m(x).$$

Replacing $A_m(x)$ by the expression $\sum_{k=0}^m \binom{m}{k} d_k(x)$ and changing the order of summation, we obtain the formula

$$(32) \quad \tilde{A}_n(x) = \sum_{k=0}^n \binom{n}{k} d_k(x) (1+x)^{n-k}.$$

Using the γ -expansion of Theorem 2.13 for the derangement polynomials, we conclude that (14) holds with

$$(33) \quad \tilde{\gamma}_{n,i} = \sum_{k=2i}^n \binom{n}{k} \xi_{k,i}.$$

Because of the third interpretation for the coefficients $\xi_{n,i}$ given in Theorem 2.13, this formula is equivalent to the combinatorial interpretation for $\tilde{\gamma}_{n,i}$ predicted in [102]. This approach can be generalized in the context of r -colored permutations; details will appear elsewhere. \square

2.1.5. *Involutions.* Let $\mathcal{I}_n := \{w \in \mathfrak{S}_n : w^{-1} = w\}$ be the set of involutions in \mathfrak{S}_n and

$$(34) \quad I_n(x) = \sum_{w \in \mathcal{I}_n} x^{\text{des}(w)}$$

be the polynomial defining the Eulerian distribution on \mathcal{I}_n . For the first few values of n , we have

$$I_n(x) = \begin{cases} 1, & \text{if } n = 1, \\ 1 + x, & \text{if } n = 2, \\ 1 + 2x + x^2, & \text{if } n = 3, \\ 1 + 4x + 4x^2 + x^3, & \text{if } n = 4, \\ 1 + 6x + 12x^2 + 6x^3 + x^4, & \text{if } n = 5, \\ 1 + 9x + 28x^2 + 28x^3 + 9x^4 + x^5, & \text{if } n = 6, \\ 1 + 12x + 57x^2 + 92x^3 + 57x^4 + 12x^5 + x^6, & \text{if } n = 7. \end{cases}$$

The polynomial $I_n(x)$ seems to have first appeared in [136], where V. Strehl proved its symmetry (conjectured by D. Dumont). Strehl’s argument uses basic properties of the Robinson–Schensted correspondence to derive the alternative formula

$$(35) \quad I_n(x) = \sum_{Q \in \text{SYT}(n)} x^{\text{des}(Q)},$$

where $\text{SYT}(n)$ stands for the set of all standard Young tableaux with n squares and $\text{des}(Q)$ is the numbers of entries (called descents) $i \in [n - 1]$ such that $i + 1$ appears in Q in a lower row than i , for $Q \in \text{SYT}(n)$. This formula makes the symmetry of $I_n(x)$ apparent, since transposing a standard Young tableau interchanges descents with ascents (nondescents).

Two basic results about $I_n(x)$ are as follows. The proof of the second part, given in [68], uses the first part to find recursions for the coefficients of $I_n(x)$ and proceeds by induction on n .

Theorem 2.19. (a) (DÉSARMÉNIEN–FOATA [46]) *We have*

$$(36) \quad \sum_{n \geq 0} \frac{I_n(x)}{(1-x)^{n+1}} z^n = \sum_{m \geq 0} \frac{x^m}{(1-z)^{m+1}(1-z^2)^{m(m+1)/2}},$$

where $I_0(x) := 1$.

(b) (GUO–ZENG [68]) *The polynomial $I_n(x)$ is (symmetric and) unimodal for every positive integer n .*

The polynomials $I_n(x)$ are not real-rooted (in fact, not even log-concave [13]) for n large enough, but the following data suggests they may be γ -positive:

$$I_n(x) = \begin{cases} 1, & \text{if } n = 1, \\ 1 + x, & \text{if } n = 2, \\ (1 + x)^2, & \text{if } n = 3, \\ (1 + x)^3 + x(1 + x), & \text{if } n = 4, \\ (1 + x)^4 + 2x(1 + x)^2 + 2x^2, & \text{if } n = 5, \\ (1 + x)^5 + 4x(1 + x)^3 + 6x^2(1 + x), & \text{if } n = 6, \\ (1 + x)^6 + 6x(1 + x)^4 + 18x^2(1 + x)^2, & \text{if } n = 7. \end{cases}$$

Somewhat surprisingly, the following intriguing conjecture is still open (an analogous statement for fixed-point free involutions is conjectured [68, Conjecture 4.3] for large n).

Conjecture 2.20 (GUO–ZENG [68, Conjecture 4.1]). *The polynomial $I_n(x)$ is γ -positive for every positive integer n .*

The previous conjecture suggests that it may be interesting to study the distribution of the descent set $\text{Des}(w)$ for $w \in \mathcal{I}_n$. For instance, for $S \subseteq [n-1]$ let $\beta_n(S)$ denote the number of permutations $w \in \mathfrak{S}_n$ such that $\text{Des}(w) = S$. It is well known (see [127, Section 1.6.3]) that $\beta_n(S) = \beta_n([n-1] \setminus S)$ for every $S \subseteq [n-1]$ and that for each $n \geq 1$, when S ranges over all subsets of $[n-1]$, the quantity $\beta_n(S)$ attains its maximum for $S = \{1, 3, 5, \dots\} \cap [n-1]$.

Similarly, for $S \subseteq [n-1]$ let $\delta_n(S)$ denote the number of involutions $w \in \mathcal{I}_n$ such that $\text{Des}(w) = S$. Strehl's proof of the symmetry of $I_n(x)$ shows that $\delta_n(S) = \delta_n([n-1] \setminus S)$ for every $S \subseteq [n-1]$.

Question 2.21. *Is it true that for each $n \geq 1$, when S ranges over all subsets of $[n-1]$, the quantity $\delta_n(S)$ attains its maximum for $S = \{1, 3, 5, \dots\} \cap [n-1]$?*

It is natural to consider the \mathcal{B}_n -analog

$$(37) \quad I_n^B(x) = \sum_{w \in \mathcal{I}_n^B} x^{\text{des}_B(w)}$$

of $I_n(x)$, where $\mathcal{I}_n^B := \{w \in \mathcal{B}_n : w^{-1} = w\}$ is the set of involutions in the hyperoctahedral group \mathcal{B}_n . Presumably (but not obviously), the right-hand side of the defining equation (37) is unaffected when des_B is replaced with the Coxeter group descent statistic des^B for \mathcal{B}_n . For the first few values of n , we have

$$I_n^B(x) = \begin{cases} 1 + x, & \text{if } n = 1, \\ 1 + 4x + x^2, & \text{if } n = 2, \\ 1 + 9x + 9x^2 + x^3, & \text{if } n = 3, \\ 1 + 17x + 40x^2 + 17x^3 + x^4, & \text{if } n = 4, \\ 1 + 28x + 127x^2 + 127x^3 + 28x^4 + x^5, & \text{if } n = 5, \\ 1 + 43x + 331x^2 + 634x^3 + 331x^4 + 43x^5 + x^6, & \text{if } n = 6. \end{cases}$$

The symmetry of $I_n^B(x)$ can be demonstrated by an analog of Strehl’s argument, using basic properties (see [1, Proposition 5.1]) of the Robinson–Schensted correspondence of type B and replacing $\text{SYT}(n)$ with the set of all standard Young bitableaux with a total of n squares; see [87, Section 3.2] for the details.

The analog

$$(38) \quad \sum_{n \geq 0} \frac{I_n^B(x)}{(1-x)^{n+1}} z^n = \sum_{m \geq 0} \frac{x^m}{(1-z)^{2m+1}(1-z^2)^{m^2}},$$

of Equation (36) for $I_n^B(x)$, where $I_0^B(x) := 1$, was found by V. Moustakas [87], who also showed that $I_n^B(x)$ is unimodal for every positive integer n . As expected, the following data suggests that $I_n^B(x)$ may be γ -positive for every n :

$$I_n^B(x) = \begin{cases} 1+x, & \text{if } n=1, \\ (1+x)^2+2x, & \text{if } n=2, \\ (1+x)^3+6x(1+x), & \text{if } n=3, \\ (1+x)^4+13x(1+x)^2+8x^2, & \text{if } n=4, \\ (1+x)^5+23x(1+x)^3+48x^2(1+x), & \text{if } n=5, \\ (1+x)^6+37x(1+x)^4+168x^2(1+x)^2+56x^3, & \text{if } n=6. \end{cases}$$

The formulas in the following proposition, which seem not to have appeared in the literature explicitly before, express $I_n(x)$ and $I_n^B(x)$ in terms of Eulerian polynomials of types A and B , respectively. We sketch their proofs, which we find interesting. We recall (see, for instance, [1, Section 2.3]) that the cycle form of elements of \mathcal{B}_n involves positive (paired) cycles and negative (balanced) cycles.

Proposition 2.22. *For $n \geq 1$,*

$$(39) \quad I_n(x) = \frac{1}{n!} \sum_{w \in \mathfrak{S}_n} (1-x)^{n-c(w^2)} A_{c(w^2)}(x),$$

$$(40) \quad I_n^B(x) = \frac{1}{2^n n!} \sum_{w \in \mathcal{B}_n} (1-x)^{n-c_+(w^2)} B_{c_+(w^2)}(x),$$

where $c(u)$ stands for the number of cycles of $u \in \mathfrak{S}_n$ and $c_+(v)$ stands for the number of positive cycles of $v \in \mathcal{B}_n$.

Proof. Recall from [125, Section 7.19] that the fundamental quasisymmetric function associated to $S \subseteq [n-1]$ is defined as

$$(41) \quad F_{n,S}(\mathbf{x}) = \sum_{\substack{1 \leq i_1 \leq i_2 \leq \dots \leq i_n \\ j \in S \Rightarrow i_j < i_{j+1}}} x_{i_1} x_{i_2} \cdots x_{i_n},$$

where $\mathbf{x} = (x_1, x_2, \dots)$ is a sequence of commuting independent indeterminates, and that

$$(42) \quad \sum_{m \geq 1} F_{n,S}(1^m) x^{m-1} = \frac{x^{|S|}}{(1-x)^{n+1}}.$$

Applying this equality for $S = \text{Des}(w)$, summing over all involutions $w \in \mathcal{I}_n$, changing the order of summation on the left-hand side and using the correspondence between involutions and standard Young tableaux, we get

$$\begin{aligned} \frac{I_n(x)}{(1-x)^{n+1}} &= \sum_{m \geq 1} \sum_{w \in \mathcal{I}_n} F_{n, \text{Des}(w)}(1^m) x^{m-1} = \sum_{m \geq 1} \sum_{Q \in \text{SYT}(n)} F_{n, \text{Des}(Q)}(1^m) x^{m-1} \\ &= \sum_{m \geq 1} \sum_{\lambda \vdash n} \sum_{Q \in \text{SYT}(\lambda)} F_{n, \text{Des}(Q)}(1^m) x^{m-1}. \end{aligned}$$

By using the well known expansion [125, Theorem 7.19.7]

$$(43) \quad s_\lambda(\mathbf{x}) = \sum_{Q \in \text{SYT}(\lambda)} F_{n, \text{Des}(Q)}(\mathbf{x})$$

of the Schur function $s_\lambda(\mathbf{x})$ associated to $\lambda \vdash n$, our formula for $I_n(x)$ may be rewritten as

$$\frac{I_n(x)}{(1-x)^{n+1}} = \sum_{m \geq 1} \sum_{\lambda \vdash n} s_\lambda(1^m) x^{m-1}.$$

Expanding now $s_\lambda(\mathbf{x})$ in the power-sum basis (see [125, Theorem 7.18.5]) and changing the order of summation, we obtain

$$\frac{I_n(x)}{(1-x)^{n+1}} = \frac{1}{n!} \sum_{m \geq 1} \sum_{\lambda \vdash n} \sum_{u \in \mathfrak{S}_n} \chi^\lambda(u) m^{c(u)} x^{m-1} = \frac{1}{n!} \sum_{u \in \mathfrak{S}_n} \sum_{m \geq 1} m^{c(u)} x^{m-1} \sum_{\lambda \vdash n} \chi^\lambda(u),$$

where χ^λ is the irreducible character of \mathfrak{S}_n corresponding to $\lambda \vdash n$. The desired expression (39) for $I_n(x)$ follows by applying Worpitzky's identity $\sum_{m \geq 1} m^k x^{m-1} = A_k(x)/(1-x)^{k+1}$ and the fact (see [72, p. 58]) that $\sum_{\lambda \vdash n} \chi^\lambda(u)$ is equal to the number of $w \in \mathfrak{S}_n$ satisfying $w^2 = u$, for every $u \in \mathfrak{S}_n$.

The proof of the expression (40) for $I_n^B(x)$ is similar, provided one uses Poirier's signed quasisymmetric functions [99] to replace the functions $F_{n,S}(\mathbf{x})$. In the sequel, we assume some familiarity with [1, Section 2], especially with the notion of signed descent set for signed permutations and standard Young bitableaux.

Let $\mathbf{x} = (x_1, x_2, \dots)$ and $\mathbf{y} = (y_1, y_2, \dots)$ be sequences of commuting independent indeterminates. Given $w \in \mathcal{B}_n$, let $\varepsilon = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n) \in \{-, +\}^n$ be the vector with i th coordinate equal to the sign of $w(i)$, and let $\text{Des}(w)$ be the set consisting of those indices $i \in [n-1]$ for which either $\varepsilon_i = +$ and $\varepsilon_{i+1} = -$, or $\varepsilon_i = \varepsilon_{i+1}$ and $|w(i)| > |w(i+1)|$. The signed quasisymmetric function associated (in a more general setting) to w by Poirier [99] may be defined as

$$(44) \quad F_w(\mathbf{x}, \mathbf{y}) = \sum_{\substack{1 \leq i_1 \leq i_2 \leq \dots \leq i_n \\ j \in \text{Des}(w) \Rightarrow i_j < i_{j+1}}} z_{i_1} z_{i_2} \cdots z_{i_n},$$

where $z_{i_j} = x_{i_j}$ if $\varepsilon_j = +$, and $z_{i_j} = y_{i_j}$ if $\varepsilon_j = -$; see [1, Sections 2.2 and 2.4]. There is a similar definition of $F_Q(\mathbf{x}, \mathbf{y})$ for every standard Young bitableau Q ; see [1, Section 2] for

a uniform treatment of these functions. One can check that

$$(45) \quad \sum_{m \geq 1} F_w(1^m, 01^{m-1})x^{m-1} = \frac{x^{\text{des}_B(w)}}{(1-x)^{n+1}},$$

where $F_w(1^m, 01^{m-1})$ stands for the specialization $x_1 = \cdots = x_m = y_2 = \cdots = y_m = 1$, $y_1 = 0$ and $x_i = y_i = 0$ for $i > m$ of the function $F_w(\mathbf{x}, \mathbf{y})$. Let $\text{SYB}(\lambda, \mu)$ denote the set of standard Young bitableaux of shape (λ, μ) . Following the proof for $I_n(x)$ described earlier and using the one-to-one correspondence between involutions in \mathcal{B}_n and standard Young bitableaux with a total of n squares, as well as the expansion

$$(46) \quad s_\lambda(\mathbf{x})s_\mu(\mathbf{y}) = \sum_{Q \in \text{SYB}(\lambda, \mu)} F_Q(\mathbf{x}, \mathbf{y})$$

of [1, Proposition 4.2] instead of (43), we arrive at the equation

$$\frac{I_n^B(x)}{(1-x)^{n+1}} = \sum_{m \geq 1} \sum_{(\lambda, \mu) \vdash n} s_\lambda(1^m)s_\mu(1^{m-1})x^{m-1}.$$

Finally, denote by $\chi^{\lambda, \mu}$ the irreducible \mathcal{B}_n -character associated to the bipartition (λ, μ) of n . One uses the characteristic map [1, Equation (2.5)] for \mathcal{B}_n to expand $s_\lambda(\mathbf{x})s_\mu(\mathbf{y}) = \text{ch}(\chi^{\lambda, \mu})$ in the power sum basis, the type B Worpitzky identity $\sum_{m \geq 1} (2m-1)^k x^{m-1} = B_k(x)/(1-x)^{k+1}$ and the fact (see again [72, p. 58]) that $\sum_{(\lambda, \mu) \vdash n} \chi^{\lambda, \mu}(v)$ is equal to the number of square roots of v in \mathcal{B}_n , for every $v \in \mathcal{B}_n$, to reach the desired conclusion. \square

Remark 2.23. Another proof of the symmetry of $I_n(x)$ and $I_n^B(x)$ can be inferred from Proposition 2.22 as follows. Replacing x by $1/x$ in the right-hand sides of Equations (39) and (40) and using the symmetry of the Eulerian polynomials $A_k(x)$ and $B_k(x)$, as well as the fact that $n - c(u^2)$ and $n - c_+(v^2)$ are even for all $u \in \mathfrak{S}_n$ and $v \in \mathcal{B}_n$, we see that $x^{n-1}I_n(1/x) = I_n(x)$ and $x^n I_n^B(1/x) = I_n^B(x)$, as desired. \square

The polynomial $I_n(x)$ affords a natural Coxeter group generalization. Given a finite Coxeter group W (together with a set of simple generators), let $\mathcal{I}_W := \{w \in W : w^{-1} = w\}$ be the set of involutions in W and define

$$(47) \quad I_W(x) := \sum_{w \in \mathcal{I}_W} x^{\text{des}(w)},$$

where, as in Equation (16), $\text{des}(w)$ is the number of right descents of w (the notions of left and right descents coincide for involutions). The polynomial $I_W(x)$ is equal to $I_n(x)$, when W is the symmetric group \mathfrak{S}_n , and presumably to $I_n^B(x)$, when W is the hyperoctahedral group \mathcal{B}_n .

The symmetry of $I_W(x)$ follows from the work of Hultman [71, Section 5], who showed (in a more general context) that $I_W(x)$ is equal to the h -polynomial of a Boolean cell complex which is homeomorphic to a sphere. Thus, it seems natural to ask the following question.

Question 2.24. *For which finite Coxeter groups W is $I_W(x)$ unimodal, or even γ -positive?*

Another possible generalization of $I_n(x)$ is provided by the polynomial

$$(48) \quad I_{n,k}(x) := \sum_{w \in \mathfrak{S}_n : w^k = e} x^{\text{des}(w)},$$

defined for positive integers k , where $e \in \mathfrak{S}_n$ stands for the identity permutation. This polynomial is no longer symmetric for $k \geq 3$, but it may still be worthwhile to investigate its unimodality. Following the proof of the first part of Proposition 2.22 and using the main result of [106], in the equivalent form stated in [1, Theorem 7.7], to evaluate the quasisymmetric generating function of the descent set over all k -roots of $e \in \mathfrak{S}_n$, we obtain in the generalization

$$(49) \quad I_{n,k}(x) = \frac{1}{n!} \sum_{w \in \mathfrak{S}_n} (1-x)^{n-c(w^k)} A_{c(w^k)}(x)$$

of Equation (39).

2.1.6. Multiset permutations. Let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k) \vdash n$ be an integer partition of n . We denote by M_λ the multiset consisting of λ_i copies of i for each $i \in [k]$ and by $\mathfrak{S}(M_\lambda)$ the set of all permutations of M_λ , written in one-line notation. The sets of *ascents*, *descents* and *excedances* of $w = (a_1, a_2, \dots, a_n) \in \mathfrak{S}(M_\lambda)$ are defined as

- $\text{Asc}(w) := \{i \in [n-1] : a_i \leq a_{i+1}\}$,
- $\text{Des}(w) := \{i \in [n-1] : a_i > a_{i+1}\}$,
- $\text{Exc}(w) := \{i \in [n-1] : a_i > j_i\}$,

where (j_1, j_2, \dots, j_n) is the unique permutation of M_λ with no descents; the cardinalities of these sets are denoted by $\text{asc}(w)$, $\text{des}(w)$ and $\text{exc}(w)$, respectively. We say that w is a *Smirnov permutation* of type λ if it has no two equal successive entries, and a *derangement* of type λ if $a_i \neq j_i$ for all indices $i \in [n]$. We denote by \mathcal{S}_λ and \mathcal{D}_λ the set of all Smirnov permutations and derangements of type λ , respectively.

The following result, which is a consequence of [84, Theorem 5.1] and its proof, generalizes Theorems 2.1 and 2.13 in the context of permutations of multisets.

Theorem 2.25 (LINUSSON–SHARESHIAN–WACHS [84, Section 5]). *For all partitions $\lambda \vdash n$,*

$$(50) \quad \sum_{w \in \mathcal{S}_\lambda} x^{\text{des}(w)} = \sum_{i=0}^{\lfloor (n-1)/2 \rfloor} \gamma_{\lambda,i} x^i (1+x)^{n-1-2i}$$

and

$$(51) \quad \sum_{w \in \mathcal{D}_\lambda} x^{\text{exc}(w)} = \sum_{i=0}^{\lfloor n/2 \rfloor} \xi_{\lambda,i} x^i (1+x)^{n-2i},$$

where

- $\gamma_{\lambda,i}$ is the number of permutations $w \in \mathfrak{S}(M_\lambda)$ for which $\text{Asc}(w) \in \text{Stab}([n-2])$ has i elements, and
- $\xi_{\lambda,i}$ is the number of permutations $w \in \mathfrak{S}(M_\lambda)$ for which $\text{Asc}(w) \in \text{Stab}([2, n-2])$ has $i-1$ elements.

2.1.7. *Colored permutations.* Theorem 2.13 has been generalized to the wreath product group $\mathbb{Z}_r \wr \mathfrak{S}_n$. Recall that the elements of $\mathbb{Z}_r \wr \mathfrak{S}_n$ can be viewed as r -colored permutations of the form $\sigma \times \mathbf{z}$, where $\sigma = (\sigma(1), \sigma(2), \dots, \sigma(n)) \in \mathfrak{S}_n$ and $\mathbf{z} = (z_1, z_2, \dots, z_n) \in \{0, 1, \dots, r-1\}^n$ (the number z_i is thought of as the color assigned to $\sigma(i)$). To define the notions of descent and excedance for colored permutations, we follow [129, Section 2] [130, Section 3.1]. A *descent* of $\sigma \times \mathbf{z} \in \mathbb{Z}_r \wr \mathfrak{S}_n$ is any index $i \in [n]$ such that either $z_i > z_{i+1}$, or $z_i = z_{i+1}$ and $\sigma(i) > \sigma(i+1)$, where $\sigma(n+1) := n+1$ and $z_{n+1} := 0$ (in particular, n is a descent if and only if $\sigma(n)$ has nonzero color). An *excedance* of $\sigma \times \mathbf{z}$ is any index $i \in [n]$ such that either $\sigma(i) > i$, or $\sigma(i) = i$ and $z_i > 0$. Let $\text{des}(w)$ and $\text{exc}(w)$ be the number of descents and excedances, respectively, of $w \in \mathbb{Z}_r \wr \mathfrak{S}_n$. The Eulerian polynomial

$$(52) \quad A_{n,r}(x) := \sum_{w \in \mathbb{Z}_r \wr \mathfrak{S}_n} x^{\text{des}(w)} = \sum_{w \in \mathbb{Z}_r \wr \mathfrak{S}_n} x^{\text{exc}(w)}$$

for $\mathbb{Z}_r \wr \mathfrak{S}_n$ (the second equality being the content of [129, Theorem 3.15] [130, Theorem 15]) was defined and studied by E. Steingrímsson [129, 130], who showed it to be real-rooted for all n, r ; it reduces to the Eulerian polynomials $A_n(x)$ and $B_n(x)$ when $r = 1$ and $r = 2$, respectively. The derangement polynomial for $\mathbb{Z}_r \wr \mathfrak{S}_n$ is defined as

$$(53) \quad d_{n,r}(x) := \sum_{w \in \mathcal{D}_{n,r}} x^{\text{exc}(w)},$$

where $\mathcal{D}_{n,r}$ denotes the set of all derangements (colored permutations without fixed points of zero color) in $\mathbb{Z}_r \wr \mathfrak{S}_n$. This polynomial was introduced and studied by C. Chow and T. Mansour [41], who showed that it is real-rooted for all positive integers n, r ; it reduces to $d_n(x)$ for $r = 1$. For $r = 2$ it was first studied by W. Chen, R. Tang and A. Zhao [38] and, independently, by C. Chow [40]. For the first few values of n , we have

$$d_{n,2}(x) = \begin{cases} x, & \text{if } n = 1, \\ 4x + x^2, & \text{if } n = 2, \\ 8x + 20x^2 + x^3, & \text{if } n = 3, \\ 16x + 144x^2 + 72x^3 + x^4, & \text{if } n = 4, \\ 32x + 752x^2 + 1312x^3 + 232x^4 + x^5, & \text{if } n = 5, \\ 64x + 3456x^2 + 14576x^3 + 9136x^4 + 716x^5 + x^6, & \text{if } n = 6. \end{cases}$$

Even though $d_{n,r}(x)$ and $A_{n,r}(x)$ are no longer symmetric for $r \geq 2$ and $r \geq 3$, respectively, the theory of γ -positivity is still relevant to their study. Indeed, the following theorem shows that $d_{n,r}(x)$ is equal to the sum of two γ -positive, hence symmetric and unimodal, polynomials whose centers of symmetry differ by $1/2$ and thus implies its unimodality. We denote by $\text{Des}(w)$ the set of descents of $w \in \mathbb{Z}_r \wr \mathfrak{S}_n$ and set $\text{Asc}(w) := [n] \setminus \text{Des}(w)$.

For $r = 1$, these notions differ from the ones already defined for permutations $w \in \mathfrak{S}_n$ only in the fact that $\text{Asc}(w)$ is now forced to contain n . We call $w \in \mathbb{Z}_r \wr \mathfrak{S}_n$ *down-up*, if $\text{Des}(w) = \{1, 3, 5, \dots\} \cap [n]$.

Theorem 2.26 (ATHANASIADIS [6, Theorem 1.3 and Corollary 6.1]). *For all positive integers n, r ,*

$$(54) \quad d_{n,r}(x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \xi_{n,r,i}^+ x^i (1+x)^{n-2i} + \sum_{i=0}^{\lfloor (n+1)/2 \rfloor} \xi_{n,r,i}^- x^i (1+x)^{n+1-2i},$$

where $\xi_{n,r,i}^+$ is the number of permutations $w \in \mathbb{Z}_r \wr \mathfrak{S}_n$ for which $\text{Asc}(w) \in \text{Stab}([2, n])$ has i elements and contains n , and $\xi_{n,r,i}^-$ is the number of permutations $w \in \mathbb{Z}_r \wr \mathfrak{S}_n$ for which $\text{Asc}(w) \in \text{Stab}([2, n-1])$ has $i-1$ elements.

In particular, $d_{n,r}(x)$ is unimodal with a peak at $\lfloor (n+1)/2 \rfloor$. Moreover, if $r \geq 2$, then $(-1)^{\lfloor \frac{n+1}{2} \rfloor} d_{n,r}^*(-1)$ is equal to the number of down-up colored permutations in $\mathbb{Z}_r \wr \mathfrak{S}_n$.

The following question seems natural to ask.

Question 2.27. *Is there a positive expansion for $A_{n,r}(x)$ of the type provided by Equation (54) for $d_{n,r}(x)$? Is there a (p, q) -analog which reduces to Theorem 2.3 for $r = 1$?*

Problem 2.28. *Find a (p, q) -analog of Theorem 2.26 which reduces to Theorem 2.15 for $r = 1$.*

The proof of Theorem 2.26 uses a general result of Linusson, Shareshian and Wachs [84, Corollary 3.8] on the Möbius function of the Rees product for posets, which we discuss in Section 2.4.

Problem 2.29. *Find a direct combinatorial proof of Theorem 2.26.*

In the case $r = 2$, further discussed in the sequel, a different combinatorial interpretation for the numbers $\xi_{n,2,i}^+$ and $\xi_{n,2,i}^-$, in terms of permutations with no double excedance, was found by Shin and Zeng [116, Section 2]. This suggests the problem to find a combinatorial interpretation for the numbers $\xi_{n,r,i}^+$ and $\xi_{n,r,i}^-$ which generalizes the first interpretation of the numbers $\xi_{n,i}$, stated in Theorem 2.13, and prove directly its equivalence to the interpretation of Theorem 2.26.

Let us denote by $d_{n,r}^+(x)$ and $d_{n,r}^-(x)$ the first and second summand, respectively, in the right-hand side of (54). A geometric interpretation of the former will be discussed in Section 3.3. The following statement appeared as [108, Conjecture 3.7.10] in the special case $r = 2$; see also [7, Question 4.11].

Conjecture 2.30. *The polynomials $d_{n,r}^+(x)$ and $d_{n,r}^-(x)$ are real-rooted for all positive integers n, r .*

We close our discussion with generalizations of $A_n(x)$ and $d_n(x)$ to r -colored permutations which are different from, but related to (especially for $r = 2$), $A_{n,r}(x)$ and $d_{n,r}(x)$.

The *flag excedance* of a colored permutation $w = \sigma \times \mathbf{z} \in \mathbb{Z}_r \wr \mathfrak{S}_n$ was defined by E. Bagno and D. Garber [12] as

$$(55) \quad \text{fexc}(w) = r \cdot \text{exc}(\sigma) + \sum_{i=1}^n z_i,$$

where the sum of the colors z_i takes place in \mathbb{Z} . The generating polynomial of fexc over all r -colored permutations factors nicely [11, 54] (see also [6, Proposition 2.2]) as

$$(56) \quad A_{n,r}^{\text{fexc}}(x) := \sum_{w \in \mathbb{Z}_r \wr \mathfrak{S}_n} x^{\text{fexc}(w)} = (1 + x + x^2 + \cdots + x^{r-1})^n A_n(x).$$

The polynomial

$$(57) \quad f_{n,r}(x) := \sum_{w \in \mathcal{D}_{n,r}} x^{\text{fexc}(w)}$$

was studied by P. Mongelli [86, Section 3] in the case $r = 2$ and by Z. Lin [79, Section 2.4.1] and by H. Shin and J. Zeng [116] for any $r \geq 1$; it reduces to $d_n(x)$ for $r = 1$. For $r = 2$ and for the first few values of n , we have

$$f_{n,2}(x) = \begin{cases} x, & \text{if } n = 1, \\ x + 3x^2 + x^3, & \text{if } n = 2, \\ x + 7x^2 + 13x^3 + 7x^4 + x^5, & \text{if } n = 3, \\ x + 15x^2 + 57x^3 + 87x^4 + 57x^5 + 15x^6 + x^7, & \text{if } n = 4, \\ x + 31x^2 + 201x^3 + 551x^4 + 761x^5 + 551x^6 + 201x^7 + 31x^8 + x^9, & \text{if } n = 5. \end{cases}$$

The symmetry of $f_{n,r}(x)$ is nearly obvious (see [6, Proposition 2.5]); its unimodality was shown in [79, Theorem 2.4.11] [116, Corollary 4] (where the first reference offers a generalization and q -analog as well). The connection between the generating polynomials for exc and fexc , respectively, over $\mathbb{Z}_r \wr \mathfrak{S}_n$ and $\mathcal{D}_{n,r}$ are the formulas

$$(58) \quad A_{n,r}(x) = \tilde{\mathbb{E}}_r(A_{n,r}^{\text{fexc}}(x)),$$

$$(59) \quad d_{n,r}(x) = \tilde{\mathbb{E}}_r(f_{n,r}(x)),$$

where $\tilde{\mathbb{E}}_r : \mathbb{R}[x] \rightarrow \mathbb{R}[x]$ is the linear operator defined by setting $\tilde{\mathbb{E}}_r(x^m) = x^{\lceil m/r \rceil}$ for $m \in \mathbb{N}$; see [6, Proposition 2.3] and its proof. Equation (59) implies (see [6, Section 5]) that

$$(60) \quad d_{n,r}^+(x) = \sum_{w \in (\mathcal{D}_{n,r})^b} x^{\text{fexc}(w)/r},$$

$$(61) \quad d_{n,r}^-(x) = \sum_{w \in \mathcal{D}_{n,r} \setminus (\mathcal{D}_{n,r})^b} x^{\lceil \frac{\text{fexc}(w)}{r} \rceil},$$

where $(\mathcal{D}_{n,r})^b$ stands for the set of derangements $w \in \mathcal{D}_{n,r}$ such that $\text{fexc}(w)$ is divisible by r (colored permutations with this property are sometimes called *balanced*).

The polynomial $f_{n,r}(x)$ is not γ -positive for $r \geq 3$, the case $r = 2$ being more interesting. As discussed in [6, Section 5], it follows from Equations (60) and (61) that $d_{n,2}(x)$ and $f_{n,2}(x)$ determine one another in a simple way. The γ -positivity of $f_{n,2}(x)$ (which is not real-rooted for all n) follows from the formula in [86, Proposition 3.4]

$$(62) \quad f_{n,2}(x) = \sum_{k=0}^n \binom{n}{k} x^k (1+x)^{n-k} d_{n-k}(x)$$

and the γ -positivity of $d_n(x)$ (Mongelli [86, Conjecture 8.1] further conjectured that $f_{n,2}(x)$ is log-concave for all n). The corresponding γ -coefficients for the first few values of n are given by

$$f_{n,2}(x) = \begin{cases} x, & \text{if } n = 1, \\ x(1+x)^2 + x^2, & \text{if } n = 2, \\ x(1+x)^4 + 3x^2(1+x)^2 + x^3, & \text{if } n = 3, \\ x(1+x)^6 + 9x^2(1+x)^4 + 6x^3(1+x)^2 + x^4, & \text{if } n = 4, \\ x(1+x)^8 + 23x^2(1+x)^6 + 35x^3(1+x)^4 + 10x^4(1+x)^2 + x^5, & \text{if } n = 5. \end{cases}$$

The following elegant combinatorial interpretation for these coefficients was found by Shin and Zeng [116].

Theorem 2.31 (SHIN–ZENG [116, Cor. 5]). *We have $f_{n,2}(x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \hat{\xi}_{n,i} x^i (1+x)^{2n-2i}$, where $\hat{\xi}_{n,i}$ is equal to the number of permutations $w \in \mathfrak{S}_n$ with i weak excedances and no double excedance. In particular, $f_{n,2}(-1) = (-1)^n$ for $n \geq 1$.*

2.2. Coxeter–Narayana polynomials. Let W be a finite Coxeter group with set of simple generators S , and let $\ell_T : W \rightarrow \mathbb{N}$ be the length (known as *absolute length*) function with respect to the generating set $T = \{wsw^{-1} : s \in S, w \in W\}$ of all reflections in W . Choose a Coxeter element $c \in W$ and set

$$\text{NC}_W = \text{NC}_W(c) := \{w \in W : \ell_T(w) + \ell_T(w^{-1}c) = \ell_T(c)\}.$$

This set, endowed with a natural partial order which is graded by absolute length, was introduced by D. Bessis [16] and independently by T. Brady and C. Watt [23] as the *poset of noncrossing partitions* associated to W ; see [3, Chapter 2] for more information (the isomorphism type of this poset does not depend on the choice of c). The generating polynomial

$$(63) \quad \text{Cat}(W, x) := \sum_{w \in \text{NC}_W} x^{\ell_T(w)}$$

of the absolute length function over NC_W admits numerous algebraic, combinatorial and geometric interpretations and plays an important role in the subject of Catalan combinatorics of Coxeter groups [3, Chapter 5]. For the symmetric group \mathfrak{S}_n , its coefficients,

known as *Narayana numbers*, refine the n th Catalan number; explicitly, we have:

$$(64) \quad \text{Cat}(W, x) = \begin{cases} \sum_{i=0}^n \frac{1}{i+1} \binom{n}{i} \binom{n+1}{i} x^i, & \text{if } W = \mathfrak{S}_{n+1}, \\ \sum_{i=0}^n \binom{n}{i}^2 x^i, & \text{if } W = \mathcal{B}_n \\ \sum_{i=0}^n \binom{n}{i} \left(\binom{n-1}{i} + \binom{n-2}{i-2} \right) x^i, & \text{if } W = \mathcal{B}_n^e, \end{cases}$$

where (to avoid confusion with our notation for the set of derangements in \mathfrak{S}_n) \mathcal{B}_n^e stands for the group of even-signed permutations of Ω_n (which is a Coxeter group of type D_n).

The symmetry of $\text{Cat}(W, x)$ is a simple consequence of the definition. The following result can be verified with case-by-case computations; it is due essentially to R. Simion and D. Ullman [117, Corollary 3.1] [102, Proposition 11.14] for the group \mathfrak{S}_n , to A. Postnikov, V. Reiner and L. Williams [102, Proposition 11.15] for the group \mathcal{B}_n and to M. Gorsky [67] for the group \mathcal{B}_n^e (see also Remark 2.34).

Theorem 2.32. *The polynomial $\text{Cat}(W, x)$ is γ -positive for every finite Coxeter group W . Moreover, writing*

$$(65) \quad \text{Cat}(W, x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \gamma_i(W) x^i (1+x)^{n-2i},$$

where n is the rank of W , we have the explicit formulas

$$(66) \quad \gamma_i(W) = \begin{cases} \frac{1}{i+1} \binom{n}{i, i, n-2i}, & \text{if } W = \mathfrak{S}_{n+1}, \\ \binom{n}{i, i, n-2i}, & \text{if } W = \mathcal{B}_n, \\ \frac{n-i-1}{n-1} \binom{n}{i, i, n-2i}, & \text{if } W = \mathcal{B}_n^e, \end{cases}$$

for $0 \leq i \leq \lfloor n/2 \rfloor$.

For an extension of the γ -positivity of $\text{Cat}(W, x)$ to well-generated complex reflection groups, see [88, Theorem 1.3]. Elegant q -analogs of the γ -positivity of $\text{Cat}(\mathfrak{S}_n, x)$ are given in [22, Theorem 1.2] and [82, Theorem 4].

As is usual in Catalan combinatorics, the polynomial $\text{Cat}(W, x)$ has a positive analog, defined as

$$(67) \quad \text{Cat}^+(W, x) := \sum_{J \subseteq S} (-x)^{|S \setminus J|} \text{Cat}(W_J, x),$$

where W_J is the standard parabolic subgroup of W associated to $J \subseteq S$; we have the explicit formulas:

$$(68) \quad \text{Cat}^+(W, x) = \begin{cases} \sum_{i=0}^n \frac{1}{i+1} \binom{n-1}{i} \binom{n}{i} x^i, & \text{if } W = \mathfrak{S}_{n+1}, \\ \sum_{i=0}^n \binom{n-1}{i} \binom{n}{i} x^i, & \text{if } W = \mathcal{B}_n, \\ \sum_{i=0}^n \left(\binom{n-2}{i} \binom{n}{i} + \binom{n-2}{i-2} \binom{n-1}{i} \right) x^i, & \text{if } W = \mathcal{B}_n^e. \end{cases}$$

This polynomial is not symmetric, except for the case of the symmetric groups. However, the polynomial

$$(69) \quad \text{Cat}^{++}(W, x) := \sum_{J \subseteq S} (-1)^{|S \setminus J|} \text{Cat}^+(W_J, x)$$

turns out to be symmetric and to have nonnegative coefficients for all W . This polynomial was first considered in an enumerative-geometric context in [10], explained in Section 3.3, and more recently in a different context in [14], where our notation comes from. For the symmetric and hyperoctahedral groups, the work [10] provides combinatorial interpretations for the coefficients of $\text{Cat}^{++}(W, x)$ and the corresponding γ -coefficients $\xi_i(W)$, in terms of noncrossing partitions with restrictions.

Theorem 2.33 (ATHANASIADIS–SAVVIDOU [10]). *The polynomial $\text{Cat}^{++}(W, x)$ is γ -positive for every finite Coxeter group W . Moreover, writing*

$$(70) \quad \text{Cat}^{++}(W, x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \xi_i(W) x^i (1+x)^{n-2i},$$

where n is the rank of W , we have $\xi_0(W) = 0$ and the explicit formulas

$$(71) \quad \xi_i(W) = \begin{cases} \frac{1}{n-i+1} \binom{n}{i} \binom{n-i-1}{i-1}, & \text{if } W = \mathfrak{S}_{n+1}, \\ \binom{n}{i} \binom{n-i-1}{i-1}, & \text{if } W = \mathcal{B}_n, \\ \frac{n-2}{i} \binom{2i-2}{i-1} \binom{n-2}{2i-2}, & \text{if } W = \mathcal{B}_n^e, \end{cases}$$

for $1 \leq i \leq \lfloor n/2 \rfloor$.

Remark 2.34. Equations (67) and (69) can be inverted to give

$$(72) \quad \text{Cat}(W, x) = \sum_{J \subseteq S} x^{|S \setminus J|} \text{Cat}^+(W_J, x),$$

$$(73) \quad \text{Cat}^+(W, x) = \sum_{J \subseteq S} \text{Cat}^{++}(W_J, x),$$

respectively. Hence,

$$\begin{aligned} \text{Cat}(W, x) &= \sum_{J \subseteq S} x^{|S \setminus J|} \text{Cat}^+(W_J, x) = \sum_{J \subseteq S} x^{|S \setminus J|} \sum_{I \subseteq J} \text{Cat}^{++}(W_I, x) \\ &= \sum_{I \subseteq S} (x+1)^{n-|I|} \text{Cat}^{++}(W_I, x), \end{aligned}$$

where n is the rank of W . Setting

$$\gamma(W, x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \gamma_i(W) x^i, \quad \xi(W, x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \xi_i(W) x^i,$$

we see that the previous formula for $\text{Cat}(W, x)$ translates into the equation

$$(74) \quad \gamma(W, x) = \sum_{J \subseteq S} \xi(W_J, x).$$

Alternatively, given the geometric interpretations of $\text{Cat}(W, x)$ and $\text{Cat}^{++}(W, x)$ discussed in Section 3.3, Equation (74) is a special case of [5, Corollary 5.5] [7, Equation (6)]; see also Theorem 3.15 in the sequel.

In particular, the γ -positivity statement of Theorem 2.33 is stronger than that of Theorem 2.32. \square

The numbers $\gamma_1(W)$ and $\xi_1(W)$ are equal to the number of nonsimple reflections in W and the number of reflections in W which do not belong to any proper standard parabolic subgroup, respectively; see the first remark in [10, Section 5].

Problem 2.35. *Find a proof of the γ -positivity of $\text{Cat}(W, x)$ and $\text{Cat}^{++}(W, x)$, as well as algebraic or combinatorial interpretations for the numbers $\gamma_i(W)$ and $\xi_i(W)$, which do not depend on the classification of finite Coxeter groups.*

2.3. Polynomials arising from enriched poset partitions. Let (\mathcal{P}, ω) be a labeled poset with n elements, as in Section 2.1.2, and recall that $\Omega_m := \{1, -1, 2, -2, \dots, m, -m\}$. For $m \in \mathbb{N}$, denote by $\Omega'_{\mathcal{P}, \omega}(m)$ the number of maps $f : \mathcal{P} \rightarrow \Omega_m$ which are such that for all $x, y \in \mathcal{P}$ with $x <_{\mathcal{P}} y$:

- $|f(x)| \leq |f(y)|$,
- $|f(x)| = |f(y)|$ implies $f(x) \leq f(y)$,
- $f(x) = f(y) > 0$ implies $\omega(x) < \omega(y)$,
- $f(x) = f(y) < 0$ implies $\omega(x) > \omega(y)$,

where $\Omega'_{\mathcal{P},\omega}(0) := 0$. These maps are called *enriched* (\mathcal{P}, ω) -partitions; their theory was developed by Stembridge [133], in analogy with Stanley's theory of (\mathcal{P}, ω) -partitions. The function $\Omega'_{\mathcal{P},\omega}(m)$, studied in [133, Section 4], turns out to be a polynomial in m of degree at most n and hence

$$(75) \quad \sum_{m \geq 0} \Omega'_{\mathcal{P},\omega}(m) x^m = \frac{\mathcal{A}'_{\mathcal{P},\omega}(x)}{(1-x)^{n+1}}$$

for some polynomial $\mathcal{A}'_{\mathcal{P},\omega}(x)$ of degree at most n . As pointed out by Stembridge to the author, the more explicit formula

$$(76) \quad \mathcal{A}'_{\mathcal{P},\omega}(x) = x \sum_{\varepsilon: \mathcal{P} \rightarrow \{-1,1\}} \mathcal{A}_{\mathcal{P},\varepsilon\omega}(x)$$

follows from [133, Theorem 3.6], where $\mathcal{A}_{\mathcal{P},\varepsilon\omega}(x)$ is the ordinary $(\mathcal{P}, \varepsilon\omega)$ -Eulerian polynomial defined as in Section 2.1.2, with labels taken from the totally ordered set \mathbb{Z} .

The following result is a restatement of [133, Theorem 4.1]; it appeared several years before the work on γ -positivity of Brändén [24, 26] and Gal [61]. A *peak* of a permutation $w \in \mathfrak{S}_n$ is any index $2 \leq i \leq n-1$ such that $w(i-1) < w(i) > w(i+1)$.

Theorem 2.36 (STEMBRIDGE [133]). *For every n -element labeled poset (\mathcal{P}, ω)*

$$(77) \quad \mathcal{A}'_{\mathcal{P},\omega}(x) = \sum_{i=0}^{\lfloor (n-1)/2 \rfloor} p_{n,i} 2^{2i+1} x^{i+1} (1+x)^{n-1-2i},$$

where $p_{n,i}$ is the number of permutations $w \in \mathcal{L}(\mathcal{P}, \omega)$ (see Definition 2.6) with i peaks.

In particular, $\mathcal{A}'_{\mathcal{P},\omega}(x)$ is γ -positive.

This theorem reduces to the γ -positivity of the Eulerian polynomial $A_n(x)$ when \mathcal{P} is the n -element antichain, since in this case $\mathcal{A}'_{\mathcal{P},\omega}(x) = x 2^n A_n(x)$.

2.4. Polynomials arising from poset homology. A γ -positivity theorem, due to S. Linnusson, J. Shareshian and M. Wachs [84], which captures as special cases some of the basic examples we have seen so far, comes from the study of the homology of the Rees product of posets, a construction due to A. Björner and V. Welker [21]. Given finite graded posets \mathcal{P} and \mathcal{Q} with rank functions $\rho_{\mathcal{P}}$ and $\rho_{\mathcal{Q}}$, respectively, their *Rees product* is defined in [21] as $\mathcal{P} * \mathcal{Q} = \{(p, q) \in \mathcal{P} \times \mathcal{Q} : \rho_{\mathcal{P}}(p) \geq \rho_{\mathcal{Q}}(q)\}$, partially ordered by setting $(p_1, q_1) \preceq (p_2, q_2)$ if all of the following conditions are satisfied:

- $p_1 \preceq p_2$ holds in \mathcal{P} ,
- $q_1 \preceq q_2$ holds in \mathcal{Q} and
- $\rho_{\mathcal{P}}(p_2) - \rho_{\mathcal{P}}(p_1) \geq \rho_{\mathcal{Q}}(q_2) - \rho_{\mathcal{Q}}(q_1)$.

The poset $\mathcal{P} * \mathcal{Q}$ is graded with rank function given by $\rho(p, q) = \rho_{\mathcal{P}}(p)$ for $(p, q) \in \mathcal{P} * \mathcal{Q}$.

To state the result of [84], we need to recall a few more definitions. Let \mathcal{P} be a finite graded poset, as before, and assume further that \mathcal{P} is bounded, with minimum element

$\hat{0}$ and maximum element $\hat{1}$, and that it has rank $n + 1$. For $S \subseteq [n]$, we denote by $a_{\mathcal{P}}(S)$ the number of maximal chains of the rank-selected subposet

$$(78) \quad \mathcal{P}_S := \{p \in \mathcal{P} : \rho_{\mathcal{P}}(p) \in S\} \cup \{\hat{0}, \hat{1}\}$$

of \mathcal{P} , and set

$$(79) \quad b_{\mathcal{P}}(S) := \sum_{T \subseteq S} (-1)^{|S-T|} a_{\mathcal{P}}(T).$$

The numbers $a_{\mathcal{P}}(S)$ and $b_{\mathcal{P}}(S)$ are important enumerative invariants of \mathcal{P} ; see [127, Section 3.13]. The numbers $b_{\mathcal{P}}(S)$ are nonnegative, if \mathcal{P} is Cohen–Macaulay over some field, and afford a simple combinatorial interpretation, if \mathcal{P} admits an R -labeling. The number $(-1)^{n-1} b_{\mathcal{P}}([n])$ is the *Möbius number* of the poset $\bar{\mathcal{P}}$, obtained from \mathcal{P} by removing $\hat{0}$ and $\hat{1}$; it is denoted by $\mu(\bar{\mathcal{P}})$. The posets obtained from \mathcal{P} by removing $\hat{0}$ or $\hat{1}$ are denoted by \mathcal{P}_- or \mathcal{P}^- , respectively. The poset whose Hasse diagram is a complete x -ary tree of height n , rooted at the minimum element, is denoted by $\mathcal{T}_{x,n}$. For the notion of EL-shellability, the reader is referred to [84, Section 2] and references therein. The following theorem is a restatement of [84, Corollary 3.8].

Theorem 2.37 (LINUSSON–SHARESHIAN–WACHS [84]). *Let \mathcal{P} be a finite bounded graded poset of rank $n + 1$. If \mathcal{P} is EL-shellable, then*

$$(80) \quad \mu((\mathcal{P}^- * \mathcal{T}_{x,n})_-) = \sum_{S \in \text{Stab}([n-1])} b_{\mathcal{P}}([n] \setminus S) x^{|S|} (1+x)^{n-2|S|} \\ + \sum_{S \in \text{Stab}([n-2])} b_{\mathcal{P}}([n-1] \setminus S) x^{|S|+1} (1+x)^{n-1-2|S|}$$

and

$$(81) \quad \mu(\bar{\mathcal{P}} * \mathcal{T}_{x,n-1}) = \sum_{S \in \text{Stab}([2,n-2])} b_{\mathcal{P}}([n-1] \setminus S) x^{|S|+1} (1+x)^{n-2-2|S|} \\ + \sum_{S \in \text{Stab}([2,n-1])} b_{\mathcal{P}}([n] \setminus S) x^{|S|} (1+x)^{n-1-2|S|}$$

for every positive integer x .

This theorem turns out to be valid without the assumption of EL-shellability; see [9]. When \mathcal{P}^- has a maximum element, the first and the second summand of the right-hand sides of Equations (80) and (81), respectively, vanishes and hence the left-hand sides are γ -positive polynomials in x , provided \mathcal{P} is Cohen–Macaulay over some field.

Example 2.38. Suppose that \mathcal{P}^- is the Boolean lattice of subsets of the set $[n]$, ordered by inclusion. Then, the left-hand sides of Equations (80) and (81) are equal to $xA_n(x)$ and $d_n(x)$, respectively (this is the content of [110, Equation (3.1)] in the former case, and is implicit in [110] in the latter). The number $b_{\mathcal{P}}(S)$ is known to count permutations $w \in \mathfrak{S}_n$ such that $\text{Des}(w) = S$ [127, Corollary 3.13.2]. Thus, Theorem 2.37 reduces to

the γ -positivity of $A_n(x)$ and $d_n(x)$ (specifically, to the third interpretations given for $\gamma_{n,i}$ and $\xi_{n,i}$ in Theorems 2.1 and 2.13, respectively) in this case. \square

The $p = 1$ specializations of Theorems 2.3 and 2.15 can be viewed as the special case of Theorem 2.37 in which \mathcal{P}^- is the lattice of subspaces of an n -dimensional vector space over a field with q elements. This follows from Equations (1.3) and (1.4) in [84] and from the known interpretation [127, Theorem 3.13.3] of $b_{\mathcal{P}}(S)$ for this poset. Theorem 2.26 is derived in [6, Section 4] from the special case of Theorem 2.37 in which \mathcal{P}^- is the set of r -colored subsets of $[n]$, ordered by inclusion.

Similarly, Theorem 2.25 can be viewed as the special case of Theorem 2.37 in which \mathcal{P}^- is the product of chains whose lengths are the parts of λ ; see [84, Section 5]. The proof of Theorem 2.25 given there, however, involves a different approach and thus, it would be interesting to prove directly that the left-hand sides of Equations (80) and (81) are equal to those of Equations (50) and (51), respectively, in this case.

Remark 2.39. Theorem 2.25 can be applied, more generally, when \mathcal{P} is the distributive lattice of order ideals of an n -element poset \mathcal{Q} , since then $b_{\mathcal{P}}(S)$ counts linear extensions of \mathcal{Q} with descent set equal to S [127, Theorem 3.13.1]. Are there combinatorial interpretations for the coefficients of the left-hand sides of Equations (80) and (81) which reduce to those of Theorem 2.25 when \mathcal{Q} is a disjoint union of chains of cardinalities equal to the parts of λ ? \square

2.5. Polynomials arising from symmetric functions. Let $\mathbf{x} = (x_1, x_2, \dots)$ be a sequence of commuting independent indeterminates (to avoid confusion with this notation, in this section we consider polynomials in the variable t , rather than x). For integer partitions λ , consider the polynomials $P_\lambda(t)$ and $R_\lambda(t)$ which are defined by the equations

$$(82) \quad \frac{(1-t)H(\mathbf{x}; z)}{H(\mathbf{x}; tz) - tH(\mathbf{x}; z)} = \frac{\sum_{k \geq 0} h_k(\mathbf{x})z^k}{1 - \sum_{k \geq 2} (t + t^2 + \dots + t^{k-1}) h_k(\mathbf{x})z^k} = \sum_{\lambda} P_\lambda(t) s_\lambda(\mathbf{x}) z^{|\lambda|}$$

and

$$(83) \quad \frac{1-t}{H(\mathbf{x}; tz) - tH(\mathbf{x}; z)} = \frac{1}{1 - \sum_{k \geq 2} (t + t^2 + \dots + t^{k-1}) h_k(\mathbf{x})z^k} = \sum_{\lambda} R_\lambda(t) s_\lambda(\mathbf{x}) z^{|\lambda|},$$

where $H(\mathbf{x}; z) = \sum_{n \geq 0} h_n(\mathbf{x})z^n$ is the generating function for the complete homogeneous symmetric functions in \mathbf{x} , and λ ranges over all partitions (including the one without any parts). The left-hand sides of Equations (82) and (83) were first considered by Stanley [122, Propositions 12 and 13] and, since then, they have appeared in various algebraic-geometric and combinatorial contexts; see the relevant discussions in [111, Section 7] [84, Section 4] [113], as well as Section 5 in the sequel. Stanley [122] observed that for $\lambda \vdash n$, the polynomials $P_\lambda(t)$ and $R_\lambda(t)$ have nonnegative and symmetric coefficients, with

centers of symmetry $(n - 1)/2$ and $n/2$, respectively, which satisfy

$$(84) \quad \sum_{\lambda \vdash n} f^\lambda P_\lambda(t) = A_n(t)$$

and

$$(85) \quad \sum_{\lambda \vdash n} f^\lambda R_\lambda(t) = d_n(t),$$

where f^λ is the number of standard Young tableaux of shape λ . This shows that the coefficients of z^n in the functions appearing in Equations (82) and (83) correspond, via Frobenius characteristic, to graded \mathfrak{S}_n -representations whose graded dimensions are equal to $A_n(t)$ and $d_n(t)$, respectively. Thus, the left-hand sides of Equations (82) and (83) can be considered as representation-theoretic analogs of $A_n(t)$ and $d_n(t)$, respectively. Combinatorial interpretations for the coefficients of $P_\lambda(t)$ and $R_\lambda(t)$ were provided by Stembridge [131, Section 4]; a different one for $P_\lambda(t)$ can be derived as a special case of [112, Theorem 6.3].

The unimodality of $P_\lambda(t)$ and $R_\lambda(t)$, which refines that of $A_n(t)$ and $d_n(t)$, was proved by Stanley [122, Proposition 12] and Brenti [30, Corollary 1], respectively. Given that $A_n(t)$ and $d_n(t)$ are γ -positive, it is natural to ask whether $P_\lambda(t)$ and $R_\lambda(t)$ have the same property. An affirmative answer follows from explicit formulas which refine Theorems 2.1 and 2.13. These formulas are essentially equivalent to the following unpublished result of Gessel, stated without proof in [84, Section 4] [111, Section 7] [114, Section 3]. We write $E(\mathbf{x}; z) = \sum_{n \geq 0} e_n(\mathbf{x})z^n$ for the generating function for the elementary symmetric functions in \mathbf{x} . For a map $w : [n] \rightarrow \mathbb{Z}_{>0}$ we set $\mathbf{x}_w := x_{w(1)}x_{w(2)} \cdots x_{w(n)}$ and, as usual, define by $\text{Asc}(w) := [n - 1] \setminus \text{Des}(w)$ and $\text{Des}(w) := \{i \in [n - 1] : w(i) > w(i + 1)\}$ the set of *ascents* and *descents* of w and denote by $\text{asc}(w)$ and $\text{des}(w)$ the cardinalities of these sets, respectively.

Theorem 2.40 (GESSEL, UNPUBLISHED). *We have*

$$(86) \quad \frac{(1 - t)H(\mathbf{x}; z)}{H(\mathbf{x}; tz) - tH(\mathbf{x}; z)} = 1 + \sum_{n \geq 1} z^n \sum_{\substack{w : [n] \rightarrow \mathbb{Z}_{>0} \\ \text{Des}(w) \in \text{Stab}([n - 2])}} t^{\text{des}(w)} (1 + t)^{n - 1 - 2\text{des}(w)} \mathbf{x}_w,$$

$$(87) \quad \frac{1 - t}{H(\mathbf{x}; tz) - tH(\mathbf{x}; z)} = 1 + \sum_{n \geq 2} z^n \sum_{\substack{w : [n] \rightarrow \mathbb{Z}_{>0} \\ \text{Des}(w) \in \text{Stab}([2, n - 2])}} t^{\text{des}(w) + 1} (1 + t)^{n - 2 - 2\text{des}(w)} \mathbf{x}_w,$$

$$(88) \quad \frac{(1 - t)E(\mathbf{x}; z)}{E(\mathbf{x}; tz) - tE(\mathbf{x}; z)} = 1 + \sum_{n \geq 1} z^n \sum_{\substack{w : [n] \rightarrow \mathbb{Z}_{>0} \\ \text{Asc}(w) \in \text{Stab}([n - 2])}} t^{\text{asc}(w)} (1 + t)^{n - 1 - 2\text{asc}(w)} \mathbf{x}_w,$$

and

$$(89) \quad \frac{1-t}{E(\mathbf{x}; tz) - tE(\mathbf{x}; z)} = 1 + \sum_{n \geq 2} z^n \sum_{\substack{w : [n] \rightarrow \mathbb{Z}_{>0} \\ \text{Asc}(w) \in \text{Stab}([2, n-2])}} t^{\text{asc}(w)+1} (1+t)^{n-2-2\text{asc}(w)} \mathbf{x}_w.$$

Equations (86) and (87) can be shown to be equivalent to (88) and (89), respectively, via an application of the standard involution ω on symmetric functions. Perhaps not surprisingly, Equations (88) and (89) can be deduced [9, Corollary 4.1] as special cases of an equivariant version of Theorem 2.37; see [9, Theorem 1.2] and Theorem 5.3 in the sequel. A direct combinatorial proof of (88) will be sketched in Section 4.1.

Corollary 2.41. *For $\lambda \vdash n$,*

$$(90) \quad P_\lambda(t) = \sum_{i=0}^{\lfloor (n-1)/2 \rfloor} \gamma_i^\lambda t^i (1+t)^{n-1-2i}$$

and

$$(91) \quad R_\lambda(t) = \sum_{i=0}^{\lfloor n/2 \rfloor} \xi_i^\lambda t^i (1+t)^{n-2i},$$

where

- γ_i^λ is the number of tableaux $Q \in \text{SYT}(\lambda)$ for which $\text{Des}(Q) \in \text{Stab}([n-2])$ has i elements, and
- ξ_i^λ is the number of tableaux $Q \in \text{SYT}(\lambda)$ for which $\text{Des}(Q) \in \text{Stab}([2, n-2])$ has $i-1$ elements.

In particular, $P_\lambda(t)$ and $R_\lambda(t)$ are γ -positive for every partition λ .

Proof. For $B \subseteq [n-1]$, we consider the skew hook shape whose row lengths, read from bottom to top, form the composition of n corresponding to B and denote by $s_B(\mathbf{x})$ the skew Schur function associated to this shape. As in the proof of [114, Corollary 3.2], we note that, because of the obvious correspondence between monomials \mathbf{x}_w and semistandard Young tableaux of skew hook shape, Equation (86) may be rewritten as

$$(92) \quad \frac{(1-t)H(\mathbf{x}; z)}{H(\mathbf{x}; tz) - tH(\mathbf{x}; z)} = 1 + \sum_{n \geq 1} z^n \sum_{i=0}^{\lfloor (n-1)/2 \rfloor} \gamma_{n,i}(\mathbf{x}) t^i (1+t)^{n-1-2i},$$

where

$$(93) \quad \gamma_{n,i}(\mathbf{x}) = \sum_{\substack{B \in \text{Stab}([n-2]) \\ |B|=i}} s_B(\mathbf{x}).$$

Expanding $s_B(\mathbf{x})$ by the well known rule

$$(94) \quad s_B(\mathbf{x}) = \sum_{\lambda \vdash n} |\{Q \in \text{SYT}(\lambda) : \text{Des}(Q) = B\}| \cdot s_\lambda(\mathbf{x})$$

in the basis of Schur functions and comparing the coefficients of $s_\lambda(\mathbf{x})$ on the two sides of (92), we obtain the desired formula for $P_\lambda(t)$. A similar argument works for $R_\lambda(t)$. \square

The author is not aware of any results in the literature concerning the roots of $P_\lambda(t)$ and $R_\lambda(t)$.

2.6. Polynomials arising from trees. This section discusses two more instances of γ -positivity in combinatorics. For further examples, see [26, 81] and Sections 3.3 and 4.2.

A tree T on the vertex set $[n]$ is called *rooted* if it has a distinguished vertex, called *root*. An edge $\{a, b\}$, with $a < b$, of such a tree T is called a *descent* if the unique path in T which connects a with the root passes through b . Let $t_n(x)$ be the polynomial in which the coefficient of x^k equals the number of rooted trees on the vertex set $[n]$ with exactly k descents. The following elegant x -analog

$$(95) \quad t_n(x) = \prod_{i=1}^{n-1} ((n-i) + ix)$$

of Cayley's formula, which is a specialization of results of Ö. Eğecioğlu and J. Remmel [50], implies the γ -positivity of $t_n(x)$. Several explicit combinatorial interpretations for the corresponding γ -coefficients were found by R. González D'León [66].

A tree T on the vertex set $[n]$ is called *alternating*, or *intransitive* [100], if every vertex of T is either less than all its neighbors, in which case it is called a *left vertex*, or greater than all its neighbors, in which case it is called a *right vertex*. Let $g_n(x)$ be the polynomial in which the coefficient of x^k equals the number of alternating trees on the vertex set $[n]$ with exactly k left vertices. These polynomials were considered in [63, 100]; for the first few values of n , we have

$$g_n(x) = \begin{cases} 1, & \text{if } n = 1, \\ x, & \text{if } n = 2, \\ x + x^2, & \text{if } n = 3, \\ x + 5x^2 + x^3, & \text{if } n = 4, \\ x + 17x^2 + 17x^3 + x^4, & \text{if } n = 5, \\ x + 49x^2 + 146x^3 + 49x^4 + x^5, & \text{if } n = 6, \\ x + 129x^2 + 922x^3 + 922x^4 + 129x^5 + x^6, & \text{if } n = 7. \end{cases}$$

Theorem 2.42 (GESSEL–GRIFFIN–TEWARI [64, Theorem 5.9]). *The polynomial $g_n(x)$ is γ -positive for every positive integer n .*

The proof, which uses symmetric functions, yields a combinatorial formula for the γ -coefficients involving certain plane binary trees. An explicit combinatorial interpretation has also been found by V. Tewari [138].




FIGURE 2. A two-dimensional simplicial complex

3. GAMMA-POSITIVITY IN GEOMETRY

This section discusses the main geometric contexts in which γ -positivity occurs. Many of the examples treated in Section 2 reappear here as interesting special cases of more general phenomena. Familiarity with basic notions on simplicial complexes and their geometric realizations is assumed; see [19] [36, Chapter 2] [77] [140, Lecture 1] for detailed expositions. All simplicial complexes considered here are assumed to be finite. To keep our discussion as elementary as possible, we work with triangulations of spheres and balls, rather than with the more general classes of homology spheres and homology balls.

We first review some basic definitions and background related to the face enumeration of simplicial complexes. For $i \geq 0$ we denote by $f_{i-1}(\Delta)$ the number of $(i-1)$ -dimensional faces of a simplicial complex Δ , where (unless Δ is empty) $f_{-1}(\Delta) := 1$.

Definition 3.1. *The h -polynomial of a simplicial complex Δ of dimension $n-1$ is defined as*

$$(96) \quad h(\Delta, x) := \sum_{i=0}^n f_{i-1}(\Delta)x^i(1-x)^{n-i} := \sum_{i=0}^n h_i(\Delta)x^i.$$

The sequence $h(\Delta) = (h_0(\Delta), h_1(\Delta), \dots, h_n(\Delta))$ is the h -vector of Δ .

The polynomial $h(\Delta, x)$ can be thought of as an x -analog of the number $f_{n-1}(\Delta)$ of facets (meaning, faces of maximum dimension $n-1$) of Δ , since $h(\Delta, 1) = f_{n-1}(\Delta)$. We refer the reader to [126] for the significance of the h -polynomial, as well as for important algebraic and combinatorial interpretations which are valid for special classes of simplicial complexes.

Example 3.2. The simplicial complex Δ shown in Figure 2 triangulates a two-dimensional simplex with eight vertices, fifteen edges and eight two-dimensional faces. Thus, $f_{-1}(\Delta) = 1$, $f_0(\Delta) = 8$, $f_1(\Delta) = 15$ and $f_2(\Delta) = 8$ and we may compute that $h(\Delta, x) = (1-x)^3 + 8x(1-x)^2 + 15x^2(1-x) + 8x^3 = 1 + 5x + 2x^2$. \square

The relevance of the h -polynomial to the symmetry and unimodality of real polynomials stems from the following theorem, which combines important results by V. Klee, G. Reisner and R. Stanley [76, 105, 119, 120] in geometric combinatorics; see [126] for more

information. The first two parts hold more generally for the classes of Cohen–Macaulay and Eulerian, respectively, simplicial complexes.

Theorem 3.3 (KLEE, REISNER, STANLEY). *Let Δ be $(n - 1)$ -dimensional.*

- (a) *The polynomial $h(\Delta, x)$ has nonnegative coefficients, i.e., we have $h_i(\Delta) \geq 0$ for $0 \leq i \leq n$, if Δ triangulates a ball or a sphere.*
- (b) *The polynomial $h(\Delta, x)$ is symmetric, i.e., we have $h_i(\Delta) = h_{n-i}(\Delta)$ for $0 \leq i \leq n$, if Δ triangulates a sphere.*
- (c) *The polynomial $h(\Delta, x)$ is unimodal, i.e., we have*

$$(97) \quad 1 = h_0(\Delta) \leq h_1(\Delta) \leq \cdots \leq h_{\lfloor n/2 \rfloor}(\Delta),$$

if Δ is the boundary complex of a simplicial polytope.

Part (c), proved by R. Stanley [120], is known as the *Generalized Lower Bound Theorem* for simplicial polytopes, since the inequalities $h_i(\Delta) \geq h_{i-1}(\Delta)$ impose a lower bound for each face number $f_{i-1}(\Delta)$ in terms of the numbers $f_{j-1}(\Delta)$ for $1 \leq j < i$.

3.1. Flag triangulations of spheres. Let Δ be an (abstract) simplicial complex on the vertex set V . We say that Δ is *flag* if we have $F \in \Delta$ for every $F \subseteq V$ for which all two-element subsets of F belong to Δ , and refer to [4] [126, Section III.4] for a glimpse of the combinatorial properties of this fascinating class of simplicial complexes. The complex shown in Figure 2, for example, is flag.

The following major open problem in geometric combinatorics was posed by Š. Gal [61] who disproved a more optimistic conjecture, claiming that $h(\Delta, x)$ is real-rooted for every flag triangulation of a sphere.

Conjecture 3.4 (GAL [61, Conjecture 2.1.7]). *The polynomial $h(\Delta, x)$ is γ -positive for every flag simplicial complex Δ which triangulates a sphere.*

This conjecture extends an earlier conjecture of R. Charney and M. Davis [37] on the sign of $h(\Delta, -1)$; it is also open for the more restrictive class of flag boundary complexes of simplicial polytopes.

Example 3.5. Consider a triangulation Δ of the one-dimensional sphere with m vertices (cycle of length m). We have $f_{-1}(\Delta) = 1$ and $f_0(\Delta) = f_1(\Delta) = m$ and hence $h(\Delta, x) = (1 - x)^2 + mx(1 - x) + mx^2 = 1 + (m - 2)x + x^2$ is γ -positive if and only if $m \geq 4$. Note that this is exactly the necessary and sufficient condition for Δ to be flag. \square

Conjecture 3.4 can be viewed as a Generalized Lower Bound Conjecture for flag triangulations of spheres Δ . To be more precise, we may write

$$(98) \quad h(\Delta, x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \gamma_i(\Delta) x^i (1 + x)^{n-2i},$$

where $n - 1 = \dim(\Delta)$ and $\gamma_0(\Delta) = 1$. The vector $\gamma(\Delta) := (\gamma_0(\Delta), \gamma_1(\Delta), \dots, \gamma_{\lfloor n/2 \rfloor}(\Delta))$ is known as the γ -vector of Δ and Gal’s conjecture states that $\gamma_i(\Delta) \geq 0$ for all i . Just as in the case of Theorem 3.3 (c), these inequalities impose a lower bound on each $h_i(\Delta)$ in

terms of the numbers $h_j(\Delta)$ for $0 \leq j < i$, and hence a lower bound on each face number $f_{i-1}(\Delta)$ in terms of the face numbers $f_{j-1}(\Delta)$ for $1 \leq j < i$.

For instance, since $\gamma_1(\Delta) = h_1(\Delta) - n = f_0(\Delta) - 2n$, the inequality $\gamma_1(\Delta) \geq 0$ states that every flag triangulation of the $(n-1)$ -dimensional sphere has at least $2n$ vertices, a fact which is easy to prove by induction on n . The inequality $\gamma_2(\Delta) \geq 0$ already provides a challenge. Since

$$\gamma_2(\Delta) = h_2(\Delta) - (n-2)\gamma_1(\Delta) - \binom{n}{2} = f_1(\Delta) - (2n-3)f_0(\Delta) + 2n(n-2),$$

its validity turns out to be equivalent to the following conjectural flag analog of Barnette's Lower Bound Theorem [15]. We recall that the suspension (simplicial join with a zero-dimensional sphere) of a flag triangulation of a sphere is a flag triangulation of a sphere of one dimension higher.

Conjecture 3.6. *Among all flag triangulations of the $(n-1)$ -dimensional sphere with a given number m of vertices, the $(n-2)$ -fold suspension over the one-dimensional sphere with $m-2n+4$ vertices has the smallest possible number of edges.*

For a generalization of this statement, see [90, Conjecture 1.4]. A conjecture concerning the possible vectors which can arise as γ -vectors of flag triangulations of spheres is proposed in [91].

An edge subdivision of a simplicial complex Δ is a stellar subdivision of Δ on any of its edges; we refer to [5, Section 6] [37, Section 5.3] [61, Section 2.4] for the precise definition and Figure 3 for an example. Edge subdivisions preserve flagness and homeomorphism type. We will denote by Σ_n the simplicial join of n copies of the zero-dimensional sphere (this simplicial complex is combinatorially isomorphic to the boundary complex of the n -dimensional cross-polytope; it satisfies $h(\Sigma_n, x) = (1+x)^n$ or, equivalently, $\gamma(\Sigma_n) = (1, 0, \dots, 0)$). Apart from the results on barycentric subdivisions and nested complexes, discussed separately in the sequel, Gal's conjecture is known to hold for flag triangulations of spheres:

- of dimension less than five [44, Theorem 11.2.1] [61, Corollary 2.2.3],
- with at most $2n+3$ vertices, i.e., with $\gamma_1(\Delta) \leq 3$ [91, Theorem 1.3],
- which can be obtained from Σ_n by successive edge subdivisions [61, Section 2.4],
- for other special classes of flag triangulations of spheres, discussed in Section 3.3.

Remark 3.7. The proof of the third statement above is only implicit in [61, Section 2.4]. To make it more explicit, let us denote by Δ_e the edge subdivision of Δ on its edge $e \in \Delta$. Gal observes [61, Proposition 2.4.3] that

$$(99) \quad h(\Delta_e, x) = h(\Delta, x) + x h(\text{link}_\Delta(e), x),$$

where $\text{link}_\Delta(F) = \{G \setminus F : G \in \Delta, F \subseteq G\}$ is the *link* of $F \in \Delta$ in Δ . In particular, if Δ and $\text{link}_\Delta(e)$ satisfy Gal's conjecture, then so does Δ [61, Corollary 2.4.7]. One can verify that every link of a nonempty face in Δ_e is combinatorially isomorphic to either the link of a nonempty face in Δ , or to an edge subdivision or the suspension of the link of a nonempty face in Δ . Thus, it follows by induction and Equation (99) that, if Δ can be




FIGURE 3. An edge subdivision

obtained from Σ_n by successive edge subdivisions, then Δ and all links of its faces satisfy Gal’s conjecture. \square

3.1.1. *Barycentric subdivisions.* An important class of flag simplicial complexes is that of order complexes [126, Section III.4]. Recall that the *order complex* of a poset \mathcal{P} is defined as the simplicial complex $\Delta(\mathcal{P})$ of all chains (totally ordered subsets) of \mathcal{P} . When \mathcal{P} is the poset of (nonempty) faces of a regular CW-complex X , the complex $\Delta(\mathcal{P})$ is by definition the (first) barycentric subdivision of X . As explained, for instance, in [61, Section 2.3], the following result is a direct consequence of important theorems of K. Karu and R. Stanley on the nonnegativity of the cd-index of Gorenstein* posets [74] (for introductions to the cd-index, see [126, Section III.4] [127, Section 3.17]).

Theorem 3.8 (KARU [74], STANLEY [124]; SEE [61, Corollary 2.3.5]). *Conjecture 3.4 holds for every order complex Δ which triangulates a sphere. In particular, it holds for barycentric subdivisions of regular CW-spheres.*

As recorded in [37, Section 7.3] [126, p. 103], the connection between the cd-index of an Eulerian poset \mathcal{P} and the last entry of the γ -vector of $\Delta(\mathcal{P})$ was observed by E. Babson. The nonnegativity of the cd-index was proved for the augmented face posets of certain shellable CW-spheres by Stanley [124, Theorem 2.2] and for arbitrary Gorenstein* posets by Karu [74, Theorem 1.3] (Stanley’s result suffices to conclude the validity of Gal’s conjecture for barycentric subdivisions of boundary complexes of convex polytopes). For barycentric subdivisions of Boolean complexes which triangulate a sphere, more elementary proofs of Theorem 3.8 can be found in [33, 92].

There is an interesting analog of Theorem 3.8 for triangulations of balls, rather than spheres. As explained in [73, Section 4], the following result is a direct consequence of [51, Theorem 2.5]. Here $\partial\Delta$ denotes the boundary of a triangulated ball Δ .

Theorem 3.9 (EHRENBORG–KARU [51]; see [73, THEOREM 4.6]). *The polynomial $h(\Delta, x) - h(\partial\Delta, x)$ is γ -positive for every order complex Δ which triangulates a ball. In particular, this holds for barycentric subdivisions of regular CW-balls.*

3.1.2. *Flag nested complexes.* We begin with a few definitions, following [141]. A *building set* on the ground set $[n]$ is a collection \mathcal{B} of nonempty subsets of $[n]$ such that:




FIGURE 4. A nested complex

- \mathcal{B} contains all singletons $\{i\} \subseteq [n]$, and
- if $I, J \in \mathcal{B}$ and $I \cap J \neq \emptyset$, then $I \cup J \in \mathcal{B}$.

We denote by \mathcal{B}_{\max} the set of maximal (with respect to inclusion) elements of \mathcal{B} and say that \mathcal{B} is *connected* if $[n] \in \mathcal{B}$. A set $N \subseteq \mathcal{B} \setminus \mathcal{B}_{\max}$ is called *nested* if for all $k \geq 2$ and all $I_1, I_2, \dots, I_k \in N$ such that none of the I_i contains another, their union $I_1 \cup I_2 \cup \dots \cup I_k$ is not in \mathcal{B} . The simplicial complex $\Delta_{\mathcal{B}}$ on the vertex set $\mathcal{B} \setminus \mathcal{B}_{\max}$, consisting of all nested sets, is the *nested complex* associated to \mathcal{B} . For example, if \mathcal{B} consists of all nonempty subsets of $[n]$, then $\Delta_{\mathcal{B}}$ is the barycentric subdivision of the boundary of the simplex $2^{[n]}$.

The nested complex $\Delta_{\mathcal{B}}$ is isomorphic to the boundary complex of a simplicial polytope of dimension $n - |\mathcal{B}_{\max}|$ [52, Theorem 3.14] [101, Theorem 7.4] [141, Theorem 6.1]; the corresponding polar-dual simple polytope is called a *nestohedron*. Nestohedra form an important class of Postnikov's generalized permutohedra [101] which includes permutohedra, graph-associahedra and other well studied simple polytopes in geometric combinatorics. As shown in [53, Theorem 4] and explained in [102, Remark 6.6], the complex $\Delta_{\mathcal{B}}$ can be constructed as follows. Assume (without loss of generality) that \mathcal{B} is connected, and consider the boundary $\partial\sigma$ of the simplex σ on the vertex set $[n]$. Choose any ordering of the nonsingleton elements of $\mathcal{B} \setminus \mathcal{B}_{\max}$ which respects the reverse of the inclusion order. Starting with $\partial\sigma$, and following the chosen ordering, for each nonsingleton $I \in \mathcal{B} \setminus \mathcal{B}_{\max}$ perform a stellar subdivision on the face I . The resulting simplicial complex is $\Delta_{\mathcal{B}}$.

Example 3.10. For $n = 4$, consider the building set \mathcal{B} consisting of $\{1\}, \{2\}, \{3\}, \{4\}, \{3, 4\}, \{2, 3, 4\}$ and $\{1, 2, 3, 4\}$. The nested complex $\Delta_{\mathcal{B}}$ triangulates the boundary of the three-dimensional simplex on the vertex set $\{\{1\}, \{2\}, \{3\}, \{4\}\}$; it has eight facets, six of which are shown in Figure 4 (the two unlabeled vertices being $\{3, 4\}$ and $\{2, 3, 4\}$). The remaining facets are the nonvisible facets $\{\{1\}, \{2\}, \{3\}\}$ and $\{\{1\}, \{2\}, \{4\}\}$ of the simplex. Note that $\Delta_{\mathcal{B}}$ corresponds indeed to the simplicial complex obtained by stellarly subdividing the boundary complex of the simplex on the vertex set $\{1, 2, 3, 4\}$ first on the face $\{2, 3, 4\}$ and then on $\{3, 4\}$. \square

Gal’s conjecture was verified for flag nested complexes by V. Volodin [139], who showed that they can be obtained from the boundary complex of a cross-polytope by successive edge subdivisions. The main result of [102] provides an explicit combinatorial interpretation of the entries of the γ -vector of a large family of flag nested complexes. To state this result, we need to introduce a few definitions from [102]. Let \mathcal{B} be a building set on the ground set $[n]$. The *restriction* of \mathcal{B} to $I \subseteq [n]$ is defined as the family of all subsets of I which belong to \mathcal{B} . The restrictions of \mathcal{B} to the elements of \mathcal{B}_{\max} are the *connected components* of \mathcal{B} . The building set \mathcal{B} is called *chordal* if for every $\{i_1, i_2, \dots, i_k\} \in \mathcal{B}$ with $i_1 < i_2 < \dots < i_k$ and all indices $1 \leq r \leq k$ we have $\{i_r, i_{r+1}, \dots, i_k\} \in \mathcal{B}$. Nested complexes of chordal building sets are always flag [102, Proposition 9.7]. We denote by $\mathfrak{S}_{\mathcal{B}}$ the set of all permutations $w \in \mathfrak{S}_n$ such that $w(i)$ and $\max\{w(1), w(2), \dots, w(i)\}$ lie in the same connected component of the restriction of \mathcal{B} to $\{w(1), w(2), \dots, w(i)\}$, for every $i \in [n]$. The following result combines Corollary 9.6 with Theorem 11.6 in [102] and, as shown in [102, Section 10], provides a common generalization for the γ -positivity of the Eulerian and binomial Eulerian polynomials $A_n(x)$ and $\tilde{A}_n(x)$ and the Narayana polynomials $\text{Cat}(W, x)$ for the symmetric and hyperoctahedral groups, discussed in Section 2.

Theorem 3.11 (POSTNIKOV–REINER–WILLIAMS [102]). *For every connected, chordal building set \mathcal{B} on the ground set $[n]$*

$$(100) \quad h(\Delta_{\mathcal{B}}, x) = \sum_{i=0}^n h_{\mathcal{B},i} x^i = \sum_{i=0}^{\lfloor n/2 \rfloor} \gamma_{\mathcal{B},i} x^i (1+x)^{n-2i},$$

where

- $h_{\mathcal{B},i}$ is the number of permutations $w \in \mathfrak{S}_{\mathcal{B}}$ with i descents, and
- $\gamma_{\mathcal{B},i}$ is the number of permutations $w \in \mathfrak{S}_{\mathcal{B}}$ for which $\text{Des}(w) \in \text{Stab}([n-2])$ has i elements.

The nested complex $\Delta_{\mathcal{B}}$ of Example 3.10 has eight facets. The elements of $\mathfrak{S}_{\mathcal{B}}$, written in one-line notation, are 1234, 1243, 1342, 1432, 2341, 2431, 3421 and 4321, and $h(\Delta_{\mathcal{B}}, x) = 1 + 3x + 3x^2 + x^3 = (1+x)^3$, in agreement with Theorem 3.11.

3.2. Flag triangulations of simplices. This section focuses on a variant of the h -polynomial of a triangulation of a sphere, the local h -polynomial, defined for triangulations of simplices (and for more general topological, rather than geometric, simplicial subdivisions of the simplex; see [123, Part I] [5, 7]). The local h -polynomial was introduced by Stanley [123] and plays a key role in his enumerative theory of triangulations of simplicial complexes, developed in order to study their effect on the h -polynomial of a simplicial complex.

Let V be an n -element set and Γ be a triangulation of the simplex 2^V . The restriction of Γ to the face $F \subseteq V$ of the simplex is a triangulation of F , denoted by Γ_F .

Definition 3.12 (STANLEY [123, Definition 2.1]). *The local h -polynomial of Γ (with respect to V) is defined as*

$$(101) \quad \ell_V(\Gamma, x) := \sum_{F \subseteq V} (-1)^{n-|F|} h(\Gamma_F, x) := \ell_0 + \ell_1 x + \cdots + \ell_n x^n.$$

The sequence $\ell_V(\Gamma) = (\ell_0, \ell_1, \dots, \ell_n)$ is the local h -vector of Γ (with respect to V).

For example, the complex of Figure 2 is naturally a triangulation Γ of a two-dimensional simplex 2^V . Since $h(\Gamma, x) = 1 + 5x + 2x^2$ and the restrictions of Γ on the three edges of the simplex have h -polynomials 1 , $1 + x$ and $1 + 2x$, we have $\ell_V(\Gamma, x) = (1 + 5x + 2x^2) - 1 - (1 + x) - (1 + 2x) + 1 + 1 + 1 - 1 = 2x + 2x^2$.

The following theorem shows that the local h -polynomial has properties similar to those of the h -polynomial of a triangulation of the sphere, stated in Theorem 3.3. For the notion of a regular triangulation, we refer to [45, Chapter 5] [123, Definition 5.1].

Theorem 3.13 (STANLEY [123]). *Let Γ be a triangulation of an $(n - 1)$ -dimensional simplex 2^V and set $\ell_V(\Gamma, x) = \ell_0 + \ell_1 x + \cdots + \ell_n x^n$.*

- (a) *The polynomial $\ell_V(\Gamma, x)$ has nonnegative coefficients.*
- (b) *The polynomial $\ell_V(\Gamma, x)$ is symmetric, i.e., we have $\ell_i = \ell_{n-i}$ for $0 \leq i \leq n$.*
- (c) *The polynomial $\ell_V(\Gamma, x)$ is unimodal, i.e., we have $\ell_0 \leq \ell_1 \leq \cdots \leq \ell_{\lfloor n/2 \rfloor}$, if Γ is regular.*

The following analog of Conjecture 3.4 may come as no surprise to some readers; it is stated more generally in [5, Conjecture 5.4] for a class of topological simplicial subdivisions which models combinatorially geometric subdivisions.

Conjecture 3.14 (ATHANASIADIS [5]). *The polynomial $\ell_V(\Gamma, x)$ is γ -positive for every flag triangulation Γ of the simplex 2^V .*

Since $\ell_V(\Gamma, x)$ is symmetric, with center of symmetry $n/2$, we may write

$$(102) \quad \ell_V(\Gamma, x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \xi_i(\Gamma) x^i (1+x)^{n-2i}.$$

The sequence $\xi(\Gamma) := (\xi_0(\Gamma), \xi_1(\Gamma), \dots, \xi_{\lfloor n/2 \rfloor}(\Gamma))$ is termed in [5] as the *local γ -vector* of Γ (with respect to V) and the polynomial $\xi_V(\Gamma, x) := \sum_{i=0}^{\lfloor n/2 \rfloor} \xi_i(\Gamma) x^i$ as the corresponding *local γ -polynomial*. Thus, Conjecture 3.14 claims that $\xi_i(\Gamma) \geq 0$ for all $0 \leq i \leq \lfloor n/2 \rfloor$ and every flag triangulation Γ of the simplex 2^V . Apart from results on barycentric subdivisions, discussed in the sequel, Conjecture 3.14 has been verified for flag triangulations of simplices:

- of dimension less than four [5, Proposition 5.7],
- which can be obtained from the trivial triangulation by successive edge subdivisions [5, Proposition 6.1],
- for other special classes of flag triangulations of simplices, discussed in Section 3.3.

The connection between γ -vectors and their local counterparts is explained by the following result. We recall that Σ_n stands for the boundary complex of the n -dimensional cross-polytope and refer to [5] [7, Section 2] for the precise definitions of the more general notions of simplicial subdivision which appear there.

Theorem 3.15 (ATHANASIADIS [5] [7, Theorem 3.7]). *Every flag triangulation Δ of the $(n - 1)$ -dimensional sphere is a vertex-induced simplicial homology subdivision of Σ_n . Moreover,*

$$(103) \quad \gamma(\Delta, x) = \sum_{F \in \Sigma_n} \xi_F(\Delta_F, x).$$

The version of Conjecture 3.14 stated as [5, Conjecture 5.4] applies to the summands in the right-hand side of (103). As a result, [5, Conjecture 5.4] implies Gal’s conjecture, as well as its stronger version [61, Conjecture 2.1.7] for flag homology spheres. Moreover, as explained at the end of [8, Section 1], the validity of Conjecture 3.14 for regular flag triangulations of simplices implies Gal’s conjecture for flag boundary complexes of convex polytopes.

The following statement is a direct consequence of Theorem 3.15 and Stanley’s monotonicity theorem [123, Theorem 4.10] (in the form of [7, Corollary 2.9]) for h -vectors of subdivisions of simplicial complexes; it was proved more generally for flag doubly Cohen–Macaulay complexes, using different methods, in [4, Theorem 1.3].

Corollary 3.16. *Every flag triangulation Δ of the $(n - 1)$ -dimensional sphere satisfies*

$$(104) \quad h_i(\Delta) \geq \binom{n}{i}$$

for $0 \leq i \leq n$.

3.2.1. Barycentric subdivisions. Barycentric subdivisions of polyhedral subdivisions of the simplex form a natural class of its flag triangulations. The question of the validity of Conjecture 3.14 for them was raised in [5, Question 6.2] [7, Problem 3.8] and answered in the affirmative by Juhnke-Kubitzke, Murai and Sieg [73, Theorem 1.3]. The last statement in the following theorem is a consequence of Equation (105), Theorem 3.9 and the γ -positivity of the derangement polynomials $d_n(x)$.

Theorem 3.17 (KUBITZKE–MURAI–SIEG [73]). *For every triangulation Γ of an $(n - 1)$ -dimensional simplex 2^V ,*

$$(105) \quad \ell_V(\Gamma, x) = \sum_{F \subseteq V} (h(\Gamma_F, x) - h(\partial(\Gamma_F), x)) d_{n-|F|}(x).$$

In particular, Conjecture 3.14 holds for barycentric subdivisions of regular CW-complexes which subdivide the simplex.

3.2.2. *Nested subdivisions.* Let \mathcal{B} be a connected building set on the ground set $[n]$. We denote by $\Gamma_{\mathcal{B}}$ the cone of the nested complex $\Delta_{\mathcal{B}}$. Alternatively, $\Gamma_{\mathcal{B}}$ is the simplicial complex on the vertex set \mathcal{B} , consisting of all nested subsets of \mathcal{B} (rather than $\mathcal{B} \setminus \mathcal{B}_{\max}$), defined by the condition described in Section 3.1.2. Since $\Delta_{\mathcal{B}}$ triangulates the boundary of an $(n - 1)$ -dimensional simplex, the complex $\Gamma_{\mathcal{B}}$ is a triangulation of this simplex (for example, it is the barycentric subdivision, if \mathcal{B} consists of all nonempty subsets of $[n]$). We call $\Gamma_{\mathcal{B}}$ the *nested subdivision* associated to \mathcal{B} .

No doubt, just as is the case with $\Delta_{\mathcal{B}}$, one can show that whenever $\Gamma_{\mathcal{B}}$ is a flag complex, it can be obtained from the trivial subdivision of the $(n - 1)$ -dimensional simplex by successive edge subdivisions. Thus, the γ -positivity of the local h -polynomial of $\Gamma_{\mathcal{B}}$ follows from [5, Proposition 6.1]. In view of Theorem 3.11, it seems natural to pose the following problem.

Problem 3.18. Find a combinatorial interpretation for the local γ -polynomial of $\Gamma_{\mathcal{B}}$, for any connected chordal building set \mathcal{B} .

3.3. **Examples.** This section discusses further examples of flag triangulations of spheres or simplices, which appear naturally in algebraic-geometric contexts, and the γ -positivity of their h -polynomials or local h -polynomials. These examples provide algebraic-geometric interpretations for several of the γ -positive polynomials discussed in Section 2.

3.3.1. *Barycentric and edgewise subdivisions.* As noted in Section 3.1.1, the barycentric subdivision of a regular CW-complex X , denoted here by $\text{sd}(X)$, is defined as the simplicial complex of all chains in the poset of nonempty faces of X .

Let V be an n -element set, and consider the barycentric subdivision $\text{sd}(2^V)$ of the simplex 2^V . The facets of $\text{sd}(2^V)$ are in one-to-one correspondence with the permutations of V . Thus, the h -polynomial of $\text{sd}(2^V)$ is an x -analog of the number $n!$; it is well known, in fact, that

$$(106) \quad h(\text{sd}(2^V), x) = A_n(x).$$

Since coning a simplicial complex does not affect its h -polynomial, $A_n(x)$ is also equal to the h -polynomial of the barycentric subdivision $\text{sd}(\partial(2^V))$ of the boundary complex of 2^V . Moreover, given that $\text{sd}(2^V)$ restricts to the barycentric subdivision $\text{sd}(2^F)$ for every $F \subseteq V$, Stanley [123, Proposition 2.4] showed that

$$(107) \quad \ell_V(\text{sd}(2^V), x) = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} A_k(x) = \sum_{w \in \mathcal{D}_n} x^{\text{exc}(w)} = d_n(x).$$

Consequently, the unimodality of $A_n(x)$ and $d_n(x)$ follows from parts (c) of Theorems 3.3 and 3.13, and their γ -positivity is an instance of Conjectures 3.4 and 3.14, respectively. A part of the barycentric subdivision of the boundary complex of the three-dimensional simplex is shown in Figure 5. The restrictions on the facets of the simplex are barycentric subdivisions of two-dimensional simplices.

The r -fold edgewise subdivision, denoted here by $\text{esd}_r(\Delta)$, is another elegant (but not as well known as barycentric subdivision) triangulation of a simplicial complex Δ with




FIGURE 5. Two Coxeter complexes




FIGURE 6. An edgewise subdivision

a long history in mathematics; see [8, Section 1] and references therein. To describe it in a geometrically intuitive way, consider positive integers n, r and the geometric simplex $\sigma_{n,r} = \{(x_1, x_2, \dots, x_{n-1}) \in \mathbb{R}^{n-1} : 0 \leq x_{n-1} \leq \dots \leq x_2 \leq x_1 \leq r\}$. The r -fold edgewise subdivision $\text{esd}_r(2^V)$ of an $(n-1)$ -dimensional simplex 2^V is realized as the triangulation of $\sigma_{n,r}$ whose facets are the $(n-1)$ -dimensional simplices into which $\sigma_{n,r}$ is dissected by the affine hyperplanes in \mathbb{R}^{n-1} with equations

- $x_i = k$, where $1 \leq i \leq n-1$,
- $x_i - x_j = k$, where $1 \leq i < j \leq n-1$

for $k \in \{1, 2, \dots, r-1\}$. This triangulation has exactly r^{n-1} facets; it is shown in Figure 6 for $n = 3$ and $r = 4$. For an arbitrary simplicial complex Δ , the r -fold edgewise subdivision $\text{esd}_r(\Delta)$ restricts to $\text{esd}_r(2^F)$ for every $F \in \Delta$; for a formal definition, see, for instance, [6, Section 4] [34, Section 4].

The triangulation $\text{esd}_r(\Delta)$ is flag for every flag simplicial complex Δ ; in particular, $\text{esd}_r(2^V)$ is a flag triangulation of the simplex. Combinatorial interpretations for its local h -vector and local γ -vector can be given as follows. Let us denote by $\mathcal{W}(n, r)$ the set of sequences $w = (w_0, w_1, \dots, w_n) \in \{0, 1, \dots, r-1\}^{n+1}$ satisfying $w_0 = w_n = 0$, and by $\mathcal{S}(n, r)$ the set of those elements of $\mathcal{W}(n, r)$ (known as *Smirnov words*) having no two consecutive entries equal. As usual, we call $i \in \{0, 1, \dots, n-1\}$ an *ascent* of $w \in \mathcal{W}(n, r)$ if $w_i \leq w_{i+1}$ and $i \in [n-1]$ a *double ascent* if $w_{i-1} \leq w_i \leq w_{i+1}$ (descents and double descents are defined similarly, using strict inequalities). We also denote by E_r the linear operator on the space $\mathbb{R}[x]$ of polynomials in x with real coefficients defined by setting $E_r(x^m) = x^{m/r}$,

FIGURE 7. The triangulation K_3

if m is divisible by r , and $E_r(x^m) = 0$ otherwise. For the second interpretation of $\xi_{n,r,i}$ which appears in the following result (but not in [7, 8]), see Remark 4.1.

Theorem 3.19 (ATHANASIADIS [7, Theorem 4.6] [8]). *The local h -polynomial of the r -fold edgewise subdivision $\text{esd}_r(2^V)$ of the $(n-1)$ -dimensional simplex on the vertex set V can be expressed as*

$$\begin{aligned} \ell_V(\text{esd}_r(2^V), x) &= E_r(x + x^2 + \cdots + x^{r-1})^n = \sum_{w \in \mathcal{S}(n,r)} x^{\text{asc}(w)} \\ &= \sum_{i=0}^{\lfloor n/2 \rfloor} \xi_{n,r,i} x^i (1+x)^{n-2i}, \end{aligned}$$

where $\text{asc}(w)$ is the number of ascents of $w \in \mathcal{S}(n,r)$ and $\xi_{n,r,i}$ equals each of the following:

- the number of sequences $w = (w_0, w_1, \dots, w_n) \in \mathcal{S}(n,r)$ with exactly i ascents which have the following property: for every double ascent k of w there exists a double descent $\ell > k$ such that $w_k = w_\ell$ and $w_k \leq w_j$ for all $k < j < \ell$,
- the number of $w \in \mathcal{W}(n,r)$ for which $\text{Asc}(w) \in \text{Stab}([n-2])$ has i elements, where $\text{Asc}(w)$ is the set of ascents of w .

The interpretations of the polynomials $A_n(x)$ and $d_n(x)$, described in this section, have interesting hyperoctahedral group analogs. Let \mathcal{C}_n denote the boundary complex of the standard n -dimensional unit cube. The facets of the barycentric subdivision $\text{sd}(\mathcal{C}_n)$ are in one-to-one correspondence with the signed permutations $w \in \mathcal{B}_n$. As a result, the h -polynomial of $\text{sd}(\mathcal{C}_n)$ is an x -analog of the number $2^n n!$ and, in fact (see our discussion in Section 3.3.2),

$$(108) \quad h(\text{sd}(\mathcal{C}_n), x) = B_n(x),$$

where $B_n(x)$ is the \mathcal{B}_n -Eulerian polynomial, discussed in Section 2.1.3. Consider now the cubical barycentric subdivision of an $(n-1)$ -dimensional simplex 2^V . This is a cubical complex whose face poset is isomorphic to the set of nonempty closed intervals in the poset of nonempty subsets of V , partially ordered by inclusion. Its (simplicial) barycentric subdivision, denoted by K_n , is a triangulation of 2^V with exactly $2^{n-1} n!$ facets. Figure 7



FIGURE 8. Barycentric and edgewise subdivisions

shows these subdivisions for $n = 3$ (note that K_3 is combinatorially isomorphic to the visible part of the complex $\text{sd}(\mathcal{C}_3)$ which is shown in Figure 5). We then have [7, Remark 4.5] [108, Theorem 3.1.4]

$$(109) \quad \ell_V(K_n, x) = d_{n,2}^+(x),$$

where $d_{n,2}(x) = d_{n,2}^+(x) + d_{n,2}^-(x)$ is the decomposition of the derangement polynomial for \mathcal{B}_n discussed in Section 2.1.7. Thus, the γ -positivity of $B_n(x)$ and $d_{n,2}^+(x)$, discussed in Section 2, are again instances of Conjectures 3.4 and 3.14, respectively.

The problem to interpret the polynomial $d_{n,r}^+(x)$, discussed in Section 2.1.7, as a local h -polynomial for all $r \geq 1$ was studied in [6] and provided much of the motivation for that paper. It is not clear how to generalize the triangulation K_n for this purpose. However, one can show directly (see [7, Remark 4.5]) that $\ell_V(K_n, x) = \ell_V(\text{esd}_2(\text{sd}(2^V)))$. Thus, the r -fold edgewise subdivision $\text{esd}_r(\text{sd}(2^V))$ of the barycentric subdivision of the simplex 2^V is a reasonable candidate to replace K_n and, indeed, [6, Theorem 1.2] shows that

$$(110) \quad \ell_V(\text{esd}_r(\text{sd}(2^V)), x) = d_{n,r}^+(x).$$

The barycentric subdivision of the two-dimensional simplex and its 2-fold edgewise subdivision are shown in Figure 8. Further examples of barycentric and edgewise subdivisions appear in Section 4.2.

3.3.2. Coxeter complexes. Let W be a finite Coxeter group. Then W can be realized as a reflection group in an n -dimensional Euclidean space, where n is the rank of W . The reflecting hyperplanes form a simplicial arrangement, known as the *Coxeter arrangement*, whose face poset is isomorphic to that of a simplicial complex $\text{Cox}(W)$, called the *Coxeter complex*. This complex is combinatorially isomorphic to the barycentric subdivision of the boundary complex of the n -dimensional simplex or cube (see Figure 5 for two three-dimensional pictures), when W is the symmetric group \mathfrak{S}_{n+1} or the hyperoctahedral group \mathcal{B}_n , respectively.

The facets of $\text{Cox}(W)$ are in a natural one-to-one correspondence with the elements of W . Thus, the h -polynomial of $\text{Cox}(W)$ is an x -analog of the order of the group W and, in fact, as was essentially shown by A. Björner [18] (see [31, Theorem 2.3]), we have

$$(111) \quad h(\text{Cox}(W), x) = W(x),$$

where $W(x)$ is the W -Eulerian polynomial, discussed in Section 2.1.3.

Clearly, $\text{Cox}(W)$ is a triangulation of the $(n-1)$ -dimensional sphere and, as explained in [97, Section 8.5], this triangulation is flag. Thus, the γ -positivity of $W(x)$ (Theorem 2.8) verifies Gal's conjecture in a special case.

3.3.3. Cluster complexes. Let W be a finite Coxeter group of rank n , viewed as a reflection group in an n -dimensional Euclidean space. Consider the root system Φ defined by the Coxeter arrangement associated to W and a choice Φ^+ of a positive system, along with corresponding simple system Π . The cluster complex Δ_W is a flag simplicial complex on the vertex set $\Phi^+ \cup (-\Pi)$, defined by S. Fomin and A. Zelevinsky [60]; we refer to [59] for an overview of the relevant theory and its connections to cluster algebras. This complex triangulates the $(n-1)$ -dimensional sphere; its restriction on the vertex set Φ^+ is naturally a flag triangulation of the simplex 2^Π , termed in [10] as the *cluster subdivision*. One then has [59, Theorem 5.9]

$$(112) \quad h(\Delta_W, x) = \text{Cat}(W, x)$$

and [10, Section 1.1]

$$(113) \quad \ell_\Pi(\Gamma_W, x) = \text{Cat}^{++}(W, x),$$

where $\text{Cat}(W, x)$ and $\text{Cat}^{++}(W, x)$ were defined in Section 2.2. Thus, the γ -positivity of these polynomials (Theorems 2.32 and 2.33) verifies Conjectures 3.4 and 3.14, respectively, in special cases.

4. METHODS

A remarkable variety of methods has been employed to prove γ -positivity and often to describe the γ -coefficients in some explicit way. This section reviews in some detail three such methods, namely valley hopping, methods of geometric combinatorics stemming from Stanley's seminal work [123], and (quasi)symmetric function methods. Although it seems difficult to provide an exhaustive list, or attempt some kind of classification, the author is aware of the following methodological approaches which have been followed successfully to prove the γ -positivity of a polynomial $f(x)$:

- *Combinatorial decompositions:* Assuming $f(x)$ enumerates a set of combinatorial objects according to some statistic, one may try to combinatorially decompose this set into parts, each contributing a binomial $x^i(1+x)^{n-2i}$ to this enumeration. One instance of this approach is valley hopping (see Section 4.1), and another is symmetric Boolean decompositions of posets [88, 95, 117]. Other instances appear in [10, Section 3] [48] [135, Appendix A] (see also [97, Section 13.2]).
- *Explicit combinatorial formulas:* One may try to express $f(x)$ explicitly as a sum of products of polynomials known to be γ -positive, all products having the same center of symmetry, using direct combinatorial arguments, generating functions, continued fraction expansions, and so on. The prototypical example of this approach is Brändén's [24] beautiful proof of Theorem 2.7; see also [39, 115, 116, 135].

- *The theory of the cd-index of an Eulerian poset*: see Section 3.1.1 and [48] [135, Appendix A] for related approaches.
- *Poset shellability and homology methods*: The prototypical example is the use of shellability of Rees products of posets in [84], leading to Theorem 2.37 and related results discussed in Section 2.4; see also Theorem 5.3 in the sequel.
- *Geometric methods*: Apart from those discussed in Section 4.2, geometric methods are employed in [2, 91], where the γ -positivity of $f(x)$ is proved by constructing a simplicial complex whose f -vector is shown to equal the γ -vector of $f(x)$.
- *Recursive methods*: One can prove positivity of the γ -coefficients via recursions, without providing any explicit interpretations or formulas. This is the case with the proof of Theorem 2.12 in [80] and the γ -positivity of the restricted Eulerian polynomials of [92, Section 4].
- *Symmetric and quasisymmetric function methods*: see Section 4.3.

4.1. **Valley hopping.** The idea of valley hopping is due to D. Foata and V. Strehl [57, 58], who used it to interpret combinatorially the γ -coefficients for the Eulerian polynomials (see Theorem 2.1). It was rediscovered by L. Shapiro, W. Woan and S. Getu [109] and has found numerous applications to the enumeration of classes of permutations [10, Section 4] [26, 83] [102, Section 11] [137] and related combinatorial objects [8, Section 3].

To explain this idea, let $w = (w_1, w_2, \dots, w_n) \in \mathfrak{S}_n$, where $w_i = w(i)$ for $1 \leq i \leq n$, be a permutation and set $w_0 = w_{n+1} = \infty$ and $\tilde{w} = (w_0, w_1, \dots, w_{n+1})$. Given a double ascent or double descent i of \tilde{w} (meaning, an index $i \in [n]$ such that $w_{i-1} < w_i < w_{i+1}$ or $w_{i-1} > w_i > w_{i+1}$, respectively), we define the permutation $\psi_i(w) \in \mathfrak{S}_n$ as follows: If i is a double ascent of \tilde{w} , then $\psi_i(w)$ is the permutation obtained from \tilde{w} by deleting w_i and inserting it between w_j and w_{j+1} , where j is the largest index satisfying $0 \leq j < i$ and $w_j > w_i$. Similarly, if i is a double descent of \tilde{w} , then $\psi_i(w)$ is the permutation obtained from \tilde{w} by deleting w_i and inserting it between w_j and w_{j+1} , where j is the smallest index satisfying $i < j \leq n$ and $w_i < w_{j+1}$. For the example of Figure 9 we have $\psi_1(w) = (3, 2, 5, 4, 7, 8, 1, 6)$ and $\psi_8(w) = (7, 3, 2, 5, 4, 8, 6, 1)$.

We call two permutations in \mathfrak{S}_n equivalent if one can be obtained from the other by a sequence of moves of the form $w \mapsto \psi_i(w)$. We leave to the reader to verify that this defines an equivalence relation on \mathfrak{S}_n , each equivalence class of which contains a unique permutation u such that \tilde{u} has no double ascent. Moreover, if \tilde{u} has k double descents for such $u \in \mathfrak{S}_n$, then the equivalence class $O(u)$ of u has 2^k elements and exactly $\binom{k}{j}$ of them have j ascents more than u . Therefore,

$$(114) \quad \sum_{w \in O(u)} x^{\text{asc}(w)} = x^{\text{asc}(u)}(1+x)^k = x^{\text{asc}(u)}(1+x)^{n-1-2\text{asc}(u)}.$$

Summing over all equivalence classes, we obtain the first combinatorial interpretation for $\gamma_{n,i}$, given in Theorem 2.1.




FIGURE 9. Valley hopping for $w = (7, 3, 2, 5, 4, 8, 1, 6)$

To illustrate the power of this method, let us modify it to prove Gessel's identity (88). We will use Stanley's interpretation (see [111, Theorem 7.2])

$$(115) \quad \frac{(1-t)E(\mathbf{x}; z)}{E(\mathbf{x}; tz) - tE(\mathbf{x}; z)} = 1 + \sum_{n \geq 1} z^n \sum_{w \in \mathcal{S}(n)} t^{\text{asc}(w)} \mathbf{x}_w$$

of the left-hand side of (88), where we have used notation of Section 2.5 and $\mathcal{S}(n)$ stands for the set of maps $w : [n] \rightarrow \mathbb{Z}_{>0}$ satisfying $w(i) \neq w(i+1)$ for every $i \in [n-1]$.

Proof of Equation (88). Given Equation (115), it suffices to show that

$$(116) \quad \sum_{w \in \mathcal{S}(n)} t^{\text{asc}(w)} \mathbf{x}_w = \sum_{\substack{u : [n] \rightarrow \mathbb{Z}_{>0} \\ \text{Asc}(u) \in \text{Stab}([n-2])}} t^{\text{asc}(u)} (1+t)^{n-1-2\text{asc}(u)} \mathbf{x}_u$$

for all $n \geq 1$. For $w \in \mathcal{S}(n)$ we write $w = (w_1, w_2, \dots, w_n)$ and define \tilde{w} , as we did for permutations. For a double ascent i of \tilde{w} , we denote by $\psi_i(w)$ the sequence obtained from \tilde{w} by deleting w_i and inserting it between positions j and $j+1$, where j is the largest index satisfying $0 \leq j < i$ and $w_j > w_i$. We define $\psi_i(w)$ in a similar way for double descents i of \tilde{w} and say that a move $w \mapsto \psi_i(w)$ is *valid* if i is a double ascent or descent of \tilde{w} and $\psi_i(w) \in \mathcal{S}(n)$. For the example of Figure 10 we have $\psi_4(w) = (3, 2, 3, 2, 4, 3, 2, 1, 2)$, $\psi_6(w) = (2, 3, 2, 3, 4, 2, 1, 2, 3)$, $\psi_7(w) = (2, 3, 2, 3, 4, 3, 1, 2, 2)$ and $\psi_9(w) = (2, 3, 2, 3, 4, 3, 2, 2, 1)$. The move $w \mapsto \psi_i(w)$ is valid for $i \in \{4, 6\}$ and invalid for $i \in \{7, 9\}$.

We call two elements of $\mathcal{S}(n)$ equivalent if one can be obtained from the other by a sequence of valid moves of the form $w \mapsto \psi_i(w)$. We leave again to the reader to verify that this defines an equivalence relation on $\mathcal{S}(n)$, each equivalence class of which contains a unique element v , call it the minimal representative, such that \tilde{v} has a minimum number of double ascents. For example, the minimal representative of the equivalence class of the sequence shown in Figure 10 is $\psi_4(w)$. The reasoning which led to Equation (114) also




FIGURE 10. Valley hopping for $w = (2, 3, 2, 3, 4, 3, 2, 1, 2)$

shows that the equivalence class $O(v)$ of a minimal representative v satisfies

$$(117) \quad \sum_{w \in O(v)} t^{\text{asc}(w)} \mathbf{x}_w = x^{\text{asc}(v)} (1+x)^{n-1-2\text{asc}(v)} \mathbf{x}_v.$$

Summing over all equivalence classes, we get

$$(118) \quad \sum_{w \in \mathcal{S}(n)} t^{\text{asc}(w)} \mathbf{x}_w = \sum_{v \in \mathcal{S}(n)} t^{\text{asc}(v)} (1+t)^{n-1-2\text{asc}(v)} \mathbf{x}_v,$$

where the sum on the right-hand side runs over all minimal representatives v of equivalence classes. Application of all possible invalid moves to such an element v which shift entries to the left (the order not making a difference) results in a map $u = f(v) : [n] \rightarrow \mathbb{Z}_{>0}$ such that $\text{Asc}(u) \in \text{Stab}([n-2])$ and $\mathbf{x}_u = \mathbf{x}_v$. Moreover, the map f is a bijection from the set of minimal representatives v to the set of such maps u which preserves the weight \mathbf{x}_v and the proof follows. \square

Remark 4.1. The variant of valley hopping which was employed in the previous proof appeared in [8, Section 3] to derive the first interpretation for $\xi_{n,r,i}$, given in Theorem 3.19. The last part of the argument in this proof yields the second interpretation. \square

4.2. Methods of geometric combinatorics. The enumerative theory of [123, Part I], developed by Stanley in order to study h -vectors of triangulations of simplicial complexes, together with recent developments [5], provides a powerful method to prove γ -positivity of polynomials which can often be defined purely in combinatorial terms. This point of view, which is implicit in [7, Section 4] [108, Chapter 3], is further explained in this section. Familiarity with basic definitions from Section 3 is assumed.

The following statement explains the significance of the concept of local h -vector in the theory of [123].

Theorem 4.2 (STANLEY [123, Theorem 3.2]). *For every triangulation Δ' of a pure simplicial complex Δ ,*

$$(119) \quad h(\Delta', x) = \sum_{F \in \Delta} \ell_F(\Delta'_F, x) h(\text{link}_\Delta(F), x).$$

This result implies that $h(\Delta', x)$ is γ -positive (as a sum of γ -positive polynomials with the same center of symmetry), if so are $\ell_F(\Delta'_F, x)$ and $h(\text{link}_\Delta(F), x)$ for every $F \in \Delta$. For instance, we have the following statement.

Corollary 4.3. *Suppose Δ is either:*

- *the simplicial join Σ_n of n copies of the zero-dimensional sphere, or*
- *the barycentric subdivision of the boundary of a simplex.*

Then, $h(\Delta', x)$ is γ -positive for every triangulation Δ' of Δ for which $\ell_F(\Delta'_F, x)$ is γ -positive for every $F \in \Delta$.

Proof. As already discussed, it suffices to confirm that $h(\text{link}_\Delta(F), x)$ is γ -positive for all $F \in \Delta$. This is obvious if $\Delta = \Sigma_n$, since then $\text{link}_\Delta(F)$ is isomorphic to $\Sigma_{n-|F|}$ and hence $h(\text{link}_\Delta(F), x) = (1+x)^{n-|F|}$ for every $F \in \Delta$.

Suppose now that $\Delta = \text{sd}(\partial(2^V))$ is the barycentric subdivision of the boundary complex of an n -dimensional simplex 2^V . Then, faces of Δ have the form $F = \{S_1, S_2, \dots, S_k\}$, where $S_1 \subset S_2 \subset \dots \subset S_k$ are nonempty proper subsets of V . Setting $S_0 := \emptyset$ and $S_{k+1} := V$, we have that $\text{link}_\Delta(F)$ is the simplicial join of the complexes $\text{sd}(\partial(2^{S_i \setminus S_{i-1}}))$ for $i \in [k+1]$ and hence

$$(120) \quad h(\text{link}_\Delta(F), x) = \prod_{i=1}^{k+1} A_{n_i}(x),$$

where $n_i = |S_i \setminus S_{i-1}|$ for $i \in [k+1]$. This polynomial is indeed γ -positive, as a product of γ -positive polynomials. Alternatively, $h(\text{link}_\Delta(F), x)$ can be shown to be γ -positive for the barycentric subdivision Δ of any regular CW-sphere by a stronger version of Theorem 3.8 (applying to Gorenstein* order complexes). \square

Example 4.4. The following explicit formula for the h -polynomial of the r -fold edge-wise subdivision of an $(n-1)$ -dimensional simplicial complex Δ is a consequence of [34, Corollary 1.2] and [35, Corollary 6.8]:

$$(121) \quad h(\text{esd}_r(\Delta), x) = \text{E}_r \left((1+x+x^2+\dots+x^{r-1})^n h(\Delta, x) \right),$$

where the linear operator $\text{E}_r : \mathbb{R}[x] \rightarrow \mathbb{R}[x]$ was defined before Theorem 3.19. Thus, Corollary 4.3, combined with Theorem 3.19, implies the γ -positivity of

$$(122) \quad h(\text{esd}_r(\Sigma_n), x) = \text{E}_r \left((1+x+x^2+\dots+x^{r-1})^n (1+x)^n \right)$$

and

$$(123) \quad h(\text{esd}_r(\text{sd}(\partial(2^V))), x) = \text{E}_r \left((1+x+x^2+\dots+x^{r-1})^n A_{n+1}(x) \right),$$

where 2^V is an n -dimensional simplex. It would be interesting to find explicit combinatorial interpretations for the coefficients of the right-hand sides of Equations (122) and (123) and their corresponding γ -polynomials. Similar remarks apply to the second barycentric subdivision of the boundary complex of a simplex. \square

We now consider the following generalization of the concept of local h -polynomial. Let V be an n -element set, Γ be a triangulation of the simplex 2^V and $E \in \Gamma$.

Definition 4.5 (ATHANASIADIS [5, Remark 3.7] [7, Definition 2.14], NILL–SCHEPERS [93, Section 3]). *The relative local h -polynomial of Γ (with respect to V) at $E \in \Gamma$ is defined as*

$$\ell_V(\Gamma, E, x) = \sum_{\sigma(E) \subseteq F \subseteq V} (-1)^{n-|F|} h(\text{link}_{\Gamma_F}(E), x),$$

where $\sigma(E)$ is the smallest face of 2^V containing E .

This polynomial, which reduces to $\ell_V(\Gamma, x)$ for $E = \emptyset$, has properties similar to those of $\ell_V(\Gamma, x)$. Specifically, it has nonnegative and symmetric coefficients, with center of symmetry $(n - |E|)/2$ (see [7, Theorem 2.15]) and, moreover, it is unimodal if Γ is a regular triangulation [75, Remark 6.5] (see [7, Section 2.3] for more information). The significance of the relative local h -polynomial for us stems from the following statement. Note that, if Γ is a triangulation of the simplex 2^V , then every triangulation Γ' of Γ induces a triangulation of 2^V whose local h -polynomial is denoted by $\ell_V(\Gamma', x)$.

Proposition 4.6 (ATHANASIADIS [5, Section 3] [7, Proposition 2.15 (a)]). *For every triangulation Γ of the simplex 2^V and every triangulation Γ' of Γ ,*

$$(124) \quad \ell_V(\Gamma', x) = \sum_{E \in \Gamma} \ell_E(\Gamma'_E, x) \ell_V(\Gamma, E, x).$$

As a result, $\ell_V(\Gamma', x)$ is γ -positive if so are $\ell_E(\Gamma'_E, x)$ and $\ell_V(\Gamma, E, x)$ for every $E \in \Gamma$. Using an argument similar to the one employed in the proof of Equation (120), one can show (see [108, Example 3.5.2]) that

$$(125) \quad \ell_V(\text{sd}(2^V), E, x) = d_{n_0}(x) \cdot \prod_{i=1}^k A_{n_i}(x),$$

where $E = \{S_1, S_2, \dots, S_k\}$ with $\emptyset \subset S_1 \subset S_2 \subset \dots \subset S_k \subseteq V$ is a face of $\text{sd}(2^V)$, $d_{n_0}(x)$ is a derangement polynomial, $n_0 = |V \setminus S_k|$ and $n_i = |S_i \setminus S_{i-1}|$ for $1 \leq i \leq k$ (with the convention that $S_0 = \emptyset$).

Corollary 4.7. *The polynomial $\ell_V(\Gamma', x)$ is γ -positive for every triangulation Γ' of $\text{sd}(2^V)$ for which $\ell_E(\Gamma'_E, x)$ is γ -positive for every $E \in \text{sd}(2^V)$.*

The special case for which Γ' is the r -fold edgewise subdivision proves the γ -positivity of the polynomial $d_{n,r}^+(x)$ discussed in Section 2.1.7; see [7, Example 4.8]. We give another interesting application.

Example 4.8. Suppose that Γ' is the barycentric subdivision of $\text{sd}(2^V)$ and that V is an n -element set. Then, the induced triangulation of 2^V is the second barycentric subdivision $\text{sd}^{(2)}(2^V)$. An explicit, but rather complicated, formula for $\ell_V(\text{sd}^{(2)}(2^V), x)$ can be derived from the definition of local h -polynomial and [33, Theorem 1]. Proposition 4.6, combined with Equations (107) and (125), implies that

$$(126) \quad \ell_V(\text{sd}^{(2)}(2^V), x) = \sum \binom{n}{n_0, n_1, \dots, n_k} d_k(x) d_{n_0}(x) A_{n_1}(x) \cdots A_{n_k}(x),$$

where the sum ranges over all $k \geq 0$ and over all sequences (n_0, n_1, \dots, n_k) of integers which satisfy $n_0 \geq 0$, $n_1, \dots, n_k \geq 1$ and sum to n . The γ -positivity of the polynomial $\ell_V(\text{sd}^{(2)}(2^V), x)$ follows from this formula (equivalently, from Corollary 4.7).

Using the principle of inclusion-exclusion, one can show that the sum of the coefficients of $\ell_V(\text{sd}^{(2)}(2^V), x)$ equals the number of pairs of permutations $u, v \in \mathfrak{S}_n$ which have no common fixed point. The problem to interpret combinatorially the coefficients of this polynomial and its corresponding local γ -polynomial was posed in [7, Problem 4.12]. \square

4.3. Symmetric/quasisymmetric function methods. Symmetric and quasisymmetric functions provide standard tools for enumeration problems; see, for instance, [17, 85] [125, Chapter 7]. Given a polynomial $f(t)$, to be shown to be γ -positive, one may attempt to find a (quasi)symmetric function which gives $f(t)$ via appropriate specialization. Expansion of this function in some basis of (quasi)symmetric functions may lead to a different formula for $f(t)$ for which γ -positivity is easier to prove. The first instance of this approach seems to be Stembridge's proof of Theorem 2.36 (although a more direct proof, based on Equation (76), is also possible), where the role of the basis of quasisymmetric functions is played by Gessel's basis of fundamental quasisymmetric functions (41).

To illustrate this approach, we sketch proofs of Theorems 2.3, 2.15 and 2.16 which derive these statements from the main result on Eulerian quasisymmetric functions [111] of Shareshian and Wachs, following the reasoning of [114, Section 4]. We use the notation introduced in the beginning of the proof of Proposition 2.22 and set

$$(127) \quad F_{n,S}^*(\mathbf{x}) := F_{n,n-S}(\mathbf{x}) = \sum_{\substack{i_1 \geq i_2 \geq \dots \geq i_n \geq 1 \\ j \in S \Rightarrow i_j > i_{j+1}}} x_{i_1} x_{i_2} \cdots x_{i_n}$$

for $S \subseteq [n-1]$. The principal specializations of this function are given by the formula (see [65, Lemma 5.2])

$$(128) \quad \sum_{m \geq 1} F_{n,S}^*(1, q, \dots, q^{m-1}) p^{m-1} = \frac{p^{|S|} q^{\text{sum}(S)}}{(1-p)(1-pq) \cdots (1-pq^n)}$$

which refines Equation (42), where $\text{sum}(S)$ stands for the sum of the elements of S . We also recall the formula (see [125, Proposition 7.19.12])

$$(129) \quad \sum_{m \geq 1} s_\lambda(1, q, \dots, q^{m-1}) p^{m-1} = \frac{f^\lambda(p, q)}{(1-p)(1-pq) \cdots (1-pq^n)}$$

for the principal specializations of $s_\lambda(\mathbf{x})$, where

$$(130) \quad f^\lambda(p, q) := \sum_{P \in \text{SYT}(\lambda)} p^{\text{des}(P)} q^{\text{maj}(P)}$$

and $\text{maj}(P) := \text{sum}(\text{Des}(P))$ is the major index of P .

Proof of Theorems 2.3 and 2.15. Let us denote by $A_n(p, q, t)$ the left-hand side of Equation (9). Combining the special case $r = 1$ of [111, Theorem 1.2] with Equation (82) and

extracting the coefficient of z^n , we see that

$$(131) \quad \sum_{w \in \mathfrak{S}_n} F_{n, \text{DEX}(w)}(\mathbf{x}) t^{\text{exc}(w)} = \sum_{\lambda \vdash n} P_\lambda(t) s_\lambda(\mathbf{x})$$

for $n \geq 1$. We refer to [111, Section 2] for the definition of $\text{DEX}(w)$; the only properties of this set which are essential here are the facts that $|\text{DEX}(w)| = \text{des}^*(w)$ and that $\text{sum}(\text{DEX}(w)) = \text{maj}(w) - \text{exc}(w)$ for $w \in \mathfrak{S}_n$; see [111, Lemma 2.2]. Thus, taking the generating function of the principal specialization of order m on both sides of (131) and using Equations (128) and (129), we arrive at the identity

$$(132) \quad A_n(p, q, t) = \sum_{\lambda \vdash n} P_\lambda(t) f^\lambda(p, q).$$

By the γ -expansion of $P_\lambda(t)$, given in Corollary 2.41, and the definition of $f^\lambda(p, q)$, this formula implies that

$$(133) \quad A_n(p, q, t) = \sum_{\lambda \vdash n} \sum_{\substack{P, Q \in \text{SYT}(\lambda) \\ \text{Des}(Q) \in \text{Stab}([n-2])}} p^{\text{des}(P)} q^{\text{maj}(P)} t^{\text{des}(Q)} (1+t)^{n-1-\text{des}(Q)}.$$

Use of the Robinson–Schensted correspondence and its properties [125, Lemma 7.23.1] to replace the pair (P, Q) of standard Young tableaux of the same shape with a permutation $w \in \mathfrak{S}_n$ shows that the right-hand sides of Equations (133) and (9) are equal and the proof of Theorem 2.3 follows.

To prove Theorem 2.15, one combines the special case $r = 0$ of [111, Theorem 1.2] with Equation (83) instead to show that

$$(134) \quad \sum_{w \in \mathcal{D}_n} F_{n, \text{DEX}(w)}(\mathbf{x}) t^{\text{exc}(w)} = \sum_{\lambda \vdash n} R_\lambda(t) s_\lambda(\mathbf{x})$$

and then takes principal specialization, as before. □

For the proof of Theorem 2.16 we need the formulas (see [125, Section 7.19]) for the stable principal specializations

$$(135) \quad F_{n,S}^*(1, q, q^2, \dots) = \frac{q^{\text{sum}(S)}}{(1-q)(1-q^2) \cdots (1-q^n)}$$

and

$$(136) \quad s_\lambda(1, q, q^2, \dots) = \frac{f^\lambda(q)}{(1-q)(1-q^2) \cdots (1-q^n)},$$

where $f^\lambda(q) := f^\lambda(1, q) = \sum_{P \in \text{SYT}(\lambda)} q^{\text{maj}(P)}$.

Proof of Theorem 2.16. Let us denote by $A_{n,k}(q, t)$ the left-hand side of Equation (31). Comparing the coefficients of $r^k z^n$ on both sides of [111, Equation (1.3)] and applying

Equation (83), we get

$$\begin{aligned} \sum_{w \in \mathfrak{S}_n: \text{fix}(w)=k} F_{n, \text{DEX}(w)}(\mathbf{x}) t^{\text{exc}(w)} &= h_k(\mathbf{x}) [z^{n-k}] \frac{1-t}{H(\mathbf{x}; tz) - tH(\mathbf{x}; z)} \\ &= h_k(\mathbf{x}) \sum_{\lambda \vdash n-k} R_\lambda(t) s_\lambda(\mathbf{x}). \end{aligned}$$

Taking stable principal specialization, we get

$$\frac{A_{n,k}(q, t)}{(1-q)(1-q^2) \cdots (1-q^n)} = \frac{1}{(1-q) \cdots (1-q^k)} \sum_{\lambda \vdash n-k} R_\lambda(t) \frac{f^\lambda(q)}{(1-q) \cdots (1-q^{n-k})}.$$

Finally, we solve for $A_{n,k}(q, t)$ and specialize to $p = 1$ the expression already obtained for the sum $\sum_{\lambda \vdash n-k} R_\lambda(t) f^\lambda(p, q)$ in the proof of Theorem 2.15 and the proof follows. \square

For other instances of the use of symmetric functions in γ -positivity, see the proof of Theorem 2.42 in [64] and [66, Section 4].

5. GENERALIZATIONS

This section discusses three possible generalizations of γ -positivity, namely nonsymmetric γ -positivity, equivariant γ -positivity and q - γ -positivity, which may provide interesting directions for future research.

5.1. Nonsymmetric γ -positivity. Nonsymmetric polynomials which can be written as a sum of two γ -positive polynomials, whose centers of symmetry differ by $1/2$, have appeared in Section 2. We formalize this situation here as follows.

A nonzero polynomial $f(x) \in \mathbb{R}[x]$ is said to have center $(i+j)/2$, where i (respectively, j) is the smallest (respectively, largest) integer k for which the coefficient of x^k in $f(x)$ is nonzero. One can verify that, if $f(x)$ has center $n/2$, then it can be written uniquely as a sum

$$(137) \quad f(x) = f_\alpha^+(x) + f_\alpha^-(x)$$

of two symmetric polynomials $f_\alpha^+(x), f_\alpha^-(x)$ with centers of symmetry $(n-1)/2$ and $n/2$, respectively. Similarly, $f(x)$ can be written uniquely as a sum

$$(138) \quad f(x) = f_\beta^+(x) + f_\beta^-(x)$$

of two symmetric polynomials $f_\beta^+(x), f_\beta^-(x)$ with centers of symmetry $(n+1)/2$ and $n/2$, respectively. We call $f(x)$ *left γ -positive* (respectively, *right γ -positive*) if both $f_\alpha^+(x)$ and $f_\alpha^-(x)$ (respectively, $f_\beta^+(x)$ and $f_\beta^-(x)$) are γ -positive. We leave it to the reader to verify that $f(x)$ is γ -positive if and only if it is both left and right γ -positive. Clearly, every left γ -positive or right γ -positive polynomial is unimodal, with a peak at its center $n/2$, or at $(n \pm 1)/2$. The derangement polynomials $d_{n,r}(x)$ are left γ -positive by Theorem 2.26, while the left-hand sides of Equations (80) and (81) and h -polynomials of order complexes which triangulate a ball are right γ -positive by Theorems 2.37 and 3.9, respectively. It would be interesting to find other classes of nonsymmetric γ -positive polynomials.

5.2. Equivariant γ -positivity. Let $f(t) = \sum_i a_i t^i \in \mathbb{N}[t]$ be a γ -positive polynomial. It often happens that $a_i = \dim_{\mathbb{C}}(A_i)$ for some non-virtual representations A_i of a finite group G (all representations considered in this section are finite dimensional and defined over the field of complex numbers). Then, it is natural to consider the polynomial $F(t) := \sum_i A_i t^i$, whose coefficients belong to the representation ring of G , as an equivariant analog of $f(t)$ and ask whether

$$(139) \quad F(t) = \sum_{i=0}^{\lfloor n/2 \rfloor} \Gamma_i t^i (1+t)^{n-2i}$$

for some non-virtual G -representations Γ_i , where $n/2$ is the center of symmetry of $f(t)$. We then say that $F(t)$ is γ -positive. For the symmetric group \mathfrak{S}_n , this concept reduces, via the Frobenius characteristic map, to that of Schur γ -positivity [114, Section 3].

Especially interesting is the case where $f(t)$ is the h -polynomial of a flag triangulation of the sphere, or the local h -polynomial of a flag triangulation of the simplex, as pointed out by Shareshian and Wachs [114, Sections 5–6]. The following general discussion assumes familiarity with face rings of simplicial complexes [126, Chapter II] and local face modules of triangulations of simplices [123, Section 4] [126, Section III.10] and avoids technicalities:

- Let Δ be a flag triangulation of the $(n - 1)$ -dimensional sphere, on which a finite group G acts simplicially. Suppose that Θ is a linear system of parameters for the face ring $\mathbb{C}[\Delta]$ such that the linear span of the elements of Θ is G -invariant. Then G acts on each homogeneous component of the graded vector space

$$(140) \quad \mathbb{C}(\Delta) := \mathbb{C}[\Delta]/\langle \Theta \rangle = \bigoplus_{i=0}^n \mathbb{C}(\Delta)_i,$$

whose graded dimension is equal to the h -vector of Δ . The pair (Δ, G) exhibits the *equivariant Gal phenomenon* if there exists Θ for which $\sum_{i=0}^n \mathbb{C}(\Delta)_i t^i$ is γ -positive.

- Let Γ be a flag triangulation of the $(n - 1)$ -dimensional simplex 2^V , on which a subgroup G of the automorphism group of 2^V acts simplicially. Suppose that Θ is a special linear system of parameters, in the sense of [123, Definition 4.2], for the face ring of Γ such that the linear span of the elements of Θ is G -invariant. Then G acts on each homogeneous component of the local face module

$$(141) \quad L_V(\Gamma) = \bigoplus_{i=0}^n L_V(\Gamma)_i$$

defined by Θ , whose graded dimension is equal to the local h -vector of Γ . The pair (Γ, G) exhibits the *local equivariant Gal phenomenon* if there exists Θ for which $\sum_{i=0}^n L_V(\Gamma)_i t^i$ is γ -positive.

When Δ is the boundary complex of an n -dimensional simplicial polytope (more generally, the simplicial complex associated to a complete simplicial fan in \mathbb{R}^n), there exists Θ for which (140) is isomorphic to the cohomology ring of the projective toric variety associated to Δ , as a graded G -representation [43, Section 10]. Shareshian and Wachs [114, Section 5] speculate that, in interesting situations (but not in general) in which Δ is

flag, the pair (Δ, G) exhibits the equivariant Gal phenomenon for such linear system of parameters. For example, as pointed out by these authors, and in view of results of C. Procesi [103] and R. Stanley [122, Proposition 12] [123, Proposition 4.20], the Schur γ -positivity of the coefficient of z^n in the left-hand sides of Equations (82) and (83), already discussed in Section 2.5, are instances of the equivariant Gal phenomenon and its local analog, with corresponding flag triangulations the barycentric subdivisions of the boundary of the $(n - 1)$ -dimensional simplex and the simplex itself. Another instance of the equivariant Gal phenomenon is the Schur γ -positivity of the coefficient of z^n in

$$(142) \quad \frac{(1-t)H(\mathbf{x}; z)H(\mathbf{x}; tz)}{H(\mathbf{x}; tz) - tH(\mathbf{x}; z)},$$

established in [114, Theorem 3.4], with corresponding flag triangulation the boundary complex of the n -dimensional simplicial stellohedron. This result provides an equivariant analog for the γ -positivity of the binomial Eulerian polynomials, discussed in Section 2.1.

A very interesting example is the Coxeter complex $\text{Cox}(W)$, discussed in Section 3.3.2, on which the finite Coxeter group W acts. When W is crystallographic, its representation on the cohomology of the projective toric variety associated to $\text{Cox}(W)$ was studied by C. Procesi [103], R. Stanley [122, p. 529], I. Dolgachev and V. Lunts [49], J. Stembridge [132], G. Lehrer [78] and A. Stapledon [128, Section 8]. Given the evidence provided in the sequel, it is expected that the following conjectural equivariant analog of Theorem 2.8 can be shown using the classification of finite Coxeter groups (a proof which does not use the classification is certainly more desirable).

Conjecture 5.1. *The pair $(\text{Cox}(W), W)$ exhibits the equivariant Gal phenomenon for every finite crystallographic Coxeter group W .*

As mentioned earlier, in the symmetric group case the conjecture follows from the γ -positivity of the polynomials $P_\lambda(t)$, shown in Corollary 2.41. In the case of the hyperoctahedral group \mathcal{B}_n , by [49, Theorem 6.3] or [132, Theorem 7.6], the Frobenius characteristic of the graded \mathcal{B}_n -representation on the (even degree) cohomology of the projective toric variety associated to $\text{Cox}(\mathcal{B}_n)$ is equal to the coefficient of z^n in

$$(143) \quad \frac{(1-t)H(\mathbf{x}; z)H(\mathbf{x}; tz)}{H(\mathbf{x}, \mathbf{y}; tz) - tH(\mathbf{x}, \mathbf{y}; z)},$$

where $\mathbf{y} = (y_1, y_2, \dots)$ is another sequence of commuting independent indeterminates and $H(\mathbf{x}, \mathbf{y}; z) = \sum_{n \geq 0} h_n(\mathbf{x}, \mathbf{y})z^n = H(\mathbf{x}; z)H(\mathbf{y}; z)$ is the generating function of the complete homogeneous symmetric functions $h_n(\mathbf{x}, \mathbf{y})$ in the variable z . Thus, Conjecture 5.1 in this case is implied by the following result (the proof, which uses Theorem 5.3 stated in the sequel, will appear in the final version of [9] or elsewhere).

Proposition 5.2. *The coefficient of z^n in (143) is Schur γ -positive for every $n \in \mathbb{N}$.*

For example, writing

$$\frac{(1-t)H(\mathbf{x}; z)H(\mathbf{x}; tz)}{H(\mathbf{x}, \mathbf{y}; tz) - tH(\mathbf{x}, \mathbf{y}; z)} = 1 + \sum_{n \geq 1} z^n \sum_{i=0}^{\lfloor n/2 \rfloor} \gamma_{n,i}^{\mathcal{B}}(\mathbf{x}, \mathbf{y}) t^i (1+t)^{n-2i},$$

we have

$$\begin{aligned}\gamma_{1,0}^B(\mathbf{x}, \mathbf{y}) &= s_{(1)}(\mathbf{x}), \\ \gamma_{2,0}^B(\mathbf{x}, \mathbf{y}) &= s_{(2)}(\mathbf{x}), \\ \gamma_{2,1}^B(\mathbf{x}, \mathbf{y}) &= s_{(1,1)}(\mathbf{x}) + s_{(1)}(\mathbf{x})s_{(1)}(\mathbf{y}) + s_{(2)}(\mathbf{y}), \\ \gamma_{3,0}^B(\mathbf{x}, \mathbf{y}) &= s_{(3)}(\mathbf{x}), \\ \gamma_{3,1}^B(\mathbf{x}, \mathbf{y}) &= 2s_{(2,1)}(\mathbf{x}) + s_{(1,1)}(\mathbf{x})s_{(1)}(\mathbf{y}) + 2s_{(2)}(\mathbf{x})s_{(1)}(\mathbf{y}) + 2s_{(1)}(\mathbf{x})s_{(2)}(\mathbf{y}) + s_{(3)}(\mathbf{y}).\end{aligned}$$

At present, a combinatorial interpretation of the coefficients of the functions $\gamma_{n,i}^B(\mathbf{x}, \mathbf{y})$ in the Schur basis is lacking.

Other evidence for the validity of Conjecture 5.1 is provided by the computation of the graded multiplicity of the trivial [132, Section 3] and of the sign [78, Theorem 3.5 (iii)] and reflection representation [78, Section 4] of W in the cohomology of the toric variety associated to $\text{Cox}(W)$; see also [128, Remark 8.3].

A general result on equivariant (nonsymmetric) γ -positivity is the following equivariant version of Theorem 2.37. Let \mathcal{P} be a finite graded poset, as in Section 2.4, and assume that a finite group G acts on \mathcal{P} by order preserving bijections. Then G defines a permutation representation $\alpha_{\mathcal{P}}(S)$, induced by its action on the set of maximal chains of the rank-selected subposet \mathcal{P}_S , for every $S \subseteq [n]$. One can consider the virtual G -representation

$$(144) \quad \beta_{\mathcal{P}}(S) = \sum_{T \subseteq S} (-1)^{|S-T|} \alpha_{\mathcal{P}}(T),$$

introduced by Stanley [121]. When \mathcal{P} is Cohen–Macaulay over \mathbb{C} , $\beta_{\mathcal{P}}(S)$ coincides with the non-virtual G -representation induced on the top homology group of $\bar{\mathcal{P}}_S$ (see [121] [140, Section 3.4]); the dimensions of $\alpha_{\mathcal{P}}(S)$ and $\beta_{\mathcal{P}}(S)$ are equal to the numerical invariants $a_{\mathcal{P}}(S)$ and $b_{\mathcal{P}}(S)$, respectively, defined in Section 2.4.

Theorem 5.3 (ATHANASIADIS [9, Theorem 1.2]). *Let G be a finite group acting on a finite bounded graded poset \mathcal{P} of rank $n + 1$ by order preserving bijections. If \mathcal{P} is Cohen–Macaulay over \mathbb{C} , then*

$$(145) \quad \begin{aligned} \tilde{H}_{n-1}((\mathcal{P}^- * T_{t,n})_-; \mathbb{C}) \cong_G & \sum_{S \in \text{Stab}([n-1])} \beta_{\mathcal{P}}([n] \setminus S) t^{|S|} (1+t)^{n-2|S|} \\ & + \sum_{S \in \text{Stab}([n-2])} \beta_{\mathcal{P}}([n-1] \setminus S) t^{|S|+1} (1+t)^{n-1-2|S|} \end{aligned}$$

and

$$(146) \quad \begin{aligned} \tilde{H}_{n-1}(\bar{\mathcal{P}} * T_{t,n-1}; \mathbb{C}) \cong_G & \sum_{S \in \text{Stab}([2,n-2])} \beta_{\mathcal{P}}([n-1] \setminus S) t^{|S|+1} (1+t)^{n-2-2|S|} \\ & + \sum_{S \in \text{Stab}([2,n-1])} \beta_{\mathcal{P}}([n] \setminus S) t^{|S|} (1+t)^{n-1-2|S|} \end{aligned}$$

for every positive integer t , where \cong_G denotes isomorphism of G -representations.

In particular, the left-hand sides of (145) and (146) are right γ -positive.

As shown in [9, Corollary 4.1], the Schur γ -positivity of the coefficients of the left-hand sides of Equations (86) and (87) follows from the special case of Theorem 5.3 in which \mathcal{P}^- is the Boolean lattice of subsets of $[n]$. Two more applications, establishing the Schur γ -positivity of the coefficients of two close relatives to (143), are given in [9, Section 4]. We state these results here in a form similar to that of Gessel's identities (86)–(89) as follows (a more elementary proof should be possible). For a map $w : [n] \rightarrow \mathbb{Z} \setminus \{0\}$ we write $\mathbf{z}_w := z_{w(1)}z_{w(2)} \cdots z_{w(n)}$, where $z_{w(i)} = x_{w(i)}$ if $w(i) > 0$ and $z_{w(i)} = y_{w(i)}$ otherwise, and define $\text{Asc}_B(w)$ as the set of indices $i \in [n]$ such that $w(i) >_r w(i+1)$ in the total order (17), where $w(n+1) := 0$.

Theorem 5.4 (cf. [9, Corollaries 4.4 and 4.7]). *We have*

$$(147) \quad \frac{(1-t)E(\mathbf{x}; z)E(\mathbf{x}; tz)E(\mathbf{y}; tz)}{E(\mathbf{x}, \mathbf{y}; tz) - tE(\mathbf{x}, \mathbf{y}; z)} = 1 + \sum_{n \geq 1} z^n \sum_w t^{\text{asc}(w)} (1+t)^{n+1-2\text{asc}(w)} \mathbf{z}_w \\ + \sum_{n \geq 1} z^n \sum_w t^{\text{asc}(w)} (1+t)^{n-2\text{asc}(w)} \mathbf{z}_w,$$

where, in the two sums, $w : [n] \rightarrow \mathbb{Z} \setminus \{0\}$ runs through all maps for which $\text{Asc}_B(w) \in \text{Stab}([n])$ contains (respectively, does not contain) n , and

$$(148) \quad \frac{(1-t)E(\mathbf{x}; z)}{E(\mathbf{x}, \mathbf{y}; tz) - tE(\mathbf{x}, \mathbf{y}; z)} = 1 + \sum_{n \geq 1} z^n \sum_w t^{\text{asc}(w)} (1+t)^{n-2\text{asc}(w)} \mathbf{z}_w \\ + \sum_{n \geq 1} z^n \sum_w t^{\text{asc}(w)} (1+t)^{n-1-2\text{asc}(w)} \mathbf{z}_w,$$

where, in the two sums, $w : [n] \rightarrow \mathbb{Z} \setminus \{0\}$ runs through all maps for which $\text{Asc}_B(w) \in \text{Stab}([2, n])$ contains (respectively, does not contain) n .

5.3. q - γ -positivity. Let $a, b \in \mathbb{N}$. A polynomial $f(q, x) \in \mathbb{R}[q, x]$ is called q - γ -positive of type (a, b) if

$$(149) \quad f(q, x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \gamma_i(q) x^i \prod_{k=i}^{n-1-i} (1 + xq^{ka+b})$$

for some polynomials $\gamma_i(q) \in \mathbb{R}[q]$ with nonnegative coefficients. This concept, which reduces to that of γ -positivity for $q = 1$, was formally introduced by K. Dilks [47, Chapter 4] and, in more restrictive form, by C. Krattenthaler and M. Wachs (unpublished), after such expansions were found for certain q -analogs of the Eulerian polynomials for symmetric and hyperoctahedral groups by G. Han, F. Jouhet and J. Zeng [70]. Other polynomials which admit q -analogs which are q - γ -positive include binomials and Narayana polynomials for symmetric and hyperoctahedral groups; see [47, Chapter 4].

Acknowledgements. The author wishes to thank Francesco Brenti, Persi Diaconis, Satoshi Murai, Kyle Petersen, Richard Stanley, John Stembridge, Vasu Tewari and Jiang

Zeng for useful comments and information and the SLC organizers Christian Krattenthaler, Volker Strehl and Jean-Yves Thibon for their invitation.

Note added in revision. Conjecture 2.30 has been proven by P. Brändén and L. Solus [28, Section 3.3] and, independently, by N. Gustafsson and L. Solus [69, Section 5.1].

REFERENCES

- [1] R.M. Adin, C.A. Athanasiadis, S. Elizalde and Y. Roichman, *Character formulas and descents for the hyperoctahedral group*, Adv. Appl. Math. **87** (2017), 128–169.
- [2] N. Aisbett, *Frankl–Füredi–Kalai inequalities for the γ -vectors of flag nestohedra*, Discrete Comput. Geom. **51** (2014), 323–336.
- [3] D. Armstrong, *Generalized noncrossing partitions and combinatorics of Coxeter groups*, Mem. Amer. Math. Soc. **202** (2009), no. 949.
- [4] C.A. Athanasiadis, *Some combinatorial properties of flag simplicial pseudomanifolds and spheres*, Ark. Mat. **49** (2011), 17–29.
- [5] C.A. Athanasiadis, *Flag subdivisions and γ -vectors*, Pacific J. Math. **259** (2012), 257–278.
- [6] C.A. Athanasiadis, *Edgewise subdivisions, local h -polynomials and excedances in the wreath product $\mathbb{Z}_r \wr \mathfrak{S}_n$* , SIAM J. Discrete Math. **28** (2014), 1479–1492.
- [7] C.A. Athanasiadis, *A survey of subdivisions and local h -vectors*, in *The Mathematical Legacy of Richard P. Stanley* (P. Hersh, T. Lam, P. Pylyavskyy, V. Reiner, eds.), Amer. Math. Society, Providence, R.I., 2016, pp. 39–51.
- [8] C.A. Athanasiadis, *The local h -polynomial of the edgewise subdivision of the simplex*, Bull. Hellenic Math. Soc. (N.S.) **60** (2016), 11–19.
- [9] C.A. Athanasiadis, *Some applications of Rees products of posets to equivariant gamma-positivity*, arXiv:1707.08297.
- [10] C.A. Athanasiadis and C. Savvidou, *The local h -vector of the cluster subdivision of a simplex*, Sémin. Lothar. Combin. **66** (2012), Article B66c, 21 pp. (electronic).
- [11] E. Bagno and R. Biagioli, *Colored descent representations for complex reflection groups*, Israel J. Math. **160** (2007), 317–347.
- [12] E. Bagno and D. Garber, *On the excedance number of colored permutation groups*, Sémin. Lothar. Combin. **53** (2006), Article B53f, 17 pp. (electronic).
- [13] M. Barnabei, F. Bonetti and M. Silimbani, *The descent statistic in involutions is not log-concave*, European J. Combin. **30** (2009), 11–16.
- [14] R. Barnard and N. Reading, *Coxeter-biCatalan combinatorics*, J. Algebr. Combin. **47** (2018), 241–300.
- [15] D. Barnette, *A proof of the lower bound conjecture for convex polytopes*, Pacific J. Math. **46** (1973), 349–354.
- [16] D. Bessis, *The dual braid monoid*, Ann. Sci. Ecole Norm. Sup. **36** (2003), 647–683.
- [17] S.C. Billey and P.R.W. McNamara, *The contributions of Stanley to the fabric of symmetric and quasisymmetric functions*, in *The Mathematical Legacy of Richard P. Stanley* (P. Hersh, T. Lam, P. Pylyavskyy, V. Reiner, eds.), Amer. Math. Society, Providence, R.I., 2016, pp. 83–104.
- [18] A. Björner, *Some combinatorial and algebraic properties of Coxeter complexes and Tits buildings*, Adv. Math. **52** (1984), 173–212.
- [19] A. Björner, *Topological methods*, in *Handbook of combinatorics* (R.L. Graham, M. Grötschel and L. Lovász, eds.), North Holland, Amsterdam, 1995, pp. 1819–1872.
- [20] A. Björner and F. Brenti, *Combinatorics of Coxeter groups*, Graduate Texts in Mathematics **231**, Springer, 2005.
- [21] A. Björner and V. Welker, *Segre and Rees products of posets, with ring-theoretic applications*, J. Pure Appl. Algebra **198** (2005), 43–55.

- [22] S. Blanco and T.K. Petersen, *Counting Dyck paths by area and rank*, Ann. Combin. **18** (2014), 171–197.
- [23] T. Brady and C. Watt, *$K(\pi, 1)$'s for Artin groups of finite type*, in *Proceedings of the Conference on Geometric and Combinatorial group theory*, Part I (Haifa 2000), Geom. Dedicata **94** (2002), 225–250.
- [24] P. Brändén, *Sign-graded posets, unimodality of W -polynomials and the Charney–Davis conjecture*, in *Stanley Festschrift* (B.E. Sagan, ed.), Electron. J. Combin. **11** (2004), no. 2, Research Paper 9, 15 pp. (electronic).
- [25] P. Brändén, *Counterexamples to the Neggers–Stanley conjecture*, Electron. Res. Announc. Amer. Math. Soc. **10** (2004), 155–158 (electronic).
- [26] P. Brändén, *Actions on permutations and unimodality of descent polynomials*, European J. Combin. **29** (2008), 514–531.
- [27] P. Brändén, *Unimodality, log-concavity, real-rootedness and beyond*, in *Handbook of Combinatorics* (M. Bona, ed.), CRC Press, 2015, pp. 437–483.
- [28] P. Brändén and L. Solus, *Symmetric decompositions and real-rootedness*, arXiv:1808.04141.
- [29] F. Brenti, *Unimodal, log-concave and Pólya frequency sequences in combinatorics*, Mem. Amer. Math. Soc., no. 413, 1989.
- [30] F. Brenti, *Unimodal polynomials arising from symmetric functions*, Proc. Amer. Math. Soc. **108** (1990), 1133–1141.
- [31] F. Brenti, *q -Eulerian polynomials arising from Coxeter groups*, European J. Combin. **15** (1994), 417–441.
- [32] F. Brenti, *Log-concave and unimodal sequences in algebra, combinatorics and geometry: an update*, in *Jerusalem Combinatorics '93* (H. Barcelo and G. Kalai, eds.), Contemp. Math. **178**, Amer. Math. Society, Providence, R.I., 1994, pp. 71–89.
- [33] F. Brenti and V. Welker, *f -Vectors of barycentric subdivisions*, Math. Z. **259** (2008), 849–865.
- [34] F. Brenti and V. Welker, *The Veronese construction for formal power series and graded algebras*, Adv. Appl. Math. **42** (2009), 545–556.
- [35] M. Brun and T. Römer, *Subdivisions of toric complexes*, J. Algebr. Combin. **21** (2005), 423–448.
- [36] V.M. Buchstaber and T.E. Panov, *Toric Topology*, Mathematical Surveys and Monographs **204**, Amer. Math. Society, 2015.
- [37] R. Charney and M. Davis, *The Euler characteristic of a nonpositively curved, piecewise Euclidean manifold*, Pacific J. Math. **171** (1995), 117–137.
- [38] W.Y.C. Chen, R.L. Tang and A.F.Y. Zhao, *Derangement polynomials and excedances of type B* , Electron. J. Combin. **16** (2) (2009), Research Paper 15, 16 pp. (electronic).
- [39] C.-O. Chow, *On certain combinatorial expansions of the Eulerian polynomials*, Adv. Appl. Math. **41** (2008), 133–157.
- [40] C.-O. Chow, *On derangement polynomials of type B . II*, J. Combin. Theory Ser. A **116** (2009), 816–830.
- [41] C.-O. Chow and T. Mansour, *Counting derangements, involutions and unimodal elements in the wreath product $C_r \wr \mathfrak{S}_n$* , Israel J. Math. **179** (2010), 425–448.
- [42] F. Chung, R. Graham and D. Knuth, *A symmetrical Eulerian identity*, J. Combin. **1** (2010), 29–38.
- [43] V.I. Danilov, *The geometry of toric varieties*, Russian Math. Surveys **33** (1978), 97–154; translated from Uspekhi Mat. Nauk. **33** (1978), 85–134.
- [44] M. Davis and B. Okun, *Vanishing theorems and conjectures for the L^2 -homology of right-angled Coxeter groups*, Geom. Topol. **5** (2001), 7–74.
- [45] J.A. De Loera, J. Rambau and F. Santos, *Triangulations: Structures for Algorithms and Applications*, Algorithms and Computation in Mathematics **25**, Springer, 2010.
- [46] J. Désarménien and D. Foata, *Fonctions symétriques et séries hypergéométriques basiques multivariées*, Bull. Soc. Math. France **113** (1985), 3–22.

- [47] K. Dilks, *Involutions on Baxter objects, and q -gamma nonnegativity*, Ph.D. thesis, University of Minnesota, 2015.
- [48] K. Dilks, T.K. Petersen and J.R. Stembridge, *Affine descents and the Steinberg torus*, Adv. Appl. Math. **42** (2009), 423–444.
- [49] I. Dolgachev and V. Lunts, *A character formula for the representation of a Weyl in the cohomology of the associated toric variety*, J. Algebra **168** (1994), 741–772.
- [50] Ö. Eğecioğlu and J.B. Remmel, *Bijections for Cayley trees, spanning trees, and their q -analogues*, J. Combin. Theory Ser. A **42** (1986), 15–30.
- [51] R. Ehrenborg and K. Karu, *Decomposition theorem for the cd -index of Gorenstein* posets*, J. Algebr. Combin. **26** (2007), 225–251.
- [52] E.-M. Feichtner and B. Sturmfels, *Matroid polytopes, nested sets and Bergman fans*, Port. Math. (N.S.) **62** (2005), 437–468.
- [53] E.-M. Feichtner and S. Yuzvinsky, *Chow rings of toric varieties defined by atomic lattices*, Invent. Math. **155** (2004), 515–536.
- [54] D. Foata and G.-N. Han, *The decrease value theorem with an application to permutation statistics*, Adv. Appl. Math. **46** (2011), 296–311.
- [55] D. Foata, *Groupes de réarrangements et nombres d’Euler*, C.R. Acad. Sci. Paris Sér. A–B **275** (1972), 1147–1150.
- [56] D. Foata and M.-P. Schützenberger, *Théorie Géométrique des Polynômes Eulériens*, Lecture Notes in Mathematics **138**, Springer-Verlag, 1970.
- [57] D. Foata and V. Strehl, *Rearrangements of the symmetric group and enumerative properties of the tangent and secant numbers*, Math. Z. **137** (1974), 257–264.
- [58] D. Foata and V. Strehl, *Euler numbers and variations of permutations*, in *Colloquio Internazionale sulle Teorie Combinatorie (Roma, 1973)*, Tomo I, Atti dei Convegni Lincei, No. 17, Accad. Naz. Lincei, Rome, 1976, pp. 119–131.
- [59] S. Fomin and N. Reading, *Root systems and generalized associahedra*, in *Geometric Combinatorics* (E. Miller, V. Reiner and B. Sturmfels, eds.), IAS/Park City Mathematics Series **13**, pp. 389–496, Amer. Math. Society, Providence, R.I., 2007.
- [60] S. Fomin and A.V. Zelevinsky, *Y -systems and generalized associahedra*, Ann. of Math. **158** (2003), 977–1018.
- [61] Ś.R. Gal, *Real root conjecture fails for five- and higher-dimensional spheres*, Discrete Comput. Geom. **34** (2005), 269–284.
- [62] V. Gasharov, *On the Neggers–Stanley conjecture and the Eulerian polynomials*, J. Combin. Theory Ser. A **82** (1998), 134–146.
- [63] I.M. Gessel, *Counting forests by descents and leaves*, Electron. J. Combin. **3**(2) (1996), Research Paper 8, 5 pp. (electronic).
- [64] I.M. Gessel, S. Griffin and V. Tewari, *Labeled plane binary trees and Schur-positivity*, [arXiv:1706.03055](https://arxiv.org/abs/1706.03055).
- [65] I.M. Gessel and C. Reutenauer, *Counting permutations with given cycle structure and descent set*, J. Combin. Theory Ser. A **64** (1993), 189–215.
- [66] R.S. González D’León, *A note on the γ -coefficients of the tree Eulerian polynomial*, Electron. J. Combin. **23** (2016), Research Paper 1.20, 13 pp. (electronic).
- [67] M.A. Gorsky, *Proof of Gal’s conjecture for the D series of generalized associahedra*, Russian Math. Surveys **65** (2010), 1178–1180.
- [68] V.W.J. Guo and J. Zeng, *The Eulerian distribution on involutions is indeed unimodal*, J. Combin. Theory Ser. A **113** (2006), 1061–1071.
- [69] N. Gustafsson and L. Solus, *Derangements, Ehrhart theory and local h -polynomials*, [arXiv:1807.05246](https://arxiv.org/abs/1807.05246).

- [70] G. Han, F. Jouhet and J. Zeng, *Two new triangles of q -integers via q -Eulerian polynomials of type A and B* , Ramanujan J. **31** (2013), 115–127.
- [71] A. Hultman, *The combinatorics of twisted involutions in Coxeter groups*, Trans. Amer. Math. Soc. **359** (2007), 2787–2798.
- [72] I.M. Isaacs, *Character Theory of Finite Groups*, Pure and Applied Mathematics, Academic Press, 1976.
- [73] M. Juhnke-Kubitzke, S. Murai and R. Sieg, *Local h -vectors of quasi-geometric and barycentric subdivisions*, Discrete Comput. Geom. (to appear).
- [74] K. Karu, *The cd -index of fans and posets*, Compos. Math. **142** (2006), 701–718.
- [75] E. Katz and A. Slapledon, *Local h -polynomials, invariants of subdivisions and mixed Ehrhart theory*, Adv. Math. **286** (2016), 181–239.
- [76] V. Klee, *A combinatorial analogue of Poincaré’s duality theorem*, Canad. J. Math. **16** (1964), 517–531.
- [77] D. Kozlov, *Combinatorial Algebraic Topology, Algorithms and Computation in Mathematics* **21**, Springer, 2008.
- [78] G.I. Lehrer, *Rational points and Coxeter group actions on the cohomology of toric varieties*, Ann. Inst. Fourier (Grenoble) **58** (2008), 671–688.
- [79] Z. Lin, *Eulerian calculus arising from permutation statistics*, Doctoral Dissertation, Université Claude Bernard-Lyon 1, 2014.
- [80] Z. Lin, *Proof of Gessel’s γ -positivity conjecture*, Electron. J. Combin. **23** (2016), Research Paper 3.15, 9 pp. (electronic).
- [81] Z. Lin, *On γ -positive polynomials arising in pattern avoidance*, Adv. Appl. Math. **82** (2017), 1–22.
- [82] Z. Lin and S. Fu, *On 1212-avoiding restricted growth functions*, Electron. J. Combin. **24** (2017), Research Paper 1.53, 20 pp. (electronic).
- [83] Z. Lin and J. Zeng, *The γ -positivity of basic Eulerian polynomials via group actions*, J. Combin. Theory Ser. A **135** (2015), 112–129.
- [84] S. Linusson, J. Shareshian and M.L. Wachs, *Rees products and lexicographic shellability*, J. Combin. **3** (2012), 243–276.
- [85] A. Mendes and J. Remmel, *Counting with Symmetric Functions*, Developments in Mathematics **43**, Springer, 2015.
- [86] P. Mongelli, *Excedances in classical and affine Weyl groups*, J. Combin. Theory Ser. A **120** (2013), 1216–1234.
- [87] V-D.P. Moustakas, *The Eulerian distribution on the involutions of the hyperoctahedral group (in Greek)*, Master Thesis, University of Athens, 2017.
- [88] H. Mühle, *Symmetric decompositions and the strong Sperner property for noncrossing partition lattices*, J. Algebr. Combin. **45** (2017), 745–775.
- [89] J. Neggers, *Representations of finite partially ordered sets*, J. Combin. Inform. Syst. Sci. **3** (1978), 113–133.
- [90] E. Nevo, *Remarks on missing faces and generalized lower bounds on face numbers*, Electron. J. Combin. **16** (2009), no. 2, Research Paper 8, 11 pp. (electronic).
- [91] E. Nevo and T.K. Petersen, *On gamma vectors satisfying the Kruskal–Katona inequalities*, Discrete Comput. Geom. **45** (2011), 503–521.
- [92] E. Nevo, T.K. Petersen and B.E. Tenner, *The gamma-vector of a barycentric subdivision*, J. Combin. Theory Ser. A **118** (2011), 1364–1380.
- [93] B. Nill and J. Schepers, *Combinatorial questions related to stringy E -polynomials of Gorenstein polytopes*, in *Toric Geometry* (K. Altmann et al., eds.), Oberwolfach Reports (2012), no. 21, pp. 62–64.
- [94] T.K. Petersen, *Enriched P -partitions and peak algebras*, Adv. Math. **209** (2007), 561–610.

- [95] T.K. Petersen, *On the shard intersection order of a Coxeter group*, SIAM J. Discrete Math. **27** (2013), 1880–1912.
- [96] T.K. Petersen, *Two-sided Eulerian numbers via balls in boxes*, Math. Mag. **86** (2013), 159–176.
- [97] T.K. Petersen, *Eulerian Numbers*, Birkhäuser Advanced Texts, Birkhäuser, 2015.
- [98] T.K. Petersen, *A two-sided analogue of the Coxeter complex*, arXiv:1607.00086.
- [99] S. Poirier, *Cycle type and descent set in wreath products*, Discrete Math. **180** (1998), 315–343.
- [100] A. Postnikov, *Intransitive trees*, J. Combin. Theory Ser. A **79** (1997), 360–366.
- [101] A. Postnikov, *Permutohedra, associahedra, and beyond*, Int. Math. Res. Not. **2009** (2009), 1026–1106.
- [102] A. Postnikov, V. Reiner and L. Williams, *Faces of generalized permutohedra*, Doc. Math. **13** (2008), 207–273.
- [103] C. Procesi, *The toric variety associated to Weyl chambers*, in *Mots* (M. Lothaire, ed.), Hermès, Paris, 1990, pp. 153–161.
- [104] V. Reiner and V. Welker, *On the Charney–Davis and Neggers–Stanley conjectures*, J. Combin. Theory Ser. A **109** (2005), 247–280.
- [105] G. Reisner, *Cohen–Macaulay quotients of polynomial rings*, Adv. Math. **21** (1976), 30–49.
- [106] Y. Roichman, *A note on the number of k -roots in S_n* , Sémin. Lothar. Combin. **70** (2014), Article B70i, 5 pp. (electronic).
- [107] C.D. Savage and M. Visontai, *The s -Eulerian polynomials have only real roots*, Trans. Amer. Math. Soc. **367** (2015), 763–788.
- [108] C. Savvidou, *Barycentric subdivisions, clusters and permutation enumeration* (in Greek), Doctoral Dissertation, University of Athens, 2013.
- [109] L.W. Shapiro, W.J. Woan and S. Getu, *Runs, slides and moments*, SIAM J. Algebraic Discrete Methods **4** (1983), 459–466.
- [110] J. Shareshian and M.L. Wachs, *Poset homology of Rees products and q -Eulerian polynomials*, Electron. J. Combin. **16** (2) (2009), Research Paper 20, 29 pp. (electronic).
- [111] J. Shareshian and M.L. Wachs, *Eulerian quasisymmetric functions*, Adv. Math. **225** (2010), 2921–2966.
- [112] J. Shareshian and M.L. Wachs, *Chromatic quasisymmetric functions*, Adv. Math. **295** (2016), 497–551.
- [113] J. Shareshian and M.L. Wachs, *From poset topology to q -Eulerian polynomials to Stanley’s chromatic symmetric functions*, in *The Mathematical Legacy of Richard P. Stanley* (P. Hersh, T. Lam, P. Pylyavskyy and V. Reiner, eds.), pp. 301–321, Amer. Math. Society, Providence, R.I., 2016.
- [114] J. Shareshian and M.L. Wachs, *Gamma-positivity of variations of Eulerian polynomials*, arXiv:1702.06666v3.
- [115] H. Shin and J. Zeng, *The symmetric and unimodal expansion of Eulerian polynomials via continued fractions*, European J. Combin. **33** (2012), 111–127.
- [116] H. Shin and J. Zeng, *Symmetric unimodal expansions of excedances in colored permutations*, European J. Combin. **52** (2016), 174–196.
- [117] R. Simion and D. Ullman, *On the structure of the lattice of noncrossing partitions*, Discrete Math. **98** (1991), 193–206.
- [118] R.P. Stanley, *Ordered structures and partitions*, Mem. Amer. Math. Soc., no. 119, 1972.
- [119] R.P. Stanley, *The Upper Bound Conjecture and Cohen–Macaulay rings*, Studies in Appl. Math. **54** (1975), 135–142.
- [120] R.P. Stanley, *The number of faces of a simplicial convex polytope*, Adv. Math. **35** (1980), 236–238.
- [121] R.P. Stanley, *Some aspects of groups acting on finite posets*, J. Combin. Theory Ser. A **32** (1982), 132–161.

- [122] R.P. Stanley, *Log-concave and unimodal sequences in algebra, combinatorics, and geometry*, in *Graph Theory and its Applications: East and West*, Annals of the New York Academy of Sciences **576**, New York Acad. Sci., New York, 1989, pp. 500–535.
- [123] R.P. Stanley, *Subdivisions and local h -vectors*, J. Amer. Math. Soc. **5** (1992), 805–851.
- [124] R.P. Stanley, *Flag f -vectors and the cd -index*, Math. Z. **216** (1994), 483–499.
- [125] R.P. Stanley, *Enumerative Combinatorics*, vol. 2, Cambridge Studies in Advanced Mathematics **62**, Cambridge University Press, Cambridge, 1999.
- [126] R.P. Stanley, *Combinatorics and Commutative Algebra*, second edition, Birkhäuser, Basel, 1996.
- [127] R.P. Stanley, *Enumerative Combinatorics*, vol. 1, Cambridge Studies in Advanced Mathematics **49**, Cambridge University Press, second edition, Cambridge, 2011.
- [128] A. Stapledon, *Equivariant Ehrhart theory*, Adv. Math. **226** (2011), 3622–3654.
- [129] E. Steingrímsson, *Permutation statistics of indexed and poset permutations*, Ph.D. thesis, MIT, 1992.
- [130] E. Steingrímsson, *Permutation statistics of indexed permutations*, European J. Combin. **15** (1994), 187–205.
- [131] J.R. Stembridge, *Eulerian numbers, tableaux, and the Betti numbers of a toric variety*, Discrete Math. **99** (1992), 307–320.
- [132] J.R. Stembridge, *Some permutation representations of Weyl groups associated with the cohomology of toric varieties*, Adv. Math. **106** (1994), 244–301.
- [133] J.R. Stembridge, *Enriched P -partitions*, Trans. Amer. Math. Soc. **349** (1997), 763–788.
- [134] J.R. Stembridge, *Counterexamples to the poset conjectures of Neggers, Stanley, and Stembridge*, Trans. Amer. Math. Soc. **359** (2007), 1115–1128.
- [135] J.R. Stembridge, *Coxeter cones and their h -vectors*, Adv. Math. **217** (2008), 1935–1961.
- [136] V. Strehl, *Symmetric Eulerian distributions for involutions*, Sémin. Lothar. Combin. **1** (1980), 1 pp.
- [137] H. Sun and Y. Wang, *A group action on derangements*, Electron. J. Combin. **21** (2014), Research Paper 1.67, 5 pp. (electronic).
- [138] V. Tewari, personal communication, October 2017.
- [139] V.D. Volodin, *Cubical realizations of flag nestohedra and a proof of Gal’s conjecture for them*, Uspekhi Mat. Nauk. **65** (2010), 188–190.
- [140] M. Wachs, *Poset Topology: Tools and Applications*, in *Geometric Combinatorics* (E. Miller, V. Reiner and B. Sturmfels, eds.), IAS/Park City Mathematics Series **13**, pp. 497–615, Amer. Math. Society, Providence, R.I., 2007.
- [141] A. Zelevinsky, *Nested complexes and their polyhedral realizations*, Pure Appl. Math. Quart. **2** (2006), 655–671.
- [142] X. Zhang, *On q -derangement polynomials*, in *Combinatorics and Graph Theory ’95*, Vol. 1 (Hefei), World Sci. Publishing, River Edge, N.J., 1995, pp. 462–465.

DEPARTMENT OF MATHEMATICS, NATIONAL AND KAPODISTRIAN UNIVERSITY OF ATHENS, PANEPIS-
 TIMITOUPOLIS, 15784 ATHENS, GREECE
E-mail address: caath@math.uoa.gr