

# ENUMERATION OF STANDARD PUZZLES

GUO-NIU HAN

I.R.M.A., UMR 7501  
Université de Strasbourg and CNRS  
7 rue René-Descartes, 67084 Strasbourg, France  
Email: `guoniu.han@unistra.fr`

ABSTRACT. We introduce a large family of combinatorial objects, called standard puzzles, defined by very simple rules. We focus on the standard puzzles for which the enumeration problems can be solved by explicit formulas or by classical numbers, such as binomial coefficients, Fibonacci numbers, tangent numbers, Catalan numbers, . . .

## 1. INTRODUCTION

We introduce a large family of combinatorial objects, called *standard puzzles*, defined by very simple rules, and study their enumeration problems. The topic of the paper may be classified as belonging to *Enumerative Combinatorics*, since several of those standard puzzles can be solved by explicit formulas or by using classical numbers, such as binomial coefficients, Fibonacci numbers, tangent numbers, Catalan numbers, . . .

The general definition of a standard puzzle is inspired by the following classical topics in Enumerative Combinatorics: (1) polyominoes [11] ; (2) standard Young tableaux [14] ; (3) permutation patterns [12] ; (4) doubloons, which were introduced recently [5, 2, 3].

We do not pretend to establish a general principle that will make the enumeration of all those standard puzzles possible. We will only provide a large list of standard puzzles with their first values, and the OEIS outputs whenever the sequence is already listed in the On-Line Encyclopedia of Integer Sequences [8]. Up to order 4 there are 114 sequences identified in OEIS, meaning that those sequences have already been found in various analytical, arithmetical or combinatorial contexts and can also be derived in our puzzle model, but 1339 are still unknown, that is, do have puzzle descriptions, but have not been encountered in other contexts.

## 2. DEFINITION OF STANDARD PUZZLE

A *piece* is a square having four numbers, called *labels*, written in its corners. A *puzzle* is a connected arrangement of pieces in the  $\mathbb{Z} \times \mathbb{Z}$ -plane such that the joining corners of all the pieces have the same labels (see Fig. 1). Pieces and puzzles can be translated, but can be neither rotated, nor reflected. The *shape* of each puzzle is a polyomino obtained from the puzzle by removing the labels.

A *standard puzzle* of shape  $\lambda$  is a puzzle such that the multi-set of all its labels is simply  $\{1, 2, \dots, m\}$ . This implies that  $m$  is the number of vertices of the polyomino  $\lambda$ . In particular, the four labels of a piece occurring in a standard puzzle are all distinct. Replacing the four




FIGURE 1. Adding a piece to a puzzle

labels by  $\{1, 2, 3, 4\}$  respecting the label ordering yields a *standard piece*. This operation is called a *reduction* and will be denoted by  $\Omega$  (see Fig. 2). Two pieces are said to be *identical* if they have the same reductions. In the rest of the paper, puzzle means standard puzzle and piece means standard piece. We then have only twenty-four different pieces which are listed and coded in Table 1. Note that the letters “I” and “O” are not used.



FIGURE 2. Reduction of a piece

|  |  |  |  |  |  |
|--|--|--|--|--|--|
| $A = \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}$ | $B = \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}$ | $C = \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}$ | $D = \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}$ | $E = \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}$ | $F = \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}$ |
| $G = \begin{bmatrix} 4 & 3 \\ 2 & 1 \end{bmatrix}$ | $H = \begin{bmatrix} 3 & 4 \\ 2 & 1 \end{bmatrix}$ | $J = \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix}$ | $K = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}$ | $L = \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix}$ | $M = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}$ |
| $N = \begin{bmatrix} 4 & 2 \\ 3 & 1 \end{bmatrix}$ | $P = \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix}$ | $Q = \begin{bmatrix} 4 & 1 \\ 3 & 2 \end{bmatrix}$ | $R = \begin{bmatrix} 1 & 4 \\ 3 & 2 \end{bmatrix}$ | $S = \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix}$ | $T = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ |
| $U = \begin{bmatrix} 3 & 2 \\ 4 & 1 \end{bmatrix}$ | $V = \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix}$ | $W = \begin{bmatrix} 3 & 1 \\ 4 & 2 \end{bmatrix}$ | $X = \begin{bmatrix} 1 & 3 \\ 4 & 2 \end{bmatrix}$ | $Y = \begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix}$ | $Z = \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix}$ |

TABLE 1. The twenty-four pieces and its codes

Given a standard puzzle  $\alpha$ , a set  $\mathcal{P}$  of standard pieces is called *support* of the puzzle  $\alpha$  if the reduction of each piece occurring in the puzzle  $\alpha$  is an element of  $\mathcal{P}$ . Hence, the minimal support of  $\alpha$  is the set of all different pieces of  $\alpha$  after reduction. For example, the standard puzzle given in Fig. 3 contains seven pieces, but only four of them are different  $\begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix}$ ,  $\begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}$ ,  $\begin{bmatrix} 1 & 4 \\ 3 & 2 \end{bmatrix}$ ,  $\begin{bmatrix} 3 & 2 \\ 4 & 1 \end{bmatrix}$ . The minimal support is the set of those four pieces and each set of pieces containing those four pieces is a support of the puzzle. The main problem discussed in the paper is the following.

**Problem 1.** *Given a set of pieces  $\mathcal{P}$  and a polyomino  $\lambda$ , count the number of standard puzzles of shape  $\lambda$  whose support is equal to  $\mathcal{P}$ .*





FIGURE 3. A standard puzzle

If the support is  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix} \right\}$  and the shape is the Ferrers diagram [15] of an integer partition  $\mu = (\mu_1 \geq \mu_2 \geq \dots \geq \mu_\ell > 0)$  such that  $\mu_1 = \mu_2$  and  $\mu_\ell \geq 2$ , then a standard puzzle is just a standard Young tableau [14] of shape  $\mu$ . Hence, the number of standard puzzles is the number of standard Young tableaux of fixed shape (See Fig. 4 in Section 4). Problem 1 can be solved by using the famous hook length formula [4].

However, the problem is very hard to solve in general. In the rest of the paper we focus our attention to a very special shape, namely, the  $2 \times n$  matrix. For convenience puzzles of shape  $2 \times n$  will be represented by two-row matrices. For example  $\begin{bmatrix} 5784 \\ 1236 \end{bmatrix}$  stands for



In this case the set of all puzzles made by using the support  $\mathcal{P}$  is denoted by  $\mathcal{P}^n$ . For example, take  $\mathcal{P} = BC = \{B, C\} = \left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix} \right\}$ . Then

$$\begin{aligned} BC^2 &= \left\{ \begin{bmatrix} 34 \\ 12 \end{bmatrix}, \begin{bmatrix} 42 \\ 13 \end{bmatrix} \right\} \\ BC^3 &= \left\{ \begin{bmatrix} 563 \\ 124 \end{bmatrix}, \begin{bmatrix} 463 \\ 125 \end{bmatrix}, \begin{bmatrix} 456 \\ 123 \end{bmatrix}, \begin{bmatrix} 364 \\ 125 \end{bmatrix}, \begin{bmatrix} 356 \\ 124 \end{bmatrix} \right\} \\ BC^4 &= \left\{ \begin{bmatrix} 6784 \\ 1235 \end{bmatrix}, \begin{bmatrix} 5784 \\ 1236 \end{bmatrix}, \begin{bmatrix} 5684 \\ 1237 \end{bmatrix}, \begin{bmatrix} 5678 \\ 1234 \end{bmatrix}, \begin{bmatrix} 4785 \\ 1236 \end{bmatrix}, \begin{bmatrix} 4685 \\ 1237 \end{bmatrix}, \begin{bmatrix} 4678 \\ 1235 \end{bmatrix}, \right. \\ &\quad \left. \begin{bmatrix} 4586 \\ 1237 \end{bmatrix}, \begin{bmatrix} 4578 \\ 1236 \end{bmatrix}, \begin{bmatrix} 3785 \\ 1246 \end{bmatrix}, \begin{bmatrix} 3685 \\ 1247 \end{bmatrix}, \begin{bmatrix} 3678 \\ 1245 \end{bmatrix}, \begin{bmatrix} 3586 \\ 1247 \end{bmatrix}, \begin{bmatrix} 3578 \\ 1246 \end{bmatrix} \right\} \end{aligned}$$

The sequence  $(|BC^n|)_n = (|BC^2|, |BC^3|, |BC^4|, \dots)$ , equal to

$$(2, 5, 14, 42, 132, 429, 1430, 4862, 16796, 58786, 208012, \dots),$$

is the well-known sequence of the Catalan numbers. However, many sequences defined by other supports are not identified in OEIS. For example, with  $\mathcal{P} = CK = \left\{ \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix} \right\}$  we have

$$\begin{aligned} CK^2 &= \left\{ \begin{bmatrix} 42 \\ 13 \end{bmatrix}, \begin{bmatrix} 14 \\ 23 \end{bmatrix} \right\} \\ CK^3 &= \left\{ \begin{bmatrix} 625 \\ 134 \end{bmatrix}, \begin{bmatrix} 526 \\ 134 \end{bmatrix}, \begin{bmatrix} 426 \\ 135 \end{bmatrix}, \begin{bmatrix} 164 \\ 235 \end{bmatrix} \right\} \\ CK^4 &= \left\{ \begin{bmatrix} 8275 \\ 1346 \end{bmatrix}, \begin{bmatrix} 7285 \\ 1346 \end{bmatrix}, \begin{bmatrix} 6285 \\ 1347 \end{bmatrix}, \begin{bmatrix} 5286 \\ 1347 \end{bmatrix}, \begin{bmatrix} 4286 \\ 1357 \end{bmatrix}, \begin{bmatrix} 1847 \\ 2356 \end{bmatrix}, \begin{bmatrix} 1748 \\ 2356 \end{bmatrix}, \begin{bmatrix} 1648 \\ 2357 \end{bmatrix} \right\} \end{aligned}$$

The sequence  $(|CK^n|)_n$  is

$$(2, 4, 8, 26, 66, 276, 816, 4050, 13410, 75780, 274680 \dots)$$

which is not in OEIS.

### 3. THE TWO-LINE MATRIX SHAPE

When the shape is a  $2 \times n$  matrix, the enumeration problems for the supports  $\mathcal{P}$  and  $\mathcal{P}'$  are equivalent, denoted by  $\mathcal{P} \equiv \mathcal{P}'$ , if  $\mathcal{P}'$  is obtained from  $\mathcal{P}$  by applying the following basic transformations one or more times:

- (T1) exchanging left column and right column in every piece;
- (T2) exchanging top row and bottom row in every piece;
- (T3) replacing each label  $a$  by  $(5 - a)$  in every piece.

For example, the support  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix} \right\}$  is equivalent to  $\left\{ \begin{bmatrix} 3 & 4 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 4 & 3 \\ 2 & 1 \end{bmatrix} \right\}$  by (T1), to  $\left\{ \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \right\}$  by (T2) and to  $\left\{ \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix} \right\}$  by (T3). The enumeration problems for the following supports  $\mathcal{P}$  are all equivalent.

$$\begin{aligned} \left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix} \right\} &\equiv \left\{ \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix} \right\} \equiv \left\{ \begin{bmatrix} 4 & 3 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 2 & 1 \end{bmatrix} \right\} \equiv \left\{ \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix} \right\} \\ &\equiv \left\{ \begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix} \right\} \equiv \left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 3 \\ 2 & 1 \end{bmatrix} \right\} \equiv \left\{ \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \right\} \equiv \left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 2 & 1 \end{bmatrix} \right\}. \end{aligned}$$

A support  $\mathcal{P}$  is said to be *connected* if for every pair of pieces from  $\mathcal{P}$ , there is a puzzle containing those two pieces. The non-connected supports  $\mathcal{P}$  can be fully characterized, and have one of the following three forms:

$$\begin{aligned} (a) & \left\{ \left[ \begin{array}{cc} \dot{\wedge} & \dot{\wedge} \\ \dot{\cdot} & \dot{\cdot} \end{array} \right], \left[ \begin{array}{cc} \dot{\wedge} & \dot{\wedge} \\ \dot{\cdot} & \dot{\cdot} \end{array} \right], \dots, \left[ \begin{array}{cc} \dot{\wedge} & \dot{\wedge} \\ \dot{\cdot} & \dot{\cdot} \end{array} \right]; \left[ \begin{array}{cc} \dot{\vee} & \dot{\vee} \\ \dot{\cdot} & \dot{\cdot} \end{array} \right], \left[ \begin{array}{cc} \dot{\vee} & \dot{\vee} \\ \dot{\cdot} & \dot{\cdot} \end{array} \right], \dots, \left[ \begin{array}{cc} \dot{\vee} & \dot{\vee} \\ \dot{\cdot} & \dot{\cdot} \end{array} \right] \right\} \\ (b) & \left\{ \left[ \begin{array}{cc} \dot{\wedge} & \dot{\vee} \\ \dot{\cdot} & \dot{\cdot} \end{array} \right]; \left[ \begin{array}{cc} \dot{\wedge} & \dot{\vee} \\ \dot{\cdot} & \dot{\cdot} \end{array} \right]; \dots; \left[ \begin{array}{cc} \dot{\wedge} & \dot{\vee} \\ \dot{\cdot} & \dot{\cdot} \end{array} \right] \right\} \\ (c) & \left\{ \left[ \begin{array}{cc} \dot{\vee} & \dot{\wedge} \\ \dot{\cdot} & \dot{\cdot} \end{array} \right]; \left[ \begin{array}{cc} \dot{\vee} & \dot{\wedge} \\ \dot{\cdot} & \dot{\cdot} \end{array} \right]; \dots; \left[ \begin{array}{cc} \dot{\vee} & \dot{\wedge} \\ \dot{\cdot} & \dot{\cdot} \end{array} \right] \right\} \end{aligned}$$

For example, the support  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}; \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix} \right\}$  is not connected (type (a)). It contains two connected components. For studying the enumeration problem we need only consider the connected supports. After reduction by (T1–T3), the number of possible connected supports  $\mathcal{P}$  is shown in the following table

$$\begin{array}{lcccccccc} |\mathcal{P}| & : & 1 & 2 & 3 & 4 & 5 & 6 & \dots \\ \#\{\mathcal{P}\} & : & 6 & 37 & 259 & 1391 & 5460 & ? & \dots \end{array}$$

The list of all those sets  $\mathcal{P}$  with the sequences can be found in Section 6.

**Problem 2.** *What is the sequence of the numbers of possible connected supports  $\#\{\mathcal{P}\}$  ?*

Fix a support  $\mathcal{P}$ . Let  $f_n[\begin{smallmatrix} X \\ Y \end{smallmatrix}]$  be the number of all puzzles of shape  $2 \times n$  such that the rightmost column is  $\begin{smallmatrix} X \\ Y \end{smallmatrix}$ . Then, the number  $f_n$  of all puzzles of shape  $2 \times n$  is then

$$f_n = \sum_{X,Y} f_n[\begin{smallmatrix} X \\ Y \end{smallmatrix}], \tag{1}$$

where the sum ranges over all ordered pairs of  $[2n] \times [2n]$ .

The *inverse reduction*  $x' = \Omega^{-1}(x; X, Y)$  of  $x$  by  $\{X, Y\}$  is defined by

$$x' = \begin{cases} x; & \text{if } x \leq a - 1 \\ x + 1; & \text{if } a \leq x \leq b - 2 \\ x + 2, & \text{if } b - 1 \leq x \end{cases}$$

where  $a = \min(X, Y)$  and  $b = \max(X, Y)$ .

**Proposition 1.** *Let  $1 \leq X, Y \leq 2n$  and  $X \neq Y$ . Then*

$$f_n[\begin{smallmatrix} X \\ Y \end{smallmatrix}] = \sum_{x,y} f_{n-1}[\begin{smallmatrix} x \\ y \end{smallmatrix}] \tag{2}$$

where the sum ranges over all ordered pairs  $(x, y)$  of  $[2n-2] \times [2n-2]$  such that the reduction of  $\begin{bmatrix} \Omega^{-1}(x; X, Y) & X \\ \Omega^{-1}(y; X, Y) & Y \end{bmatrix}$  is a piece in  $\mathcal{P}$ .

Formulas (1) and (2) are used for computing the first values of the number of standard puzzles ( $f_n$ ).

#### 4. SELECTED EXAMPLES

In Section 6 we display the full list of all standard puzzle sequences including the outputs from OEIS if any. Each item in the dictionary may contain the following fields:

- Codes and pieces: The support  $\mathcal{P}$  of the standard puzzles
- See also: List of other supports  $\mathcal{P}'$  such  $|\mathcal{P}^n| = |\mathcal{P}'^n|$
- Seq: The sequence  $(|\mathcal{P}^n|)$  for  $n = 2, 3, 4, \dots$
- Var: This is a variant of the sequence of  $(|\mathcal{P}^n|)$ . Sometimes the sequence is not in OEIS but a slight modification of that sequence is in OEIS.
- OEIS: The OEIS output contains three parts: the OEIS code, the number of results found (written in square brackets [ ]), and the description of the first result found in OEIS.

In the present paper we will not reproduce all the formulas observed in the same way as formulas (3)–(10) that are presented further below. In fact, we have 114 sequences identified in OEIS up to order  $|\mathcal{P}| = 4$ . It corresponds to 309 supports  $\mathcal{P}$ , that is, we have 309 formulas to prove!

For example, we find the following item in the dictionary.

---

**BC**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix} \right\}$  See also **BD**

Seq= 2, 5, 14, 42, 132, 429, 1430, 4862, 16796, 58786, 208012

OEIS: [A000108](#) [5] Catalan numbers:  $C(n) = \text{binomial}(2n,n)/(n+1) = (2n)!/(n!(n+1)!)$ .  
Also called Segner numbers.

---

From the above item *BC* in the dictionary we extract the following results.

**Theorem 2.** *We have*

$$|BC^n| = \left| \left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix} \right\}^n \right| = C_n; \quad (3)$$

$$|BD^n| = \left| \left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix} \right\}^n \right| = C_n, \quad (4)$$

where  $C_n = \frac{1}{n+1} \binom{2n}{n}$  is the Catalan numbers [10].

*Proof.* Equation (4) is well-known since the puzzles with support  $BD = \left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix} \right\}$  are just the standard Young tableaux, which can be counted by the famous hook length formula [4]. In fact, each puzzle with support  $BD$  is just a labelling of the vertices of the following diagram such that the labels are increasing in the sense of the arrows [9, p. 227].




FIGURE 4. Puzzles having the support  $BD$

However, identity (3) cannot be directly found in the references. A further bijection is to be constructed to map a puzzle from  $BC^n$  onto a puzzle from  $BD^n$ . Each puzzle with support  $BC$  contains no piece  $C$ , or only one piece  $C$  at the rightmost position. Hence, the puzzles with support  $BC$  can be characterized by the following two diagrams:



FIGURE 5. Puzzles having the support  $BC$

In Fig. 4 and 5 the labels of certain vertices can only have fixed values, in particular maximal or minimal values. Removing those vertices yields the simplified diagrams Figures 6 and 7. The standard labellings of the diagram in Fig. 6 such that  $a < b$  (respectively  $a > b$ ) are in bijection with the standard labellings of the left diagram (respectively right diagram) in Fig. 7. This gives a bijection between  $BC^n$  and  $BD^n$ .  $\square$




FIGURE 6. Puzzles having the support  $BD$  (after simplification)



FIGURE 7. Puzzles having the support  $BC$  (after simplification)

Recall that the classical tangent numbers  $T_{2n+1}$ , secant numbers  $E_{2n}$  and the unsigned Genocchi numbers  $G_{2n}$  are defined by the following generating functions [13].

$$\begin{aligned} \tan u &= \sum_{n \geq 0} \frac{u^{2n+1}}{(2n+1)!} T_{2n+1} \\ \sec u &= \sum_{n \geq 0} \frac{u^{2n}}{(2n)!} E_{2n} \\ \frac{2u}{e^u + 1} &= u + \sum_{n \geq 1} (-1)^n \frac{u^{2n}}{(2n)!} G_{2n} \end{aligned}$$

It is well-known that  $T_{2n+1}$  and  $E_{2n}$  count the numbers of alternating permutations of length  $2n+1$  and  $2n$  respectively [13]. The unsigned Genocchi numbers  $G_{2n} = nT_{2n-1}/2^{2n-2}$  count the numbers of surjective staircases [1]. The support  $\mathcal{P} = BCEG$  involves those numbers.

---

**BCEG**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 4 & 3 \\ 2 & 1 \end{bmatrix} \right\}$  See also **BEGJ**, **BEGY**, **BEJV**, **BEVY**, **BJTV**, **EJRV**

Seq= 4, 24, 272, 4960, 132672, 4893056, 237969664, 14756156928  
 Var= 1, 3, 17, 155, 2073, 38227, 929569, 28820619, 1109652905  
 OEIS: [A110501](#) [2] Unsigned Genocchi numbers (of first kind) of even index.

---

From the above item  $BCEG$  in the dictionary we read the following seven identities as shown in Theorem 3 and Conjecture 4.

**Theorem 3.** *We have*

$$|BEGJ^n| = |\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 4 & 3 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix} \right\}^n| = nT_{2n-1}/2^{n-2}; \tag{5}$$

$$|BEGY^n| = |\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 4 & 3 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix} \right\}^n| = nT_{2n-1}/2^{n-2}; \tag{6}$$

$$|BEJV^n| = |\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \right\}^n| = nT_{2n-1}/2^{n-2}; \tag{7}$$

$$|BEVY^n| = |\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix} \right\}^n| = nT_{2n-1}/2^{n-2}; \tag{8}$$

$$|BJTV^n| = |\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \right\}^n| = nT_{2n-1}/2^{n-2}; \tag{9}$$

$$|EJRV^n| = |\left\{ \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 3 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \right\}^n| = nT_{2n-1}/2^{n-2}. \tag{10}$$

*Proof.* The enumeration of  $|BJTV^n|$  is derived in [5, 2] and identity (9) is proven to be true. Notice that no easy direct proof of (9) is known. We prove that all the six left-hand sides of equations (5)-(10) are equal to

$$|BGTY^n| = |\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 3 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix} \}^n|. \quad (11)$$

Let  $S$  be a subset of  $\{1, 2, \dots, n\}$ . The *flip map*  $\phi_S : \alpha \mapsto \beta$  is a transformation which maps a puzzle  $\alpha = \begin{bmatrix} x_1 x_2 \cdots x_n \\ y_1 y_2 \cdots y_n \end{bmatrix}$  onto  $\beta = \begin{bmatrix} a_1 a_2 \cdots a_n \\ b_1 b_2 \cdots b_n \end{bmatrix}$  such that  $a_i = x_i, b_i = y_i$  for  $i \notin S$  and  $a_i = y_i, b_i = x_i$  for  $i \in S$ . Notice that the flip map is an involution. The following diagram shows the actions of the flip maps onto some pieces:

$$\begin{array}{ccc} (B, G, T, Y) & \xrightarrow{\phi_{\{1,2\}}} & (T, Y, B, G) \\ \downarrow \phi_{\{2\}} & & \\ (E, J, R, V) & \xrightarrow{\phi_{\{1,2\}}} & (R, V, E, J) \end{array}$$

The *important facts* are that the flip map  $\phi_{\{1,2\}}$  does not change the two supports  $\{B, G, T, Y\}$  and  $\{E, J, R, V\}$ ; moreover, those two supports do not contain any common piece. We can prove that all the  $2^4 = 16$  supports  $\mathcal{P}$  derived from the Cartesian product  $\Gamma := \{B, E\} \times \{G, J\} \times \{T, R\} \times \{Y, V\}$  yield the same enumeration sequences ( $|\mathcal{P}^n|$ ). For example, we will explain why

$$|BGTY^n| = |BJRY^n|. \quad (12)$$

Notice that

$$|BGTY^n| = |BG^n| + |TY^n| = 2|BG^n|. \quad (13)$$

Let  $\alpha$  be a puzzle in  $BGTY^n$ . We construct a puzzle  $\beta$  in  $BJRY^n$  in a unique manner by applying an adequate sequence of flip maps.

Start at  $\gamma := \alpha$ ; from left to right look for the first piece in  $\gamma$  that is not in  $\{B, J, R, Y\}$ , that is, the first piece equal to  $G$  or  $T$ . Let  $(i, i+1)$  be the position of that piece. Apply the flip map  $\phi_{\{i+1, i+2, \dots, n-1\}}$  to  $\gamma$ , to obtain a new puzzle. By convention, let  $\gamma := \phi_{\{i+1, i+2, \dots, n-1\}}(\gamma)$ . Hence, the piece at position  $(i, i+1)$  becomes  $J$  or  $R$ , and all pieces on the right of that position are still in  $\{B, G, T, Y\}$ , by the important facts mentioned above. Repeat this process until no more  $G$  or  $T$  are in  $\gamma$  to get the puzzle  $\beta := \gamma$  in  $BJRY^n$ .

For example, take  $\alpha = \begin{bmatrix} 11 & 13 & 10 & 6 & 12 & 8 & 14 \\ 7 & 9 & 2 & 1 & 4 & 3 & 5 \end{bmatrix} \in BGTY^n$ . The calculation in Table 4 shows that  $\beta = \begin{bmatrix} 11 & 13 & 2 & 1 & 12 & 3 & 14 \\ 7 & 9 & 10 & 6 & 4 & 8 & 5 \end{bmatrix} \in BJRY^n$ . Note that the number of a piece, for example  $B$ , common to two different supports, is not preserved.

This process is reversible due to the important facts mentioned above. Hence, identity (12) is proved. Now, since  $BJTV, EJRV \in \Gamma$ , identities (9) and (10) are proved. In the same manner,  $BGRV, EGTY, BJRV, BJRY \in \Gamma$  and  $BGRV \equiv BEGJ, EGTY \equiv BEGY, BJRV \equiv BEJV, EJRY \equiv BEVY$ . This achieves the proof of (5)-(8).  $\square$

The next identity cannot be proved in the same manner, since the flip maps are not enough to produce the bijection. Further operations are to be constructed.

**Conjecture 4.** We have

$$|BCEG^n| = |\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 4 & 3 \\ 2 & 1 \end{bmatrix} \}^n| = nT_{2n-1}/2^{n-2}. \quad (14)$$



|   | puzzle   | first $G/T$ | position | flip                    |
|---|--|-------------|----------|-------------------------|
| → | $\begin{bmatrix} 11 & 13 & 10 & 6 & 12 & 8 & 14 \\ & B & G & G & B & G & B \\ 7 & 9 & 2 & 1 & 4 & 3 & 5 \end{bmatrix}$ | $G$         | $(2, 3)$ | $S = \{3, 4, 5, 6, 7\}$ |
| → | $\begin{bmatrix} 11 & 13 & 2 & 1 & 4 & 3 & 5 \\ & B & J & Y & T & Y & T \\ 7 & 9 & 10 & 6 & 12 & 8 & 14 \end{bmatrix}$ | $T$         | $(4, 5)$ | $S = \{5, 6, 7\}$       |
| → | $\begin{bmatrix} 11 & 13 & 2 & 1 & 12 & 8 & 14 \\ & B & J & Y & R & G & B \\ 7 & 9 & 10 & 6 & 4 & 3 & 5 \end{bmatrix}$ | $G$         | $(5, 6)$ | $S = \{6, 7\}$          |
| → | $\begin{bmatrix} 11 & 13 & 2 & 1 & 12 & 3 & 5 \\ & B & J & Y & R & J & T \\ 7 & 9 & 10 & 6 & 4 & 8 & 14 \end{bmatrix}$ | $T$         | $(6, 7)$ | $S = \{7\}$             |
| → | $\begin{bmatrix} 11 & 13 & 2 & 1 & 12 & 3 & 14 \\ & B & J & Y & R & J & R \\ 7 & 9 & 10 & 6 & 4 & 8 & 5 \end{bmatrix}$ |             |          |                         |

TABLE 2. Bijection from  $BGTY^n$  onto  $BJRY^n$

Other supports  $\mathcal{P}$  giving classical numbers are listed below without proof.

**ACX**: Fibonacci numbers;

**AB, ABCD**: double factorial numbers;

**CDW**: Koch snowflake, number of angles after  $n$  iterations;

**ACET**: number of branching configurations of RNA with  $n$  or fewer hairpins;

**ACMT**: super-Catalan numbers or little Schröder numbers;

**ADHN**: number of permutations of length  $2n - 1$  with no local maxima or minima in even positions;

**AELM**: number of permutations of length  $n$  which avoid the patterns 231, 12534;

**BDFK**: number of Dyck paths of semilength  $n$  having no DUDU's starting at level 1.

### 5. WHERE ARE THE SECANT NUMBERS?

Since the tangent numbers appear in the puzzle sequence with support  $BEGJ$  (see (5)), a natural question rises: what about the secant numbers  $(E_{2n})_{n \geq 2}$ :

$$(5, 61, 1385, 50521, 2702765, 199360981, 19391512145, \dots) ? \tag{15}$$

The initial motivation was to define a support that would generate a puzzle sequence for the secant numbers. <sup>1</sup> We developed a computer program to generate the puzzle sequences up to order  $|\mathcal{P}| = 4$ . Unfortunately, no secant number sequence could be found.

---

<sup>1</sup> A support, namely,  $\mathcal{P} = ABDGH$ , that generates the secant numbers is found recently by Jiayi Lu and Yuanzhe Ding [7].

Notice that the secant numbers appeared in the puzzle sequence for the support  $BJTV$ , but a non-trivial modification is required. By (9) we have

$$|BJTV^n| = \sum_{X,Y} |BJTV^n[X_Y]| = \frac{n}{2^{n-2}} T_{2n-1}, \quad (16)$$

where  $BJTV^n[X_Y]$  is the subset of puzzles in  $NJTV^n$  such that the rightmost column is  $[X_Y]$ . In [3] we obtained the following identity by using the signed doubloon model:

$$\sum_{X,Y} |BJTV^n[X_Y]| \times Q_n(X, Y) = 2^{-n} E_{2n}, \quad (17)$$

where

$$Q_n(X, Y) = \begin{cases} \sum_{k=X}^{Y-1} \binom{2n}{k} & \text{if } X < Y \\ 0 & \text{if } X > Y \end{cases}$$

However, the secant number sequence (15) does not directly appear in the dictionary without the coefficient  $Q_n(X, Y)$ . It is possible that order  $|\mathcal{P}| \leq 4$  is not large enough. Why do not choose a bigger support, for example  $\mathcal{P} = CEHJLPRVX$ . The puzzle sequence is

$$(9, 111, 2505, 91961, 4913789, 364074545, 35418898477, \dots) \quad (18)$$

We do not obtain the secant sequence (15). It is amusing that the quotient of (18) by the secant sequence (15) is approximately 1.8:

$$(1.800, 1.820, 1.809, 1.820, 1.818, 1.826, 1.827, \dots)$$

What does it mean?

## 6. DICTIONARY OF THE STANDARD PUZZLE SEQUENCES

### Sequence identified for $|\mathcal{P}| = 1$

**A**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix} \right\}$  See also **D**

Seq= 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1

OEIS: [A000012](#) [1450] The simplest sequence of positive numbers: the all 1's sequence.

**B**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix} \right\}$

Seq= 1, 2, 5, 14, 42, 132, 429, 1430, 4862, 16796, 58786

OEIS: [A000108](#) [5] Catalan numbers:  $C(n) = \text{binomial}(2n, n) / (n+1) = (2n)! / (n!(n+1)!)$ . Also called Segner numbers.

**C**  $\left\{ \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix} \right\}$  See also **E**, **F**

Seq= 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0

OEIS: [A121373](#) [1470] Expansion of  $f(q) = f(q, -q^2)$  in powers of  $q$  where  $f(q, r)$  is the Ramanujan two variable theta function.

Sequence identified for  $|\mathcal{P}| = 2$

**AB**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix} \right\}$

Seq= 2, 8, 48, 384, 3840, 46080, 645120, 10321920, 185794560, 3715891200

OEIS: [A000165](#) [1] Double factorial numbers:  $(2n)!! = 2^n n!$ .

**AC**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix} \right\}$  See also **AK**, **AX**, **CD**, **CX**, **DE**, **DF**, **EK**, **FK**

Seq= 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2

OEIS: [A055642](#) [327] Number of digits in decimal expansion of  $n$ .

**AD**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix} \right\}$

Seq= 2, 6, 23, 106, 567, 3434, 23137, 171174, 1376525, 11934581, 110817423

OEIS: [A125273](#) [1] Eigensequence of triangle A085478:  $a(n) = \sum_{k=0..n-1} A085478(n-1, k) * a(k)$  for  $n > 0$  with  $a(0) = 1$ .

**AE**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix} \right\}$

Seq= 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12

OEIS: [A000027](#) [618] The natural numbers. Also called the whole numbers, the counting numbers or the positive integers.

**AF**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix} \right\}$

Seq= 2, 4, 7, 11, 16, 22, 29, 37, 46, 56, 67

OEIS: [A000124](#) [6] Central polygonal numbers (the Lazy Caterer's sequence):  $n(n+1)/2 + 1$ ; or, maximal number of pieces formed when slicing a pa...

**AL**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 2, 5, 10, 17, 26, 37, 50, 65, 82, 101, 122

OEIS: [A002522](#) [2]  $n^2 + 1$ .

**AP**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix} \right\}$  See also **DQ**

Seq= 2, 7, 16, 29, 46, 67, 92, 121, 154, 191, 232

OEIS: [A130883](#) [1]  $2n^2 - n + 1$ .

**AR**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 3 & 2 \end{bmatrix} \right\}$  See also **CM**, **DJ**

Seq= 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22

OEIS: [A005843](#) [25] The even numbers:  $a(n) = 2n$ .

**BC**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix} \right\}$  See also **BD**

Seq= 2, 5, 14, 42, 132, 429, 1430, 4862, 16796, 58786, 208012

OEIS: [A000108](#) [5] Catalan numbers:  $C(n) = \text{binomial}(2n, n) / (n+1) = (2n)! / (n!(n+1)!)$ . Also called Segner numbers.

**BE**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix} \right\}$  See also **ER**

Seq= 2, 4, 10, 28, 84, 264, 858, 2860, 9724, 33592, 117572

OEIS: [A068875](#) [2] Expansion of  $(1+x^*C)^*C$ , where  $C = (1-(1-4*x)^{(1/2)})/(2*x)$  is g.f. for Catalan numbers, A000108.

**BF**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix} \right\}$

Seq= 2, 3, 7, 19, 56, 174, 561, 1859, 6292, 21658, 75582

OEIS: [A005807](#) [2] Sum of adjacent Catalan numbers.

**BG**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 3 \\ 2 & 1 \end{bmatrix} \right\}$

Seq= 2, 12, 136, 2480, 66336, 2446528, 118984832, 7378078464, 568142287360

OEIS: [A117513](#) [1] Number of ways of arranging  $2n$  tokens in a row, with 2 copies of each token from 1 through  $n$ , such that between every pair o...

**BJ**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix} \right\}$

Seq= 2, 6, 22, 84, 324, 1254, 4862, 18876, 73372, 285532, 1112412

OEIS: [A121686](#) [1] Number of branches in all binary trees with  $n$  edges. A binary tree is a rooted tree in which each vertex has at most two chi...

**BL**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 2, 4, 12, 40, 140, 504, 1848, 6864, 25740, 97240, 369512

OEIS: [A028329](#) [1] Twice central binomial coefficients.

### Sequence unknown for $|\mathcal{P}| = 2$

**AH**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 2 & 1 \end{bmatrix} \right\}$

Seq= 2, 12, 132, 2372, 62304, 2261668, 108184432, 6600715188, 500046044352

**BN**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 3 & 1 \end{bmatrix} \right\}$

Seq= 2, 9, 74, 974, 18831, 502459, 17671764, 792391014, 44129928926, 2987912108763

**BQ**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 3 & 2 \end{bmatrix} \right\}$

Seq= 2, 5, 21, 96, 440, 1989, 8855, 38896, 168948, 727090, 3105322

**CK**  $\left\{ \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix} \right\}$

Seq= 2, 4, 8, 26, 66, 276, 816, 4050, 13410, 75780, 274680

**CP**  $\left\{ \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix} \right\}$

Seq= 2, 10, 82, 1162, 23026, 657148, 23719394, 1137763610, 65032729314

**CR**  $\left\{ \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 3 & 2 \end{bmatrix} \right\}$

Seq= 2, 5, 18, 91, 563, 4299, 37686, 384543, 4357567, 55614775, 772479331

**DN**  $\left\{ \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 3 & 1 \end{bmatrix} \right\}$

Seq= 2, 14, 168, 3352, 96816, 3875904, 204185344, 13726330128, 1145508631264

**EU**  $\left\{ \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 4 & 1 \end{bmatrix} \right\}$

Seq= 2, 6, 26, 150, 1230, 10038, 125490, 1292166, 22184550, 271843110, 6022023210

**EV**  $\left\{ \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \right\}$

Seq= 2, 8, 58, 712, 12564, 310256, 10025978, 415159208, 21288518044, 1329526717840

**FU**  $\left\{ \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 4 & 1 \end{bmatrix} \right\}$

Seq= 2, 12, 106, 1680, 37434, 1171968, 48008850, 2516016384, 163509808050

**Sequence identified for  $|\mathcal{P}| = 3$**

**ABC**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix} \right\}$  See also **ABE**, **ABAFT**

Seq= 3, 12, 72, 576, 5760, 69120, 967680, 15482880, 278691840, 5573836800

OEIS: [A052676](#) [1] A simple regular expression in a labeled universe.

**ABD**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix} \right\}$

Seq= 3, 15, 105, 945, 10395, 135135, 2027025, 34459425, 654729075, 13749310575

OEIS: [A001147](#) [1] Double factorial numbers:  $(2n-1)!! = 1.3.5....(2n-1)$ .

**ABK**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix} \right\}$

Seq= 3, 10, 56, 432, 4224, 49920, 691200, 10967040, 196116480, 3901685760

Var= 3, 10, 28, 72, 176, 416, 960, 2176, 4864, 10752, 23552

OEIS: [A128135](#) [1] Row sums of A128134.

**ACE**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix} \right\}$  See also **ACZ**, **AKM**, **AKS**, **AMX**, **DEM**, **DFM**

Seq= 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13

OEIS: [A000027](#) [527] The natural numbers. Also called the whole numbers, the counting numbers or the positive integers.

**ACF**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix} \right\}$

Seq= 3, 5, 8, 12, 17, 23, 30, 38, 47, 57, 68

OEIS: [A022856](#) [4]  $a(n) = n-2 + \text{Sum of } a(i+1) \bmod(a(i)) \text{ for } i = 1 \text{ to } n-2, \text{ for } n \geq 3$ .

**ACL**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix} \right\}$  See also **AMR**, **CDM**

Seq= 3, 6, 11, 18, 27, 38, 51, 66, 83, 102, 123

OEIS: [A059100](#) [4]  $n^2+2$ .

**ACM**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$  See also **AFM**, **AFZ**

Seq= 3, 6, 14, 35, 90, 234, 611, 1598, 4182, 10947, 28658

OEIS: [A032908](#) [2] One of 4 3rd-order recurring sequences for which the first derived sequence and the Galois transformed sequence coincide.

**ACS**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix} \right\}$  See also **ATX**

Seq= 3, 7, 21, 71, 253, 925, 3433, 12871, 48621, 184757, 705433

Var= 0, 4, 18, 68, 250, 922, 3430, 12868, 48618, 184754, 705430

OEIS: [A115112](#) [1] Number of different ways to select n elements from two sets of n elements under the precondition of choosing at least one el...

**ACT**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \right\}$

Seq= 3, 6, 17, 62, 259, 1162, 5441, 26234, 129283, 648142, 3294865

Var= 2, 5, 16, 61, 258, 1161, 5440, 26233, 129282, 648141, 3294864

OEIS: [A104858](#) [1] Partial sums of the little Schroeder numbers (A001003).

**ACX**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 4 & 2 \end{bmatrix} \right\}$  See also **DEK**, **DFK**

Seq= 3, 5, 8, 13, 21, 34, 55, 89, 144, 233, 377

OEIS: [A000045](#) [15] Fibonacci numbers:  $F(n) = F(n-1) + F(n-2)$ ,  $F(0) = 0$ ,  $F(1) = 1$ ,  $F(2) = 1$ , ...

**AEF**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix} \right\}$  See also **AEQ**

Seq= 3, 6, 10, 15, 21, 28, 36, 45, 55, 66, 78

OEIS: [A000217](#) [29] Triangular numbers:  $a(n) = C(n+1,2) = n(n+1)/2 = 0+1+2+\dots+n$ .

**AEJ**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix} \right\}$  See also **AKR**, **AKV**, **ARX**, **CDJ**, **CDL**, **CEM**, **CFM**, **DEJ**, **DFJ**

Seq= 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23

OEIS: [A005408](#) [34] The odd numbers:  $a(n) = 2n+1$ .

**AEL**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix} \right\}$  See also **AFQ**

Seq= 3, 7, 13, 21, 31, 43, 57, 73, 91, 111, 133

OEIS: [A002061](#) [2] Central polygonal numbers:  $n^2 - n + 1$ .

**AEM**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 3, 5, 10, 23, 57, 146, 379, 989, 2586, 6767, 17713

Var= 0, 2, 7, 20, 54, 143, 376, 986, 2583, 6764, 17710

OEIS: [A035508](#) [2] Fibonacci(2n+2)-1.

**AES**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix} \right\}$

Seq= 3, 6, 17, 58, 212, 794, 3005, 11442, 43760, 167962, 646648

Var= 1, 4, 15, 56, 210, 792, 3003, 11440, 43758, 167960, 646646

OEIS: [A001791](#) [2] Binomial coefficients  $C(2n,n-1)$ .

**AEW**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 4 & 2 \end{bmatrix} \right\}$

Seq= 3, 7, 22, 73, 237, 746, 2287, 6867, 20286, 59157, 170713

Var= 1, 5, 20, 71, 235, 744, 2285, 6865, 20284, 59155, 170711

OEIS: [A054444](#) [1] Even indexed members of A001629(n),  $n \geq 2$ , (Fibonacci convolution).

**AEZ**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix} \right\}$

Seq= 3, 7, 19, 52, 140, 372, 981, 2577, 6757, 17702, 46358  
 Var= 0, 4, 16, 49, 137, 369, 978, 2574, 6754, 17699, 46355  
 OEIS: [A114185](#) [1]  $F(2n)-n-1$ , where  $F(n)$ =Fibonacci number.

**AFL**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 3, 8, 16, 27, 41, 58, 78, 101, 127, 156, 188  
 OEIS: [A104249](#) [1]  $(3^n \cdot 2 + n + 2)/2$ .

**AFS**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix} \right\}$

Seq= 3, 8, 31, 132, 564, 2382, 9951, 41228, 169768, 695862, 2842228  
 Var= 1, 6, 29, 130, 562, 2380, 9949, 41226, 169766, 695860, 2842226  
 OEIS: [A008549](#) [1] Number of ways of choosing at most  $n-1$  items from a set of size  $2n+1$ .

**AKP**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix} \right\}$  See also **APX, CDQ, DEQ, DFQ**

Seq= 3, 8, 17, 30, 47, 68, 93, 122, 155, 192, 233  
 OEIS: [A033816](#) [1]  $2n^2 + 3n + 3$ .

**AKT**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \right\}$  See also **BFM**

Seq= 3, 5, 10, 24, 66, 198, 627, 2057, 6919, 23715, 82501  
 OEIS: [A155587](#) [1] Expansion of  $(1+x \cdot c(x))/(1-x)$ ,  $c(x)$  the g.f. of A000108.

**AKU**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 4 & 1 \end{bmatrix} \right\}$  See also **AKX, CDE, CDF, DEF**

Seq= 3, 3, 3, 3, 3, 3, 3, 3, 3, 3, 3  
 OEIS: [A010701](#) [202] Constant sequence.

**AKW**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 4 & 2 \end{bmatrix} \right\}$  See also **APS, DFW**

Seq= 3, 9, 33, 129, 513, 2049, 8193, 32769, 131073, 524289, 2097153  
 OEIS: [A084508](#) [2] Partial sums of A084509. Positions of ones in the first differences of A084506.

**AMV**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \right\}$  See also **ARS**

Seq= 3, 8, 31, 126, 509, 2044, 8187, 32762, 131065, 524280, 2097143  
 Var= 0, 5, 28, 123, 506, 2041, 8184, 32759, 131062, 524277, 2097140  
 OEIS: [A124133](#) [1]  $a(n) = (-1/2) \cdot \sum_{i1+i2+i3=2n} ((2^n)! / (i1! i2! i3!)) \cdot B(i1)$  where B are the Bernoulli numbers.

**APR**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 3 & 2 \end{bmatrix} \right\}$  See also **CFW, DJQ**

Seq= 3, 10, 21, 36, 55, 78, 105, 136, 171, 210, 253  
 OEIS: [A014105](#) [1] Second hexagonal numbers:  $n(2n+1)$ .

**ART**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 3 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \right\}$  See also **BET**

Seq= 3, 8, 25, 84, 294, 1056, 3861, 14300, 53482, 201552, 764218  
 OEIS: [A038665](#) [1] Convolution of A007054 (super ballot numbers) with A000984 (central binomial coefficients).

**ARV**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 3 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \right\}$  See also **CEW**

Seq= 3, 7, 11, 15, 19, 23, 27, 31, 35, 39, 43

OEIS: [A004767](#) [7]  $4n+3$ .

**BCD**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix} \right\}$

Seq= 3, 9, 28, 90, 297, 1001, 3432, 11934, 41990, 149226, 534888

OEIS: [A000245](#) [3]  $3(2n)!/((n+2)!(n-1)!)$ .

**BCE**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix} \right\}$  See also **BDF**, **BEK**

Seq= 3, 7, 19, 56, 174, 561, 1859, 6292, 21658, 75582, 266798

OEIS: [A005807](#) [3] Sum of adjacent Catalan numbers.

**BDE**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix} \right\}$

Seq= 3, 8, 23, 70, 222, 726, 2431, 8294, 28730, 100776, 357238

OEIS: [A000782](#) [2]  $2^*Catalan(n)-Catalan(n-1)$ .

**BDL**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 3, 10, 35, 126, 462, 1716, 6435, 24310, 92378, 352716, 1352078

OEIS: [A001700](#) [3]  $C(2n+1, n+1)$ : number of ways to put  $n+1$  indistinguishable balls into  $n+1$  distinguishable boxes = number of  $(n+1)$ -st degree m...

**BEM**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 3, 6, 14, 37, 107, 329, 1055, 3486, 11780, 40510, 141286

OEIS: [A081293](#) [1]  $a(n) = A000108(n)+A014137(n)$ .

**BFL**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 3, 5, 14, 45, 154, 546, 1980, 7293, 27170, 102102, 386308

OEIS: [A078718](#) [1] Let  $f(i,j) = \text{Sum}(\text{binom}(2*i,k)*\text{binom}(2*j,i+j-k)*(-1)^{(i+j-k)},k=0..2*i)$  (this is essentially the same as the triangle in...

**BFT**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \right\}$

Seq= 3, 6, 15, 42, 126, 396, 1287, 4290, 14586, 50388, 176358

OEIS: [A120589](#) [1] Self-convolution of A120588, such that  $a(n) = 3^*A120588(n)$  for  $n \geq 2$ .

**BLM**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 3, 6, 16, 51, 177, 639, 2355, 8790, 33100, 125478, 478194

Var= 1, 4, 14, 49, 175, 637, 2353, 8788, 33098, 125476, 478192

OEIS: [A079309](#) [1]  $a(n) = C(1,1)+C(3,2)+C(5,3)+\dots+C(2n-1,n)$ .

**CDW**  $\left\{ \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 4 & 2 \end{bmatrix} \right\}$  See also **DEW**

Seq= 3, 6, 18, 66, 258, 1026, 4098, 16386, 65538, 262146, 1048578

OEIS: [A178789](#) [1] Koch snowflake: number of angles after  $n$  iterations.



**DJM**  $\left\{ \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 3, 8, 21, 46, 87, 148, 233, 346, 491, 672, 893

OEIS: [A179903](#) [1] (1, 3, 5, 7, 9...) convolved with (1, 0, 3, 5, 7, 9,...)

**EFK**  $\left\{ \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix} \right\}$

Seq= 3, 4, 6, 8, 12, 16, 24, 32, 48, 64, 96

OEIS: [A029744](#) [8] Numbers of the form  $2^n$  or  $3 \cdot 2^n$ .

**Sequence unknown for  $|\mathcal{P}| = 3$  (partial)**

**ABG**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 3 \\ 2 & 1 \end{bmatrix} \right\}$

Seq= 3, 23, 327, 7465, 249885, 11532671, 701867995, 54461600179, 5247921916235

**ABH**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 2 & 1 \end{bmatrix} \right\}$

Seq= 3, 24, 345, 7920, 264873, 12190108, 739050425, 57109234080, 5479466654645

**ABJ**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix} \right\}$  See also **ABL**

Seq= 3, 14, 96, 858, 9420, 122490, 1839600, 31325490, 596291220, 12546094050

**ABN**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 3 & 1 \end{bmatrix} \right\}$

Seq= 3, 19, 217, 3985, 107547, 4001027, 196224625, 12270923649, 953000374835

**ABP**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix} \right\}$

Seq= 3, 17, 116, 1048, 11712, 155520, 2388480, 41610240, 810270720, 17433722880

**ABQ**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 3 & 2 \end{bmatrix} \right\}$

Seq= 3, 14, 104, 1020, 12264, 173580, 2818080, 51535260, 1047274200, 23400192060

⋮

**Sequence identified for  $|\mathcal{P}| = 4$**

**ABCD**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix} \right\}$  See also **ABDE**, **ABDF**

Seq= 4, 20, 140, 1260, 13860, 180180, 2702700, 45945900, 872972100, 18332414100

Var= 1, 5, 35, 315, 3465, 45045, 675675, 11486475, 218243025, 4583103525

OEIS: [A051577](#) [2]  $(2^{*n+3})!/3$ , related to A001147 (odd double factorials).

**ABCE**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix} \right\}$  See also **ABCF**, **ABCM**, **ABCT**, **ABEF**, **ABEZ**, **ABRX**, **ABRZ**, **ABTX**, **ACER**, **ACRT**, **AERX**, **AERZ**, **AETX**, **BCEX**, **BCRX**, **BCTX**, **CERX**

Seq= 4, 16, 96, 768, 7680, 92160, 1290240, 20643840, 371589120, 7431782400

OEIS: [A032184](#) [2] "CIJ" (necklace, indistinct, labeled) transform of 1,3,5,7...

**ABCJ**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix} \right\}$  See also **ABCK**, **ABCL**, **ABDK**, **ABEJ**, **ABEL**, **ABFJ**, **ABFL**

Seq= 4, 18, 120, 1050, 11340, 145530, 2162160, 36486450, 689188500, 14404039650  
 Var= 2, 6, 30, 210, 1890, 20790, 270270, 4054050, 68918850, 1309458150  
 OEIS: [A097801](#) [2]  $(2^n)!/(n!2^{n-1})$ .

**ABCZ**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix} \right\}$  See also **ABEM**, **ABFM**, **ABFZ**

Seq= 4, 14, 80, 634, 6332, 75974, 1063624, 17017970, 306323444, 6126468862  
 Var= 3, 13, 79, 633, 6331, 75973, 1063623, 17017969, 306323443, 6126468861  
 OEIS: [A010844](#) [1]  $a(n) = 2^n a(n-1) + 1$  with  $a(0)=1$ .

**ABDJ**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix} \right\}$  See also **ABDL**

Seq= 4, 24, 192, 1920, 23040, 322560, 5160960, 92897280, 1857945600, 40874803200  
 OEIS: [A002866](#) [3]  $a(0) = 1$ ; for  $n > 0$ ,  $a(n) = 2^{n-1} n!$ .

**ABDP**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix} \right\}$  See also **ABDU**, **ABDV**

Seq= 4, 25, 210, 2205, 27720, 405405, 6756750, 126351225, 2618916300, 59580345825  
 Var= 1, 5, 35, 315, 3465, 45045, 675675, 11486475, 218243025, 4583103525  
 OEIS: [A051577](#) [2]  $(2^{n+3})!/3$ , related to A001147 (odd double factorials).

**ABKM**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$  See also **ABKS**

Seq= 4, 12, 60, 444, 4284, 50364, 695484, 11017404, 196811964, 3912703164  
 OEIS: [A004400](#) [1]  $1 + \sum_{k=1}^n 2^k k!$ ,  $k = 1 \dots n$ .

**ABMX**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 4 & 2 \end{bmatrix} \right\}$

Seq= 4, 13, 73, 577, 5761, 69121, 967681, 15482881, 278691841, 5573836801  
 Var= 3, 12, 72, 576, 5760, 69120, 967680, 15482880, 278691840, 5573836800  
 OEIS: [A052676](#) [1] A simple regular expression in a labeled universe.

**ACDF**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix} \right\}$  See also **ACDM**, **ACFK**, **ACKM**, **ADFM**, **ADFZ**, **ADKX**, **ADKZ**, **ADMX**, **AFKX**, **AFKZ**, **AFMX**, **CDFX**, **CDKX**, **CDMX**, **CFKX**

Seq= 4, 12, 46, 212, 1134, 6868, 46274, 342348, 2753050, 23869162, 221634846  
 Var= 2, 6, 23, 106, 567, 3434, 23137, 171174, 1376525, 11934581, 110817423  
 OEIS: [A125273](#) [1] Eigensequence of triangle A085478:  $a(n) = \sum_{k=0}^{n-1} A085478(n-1, k) a(k)$  for  $n > 0$  with  $a(0)=1$ .

**ACEF**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix} \right\}$  See also **ACEQ**

Seq= 4, 7, 11, 16, 22, 29, 37, 46, 56, 67, 79  
 OEIS: [A000124](#) [7] Central polygonal numbers (the Lazy Caterer's sequence):  $n(n+1)/2 + 1$ ; or, maximal number of pieces formed when slicing a pa...

**ACEJ**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix} \right\}$  See also **AKMX, AKRU, AKRX, AKSU, AKVX, CDEJ, CDEL, CDFJ, CDFL, CEFM, DEFJ, DEFM**

Seq= 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24  
 OEIS: [A005843](#) [41] The even numbers:  $a(n) = 2n$ .

**ACEL**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix} \right\}$  See also **ACFQ, AKMR, AMRX, CDEM, CDFM**

Seq= 4, 8, 14, 22, 32, 44, 58, 74, 92, 112, 134  
 OEIS: [A014206](#) [2]  $n^2+n+2$ .

**ACEM**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$  See also **ACEZ, AEFM**

Seq= 4, 9, 22, 56, 145, 378, 988, 2585, 6766, 17712, 46369  
 OEIS: [A055588](#) [2]  $a(n)=3a(n-1)-a(n-2)-1$ ;  $a(0)=1, a(1)=2$ .

**ACES**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix} \right\}$  See also **AMTX, BCFM**

Seq= 4, 11, 36, 127, 463, 1717, 6436, 24311, 92379, 352717, 1352079  
 OEIS: [A112849](#) [1] Number of congruence classes (epimorphisms/vertex partitionings induced by graph endomorphisms) of undirected cycles of even...

**ACET**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \right\}$

Seq= 4, 10, 32, 122, 516, 2322, 10880, 52466, 258564, 1296282, 6589728  
 OEIS: [A176006](#) [1] The number of branching configurations of RNA (see Sankoff, 1985) with  $n$  or fewer hairpins.

**ACEW**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 4 & 2 \end{bmatrix} \right\}$  See also **AEQW**

Seq= 4, 11, 34, 106, 325, 978, 2896, 8463, 24466, 70102, 199369  
 Var= 3, 10, 33, 105, 324, 977, 2895, 8462, 24465, 70101, 199368  
 OEIS: [A027989](#) [1]  $a(n) =$  self-convolution of row  $n$  of array  $T$  given by A027926.

**ACFL**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 4, 9, 17, 28, 42, 59, 79, 102, 128, 157, 189  
 Var= 0, 5, 13, 24, 38, 55, 75, 98, 124, 153, 185  
 OEIS: [A140090](#) [1]  $n(3n+7)/2$ .

**ACFM**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$  See also **AFQZ**

Seq= 4, 10, 26, 68, 178, 466, 1220, 3194, 8362, 21892, 57314  
 OEIS: [A052995](#) [3] A simple regular expression.

**ACLS**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix} \right\}$  See also **ARTX, BCEM, BDET, BDFL**

Seq= 4, 12, 40, 140, 504, 1848, 6864, 25740, 97240, 369512, 1410864  
 OEIS: [A100320](#) [2] A Catalan transform of  $(1+2x)/(1-2x)$ .

**ACMT**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \right\}$

Seq= 4, 12, 46, 198, 904, 4280, 20794, 103050, 518860, 2646724, 13648870

Var= 3, 11, 45, 197, 903, 4279, 20793, 103049, 518859, 2646723, 13648869

OEIS: [A001003](#) [2] Schroeder's second problem (generalized parentheses); also called super-Catalan numbers or little Schroeder numbers.

**ACXZ**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 4 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix} \right\}$  See also **DEFK**, **DEKM**, **DFKM**

Seq= 4, 8, 16, 32, 64, 128, 256, 512, 1024, 2048, 4096

OEIS: [A000079](#) [30] Powers of 2:  $a(n) = 2^n$ .

**ADHN**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 3 & 1 \end{bmatrix} \right\}$

Seq= 4, 42, 816, 25520, 1170240, 73992912, 6169370368, 655847011584

Var= 2, 14, 204, 5104, 195040, 10570416, 771171296, 72871890176, 8658173200896

OEIS: [A122647](#) [1] Number of permutations of length  $2n-1$  with no local maxima or minima in even positions.

**AEFJ**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix} \right\}$

Seq= 4, 8, 13, 19, 26, 34, 43, 53, 64, 76, 89

OEIS: [A034856](#) [1]  $C(n+1, 2) + n - 1$ .

**AEFL**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix} \right\}$  See also **AELQ**

Seq= 4, 10, 19, 31, 46, 64, 85, 109, 136, 166, 199

OEIS: [A005448](#) [1] Centered triangular numbers:  $3n(n-1)/2 + 1$ .

**AEFQ**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 3 & 2 \end{bmatrix} \right\}$  See also **AEJL**

Seq= 4, 9, 16, 25, 36, 49, 64, 81, 100, 121, 144

OEIS: [A000290](#) [9] The squares:  $a(n) = n^2$ .

**AEFS**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix} \right\}$  See also **AEQS**

Seq= 4, 12, 46, 188, 774, 3174, 12954, 52668, 213526, 863822, 3488874

Var= 2, 10, 44, 186, 772, 3172, 12952, 52666, 213524, 863820, 3488872

OEIS: [A068551](#) [1]  $4^n - \text{binomial}(2n, n)$ .

**AEFZ**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix} \right\}$  See also **AEQZ**

Seq= 4, 11, 31, 85, 228, 604, 1590, 4173, 10937, 28647, 75014

Var= 2, 9, 29, 83, 226, 602, 1588, 4171, 10935, 28645, 75012

OEIS: [A152891](#) [1]  $a=b=0; b(n)=b+n+a; a(n)=a+n+b$ .

**AEJS**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix} \right\}$

Seq= 4, 10, 32, 114, 422, 1586, 6008, 22882, 87518, 335922, 1293294

Var= 2, 8, 30, 112, 420, 1584, 6006, 22880, 87516, 335920, 1293292

OEIS: [A162551](#) [1]  $2 * C(2n, n-1)$ .

**AELM**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 4, 10, 28, 82, 242, 710, 2064, 5946, 16992, 48222, 136034

Var= 2, 5, 14, 41, 121, 355, 1032, 2973, 8496, 24111, 68017

OEIS: [A116845](#) [1] Number of permutations of length n which avoid the patterns 231, 12534.

**AELW**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 4 & 2 \end{bmatrix} \right\}$

Seq= 4, 12, 40, 132, 422, 1310, 3972, 11824, 34692, 100612, 289034

Var= 3, 11, 39, 131, 421, 1309, 3971, 11823, 34691, 100611, 289033

OEIS: [A166336](#) [1] Expansion of  $(1-4x+7x^2-4x^3+x^4)/(1-7x+17x^2-17x^3+7x^4-x^5)$

**AFLQ**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 3 & 2 \end{bmatrix} \right\}$  See also **AKPR**, **AKPV**, **APRX**, **CDJQ**, **CDLQ**, **CEMQ**, **CFMQ**, **DEJQ**, **DFJQ**

Seq= 4, 11, 22, 37, 56, 79, 106, 137, 172, 211, 254

OEIS: [A084849](#) [1]  $1+n+2n^2$ .

**AKMP**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix} \right\}$  See also **AMPX**

Seq= 4, 13, 56, 241, 1000, 4061, 16336, 65473, 262064, 1048477, 4194184

Var= 3, 12, 55, 240, 999, 4060, 16335, 65472, 262063, 1048476, 4194183

OEIS: [A024038](#) [1]  $4^n - n^2$ .

**AKMT**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \right\}$  See also **BDFM**

Seq= 4, 9, 23, 65, 197, 626, 2056, 6918, 23714, 82500, 290512

OEIS: [A014137](#) [1] Partial sums of Catalan numbers (A000108).

**AKMV**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \right\}$  See also **AKRS**, **AKSV**, **AMVX**, **CDEW**, **CDJW**, **CDLW**, **DEJW**

Seq= 4, 10, 34, 130, 514, 2050, 8194, 32770, 131074, 524290, 2097154

Var= 2, 5, 17, 65, 257, 1025, 4097, 16385, 65537, 262145, 1048577

OEIS: [A052539](#) [2]  $4^n + 1$ .

**AKPU**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 4 & 1 \end{bmatrix} \right\}$  See also **AKPX**, **CDEQ**, **CDFQ**, **DEFQ**

Seq= 4, 9, 18, 31, 48, 69, 94, 123, 156, 193, 234

OEIS: [A100037](#) [1] Positions of occurrences of the natural numbers as second subsequence in A100035.

**AKRV**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 3 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \right\}$  See also **ARVX**, **CDJL**, **CEJM**, **CEQW**

Seq= 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44

OEIS: [A008586](#) [6] Multiples of 4.

**AKRW**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 3 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 4 & 2 \end{bmatrix} \right\}$  See also **AMPR**, **APRS**

Seq= 4, 15, 62, 253, 1020, 4091, 16378, 65529, 262136, 1048567, 4194294

Var= 3, 14, 61, 252, 1019, 4090, 16377, 65528, 262135, 1048566, 4194293

OEIS: [A024037](#) [1]  $4^n - n$ .

**AKUX**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 4 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 4 & 2 \end{bmatrix} \right\}$  See also **CDEF**

Seq= 4, 4, 4, 4, 4, 4, 4, 4, 4, 4, 4

OEIS: [A000523](#) [213]  $\text{Log}_2(n)$  rounded down.

**AKWY**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 4 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix} \right\}$  See also **BDQW**

Seq= 4, 16, 92, 604, 4214, 30538, 227476, 1730788, 13393690, 105089230, 834086422

Var= 3, 15, 91, 603, 4213, 30537, 227475, 1730787, 13393689, 105089229, 834086421

OEIS: [A099251](#) [1] Bisection of A005043.

**APRV**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 3 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \right\}$  See also **CEFW**

Seq= 4, 13, 26, 43, 64, 89, 118, 151, 188, 229, 274

OEIS: [A091823](#) [1]  $a(n) = 2*n^2 + 3*n - 1$ .

**APSX**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 4 & 2 \end{bmatrix} \right\}$  See also **DFQW**

Seq= 4, 16, 64, 256, 1024, 4096, 16384, 65536, 262144, 1048576, 4194304

OEIS: [A000302](#) [2] Powers of 4.

**ARSV**  $\left\{ \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 3 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \right\}$

Seq= 4, 14, 60, 250, 1016, 4086, 16372, 65522, 262128, 1048558, 4194284

Var= 2, 12, 58, 248, 1014, 4084, 16370, 65520, 262126, 1048556, 4194282

OEIS: [A100103](#) [1]  $2^{(2*n)} - (2^n)$ .

**BCEG**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 4 & 3 \\ 2 & 1 \end{bmatrix} \right\}$  See also **BEGJ, BEGY, BEJV, BEVY, BJTV, EJRV**

Seq= 4, 24, 272, 4960, 132672, 4893056, 237969664, 14756156928, 1136284574720

Var= 1, 3, 17, 155, 2073, 38227, 929569, 28820619, 1109652905, 51943281731

OEIS: [A110501](#) [2] Unsigned Genocchi numbers (of first kind) of even index.

**BCEK**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix} \right\}$

Seq= 4, 14, 62, 324, 1936, 12962, 95786, 772196, 6729124, 62920648, 627487330

Var= 2, 7, 31, 162, 968, 6481, 47893, 386098, 3364562, 31460324, 313743665

OEIS: [A125275](#) [1] Eigensequence of triangle A039599:  $a(n) = \text{Sum}_{k=0..n-1} A039599(n-1, k) * a(k)$  for  $n > 0$  with  $a(0) = 1$ .

**BDEF**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix} \right\}$  See also **BDEM, BDFT, BEFK, BEKM, BFKT, DEFR, DEMR, EFKR**

Seq= 4, 10, 28, 84, 264, 858, 2860, 9724, 33592, 117572, 416024

OEIS: [A068875](#) [2] Expansion of  $(1+x*C)^*C$ , where  $C = (1-(1-4*x)^{(1/2)})/(2*x)$  is g.f. for Catalan numbers, A000108.

**BDEK**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix} \right\}$

Seq= 4, 12, 38, 126, 430, 1498, 5300, 18980, 68636, 250208, 918304

Var= 2, 6, 19, 63, 215, 749, 2650, 9490, 34318, 125104, 459152

OEIS: [A109262](#) [1] A Catalan transform of the Fibonacci numbers.

**BDFK**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix} \right\}$

Seq= 4, 11, 32, 99, 318, 1051, 3550, 12200, 42520, 149930, 533890

OEIS: [A135339](#) [1] Number of Dyck paths of semilength n having no DUDU's starting at level 1.

**BEFM**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 4, 8, 18, 46, 130, 394, 1252, 4112, 13836, 47428, 165000

Var= 2, 4, 9, 23, 65, 197, 626, 2056, 6918, 23714, 82500

OEIS: [A014137](#) [1] Partial sums of Catalan numbers (A000108).

**BEFT**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \right\}$

Seq= 4, 10, 30, 98, 336, 1188, 4290, 15730, 58344, 218348, 823004

Var= 1, 2, 5, 14, 42, 132, 429, 1430, 4862, 16796, 58786

OEIS: [A000108](#) [5] Catalan numbers:  $C(n) = \text{binomial}(2n,n)/(n+1) = (2n)!/(n!(n+1)!)$ . Also called Segner numbers.

**BERT**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 3 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \right\}$

Seq= 4, 16, 80, 448, 2688, 16896, 109824, 732160, 4978688, 34398208, 240787456

OEIS: [A025225](#) [2]  $a(n) = a(1)*a(n-1) + a(2)*a(n-2) + \dots + a(n-1)*a(1)$  for  $n \geq 2$ . Also  $a(n) = (2^n)*C(n-1)$ , where  $C = A000108$  (Catalan numbers).

**BFKM**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 4, 9, 22, 58, 163, 483, 1494, 4783, 15740, 52956, 181391

OEIS: [A059019](#) [1] Number of Dyck paths of semilength n with no peak at height 3.

**BFLM**  $\left\{ \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 4, 8, 20, 60, 200, 704, 2552, 9416, 35156, 132396, 501908

Var= 1, 2, 5, 15, 50, 176, 638, 2354, 8789, 33099, 125477

OEIS: [A024718](#) [1]  $(1/2)*(1 + \text{sum of } C(2k,k))$  for  $k = 0,1,2,\dots,n$ .

**CDEK**  $\left\{ \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix} \right\}$  See also **CDFK**

Seq= 4, 10, 32, 106, 412, 1634, 7240, 32722, 160436, 803002, 4279024

Var= 2, 5, 16, 53, 206, 817, 3620, 16361, 80218, 401501, 2139512

OEIS: [A081126](#) [1] Binomial transform of  $n!/\text{floor}(n/2)!$ .

**CDFW**  $\left\{ \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 4 & 2 \end{bmatrix} \right\}$  See also **CDQW**, **DEFW**, **DEQW**

Seq= 4, 13, 49, 193, 769, 3073, 12289, 49153, 196609, 786433, 3145729

OEIS: [A140660](#) [1]  $3*4^{n+1}$ .

**CDJM**  $\left\{ \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 4, 12, 30, 62, 112, 184, 282, 410, 572, 772, 1014

Var= 2, 6, 15, 31, 56, 92, 141, 205, 286, 386, 507

OEIS: [A056520](#) [1]  $(n+2)*(2^n-2-n+3)/6$

**CDKM**  $\left\{ \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 4, 12, 40, 152, 624, 2768, 13024, 64800, 337984, 1842368, 10444416

Var= 2, 6, 20, 76, 312, 1384, 6512, 32400, 168992, 921184, 5222208

OEIS: [A000898](#) [1]  $a(n) = 2(a(n-1) + (n-1)a(n-2))$ .

**CDLM**  $\left\{ \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$

Seq= 4, 10, 20, 34, 52, 74, 100, 130, 164, 202, 244

OEIS: [A005893](#) [1] Number of points on surface of tetrahedron:  $2n^2 + 2$  (coordination sequence for sodalite net) for  $n > 0$ .

**CMPW**  $\left\{ \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 4 & 2 \end{bmatrix} \right\}$

Seq= 4, 36, 568, 14560, 546492, 28289184, 1930982576, 168054225408, 18162775533620

Var= 2, 18, 284, 7280, 273246, 14144592, 965491288, 84027112704, 9081387766810

OEIS: [A131455](#) [1] Number of inequivalent properly oriented and labeled planar chord diagrams whose associated planar tree is a path on  $n+1$  ver...

**DEJM**  $\left\{ \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \right\}$  See also **DFJM**

Seq= 4, 10, 24, 50, 92, 154, 240, 354, 500, 682, 904

Var= 2, 5, 12, 25, 46, 77, 120, 177, 250, 341, 452

OEIS: [A116731](#) [1] Number of permutations of length  $n$  which avoid the patterns 321, 2143, 3124; or avoid the patterns 132, 2314, 4312, etc.

## 7. CONCLUDING REMARKS

Due to the computer performance limitation and the complexity of the structure of the standard puzzle, we only generate the puzzle sequences up to order  $|\mathcal{P}| = 4$ . The following two examples show that there may have interesting sequences hidden in the case of  $|\mathcal{P}| \geq 5$ .

*Example 1.* A support, namely,  $\mathcal{P} = ABDGH$ , that generates the secant numbers is found recently by Jiayi Lu and Yuanzhe Ding [7]. In fact, the sequence  $(|ABDGH^n|)_n$  is

$$(5, 61, 1385, 50521, 2702765, 199360981, 19391512145, \dots)$$

which is the sequence [A000364](#) of the secant numbers, shifted by two terms.

*Example 2.* The sequence  $(|AFHKQSUN^n|)_n$  is

$$(8, 84, 1632, 51040, 2340480, 147985824, 12338740736, \dots)$$

which is the sequence [A261683](#) shifted by one term. This sequence counts the number of *whirlpool permutations* of shape  $2 \times n$ , introduced by Donald Knuth in 2020 [6].

**Acknowledgement.** The author would like to thank the referees for carefully reading the paper and for giving helpful comments.



## REFERENCES

- [1] Dumont, Dominique, Interprétations combinatoires des nombres de Genocchi, *Duke math. J.*, **41**(2) (1974), pp. 305–318.
- [2] Foata, Dominique; Han, Guo-Niu, Doubloons and new  $q$ -tangent numbers, *Quarterly J. Math.*, **62** (2011), pp. 417–432.
- [3] Foata, Dominique; Han, Guo-Niu, Doubloons and  $q$ -secant numbers, *Münster J. Math.*, **3** (2010), pp. 89–110.
- [4] Frame, J. Sutherland; Robinson, Gilbert de Beauregard; Thrall, Robert M., The hook graphs of the symmetric groups, *Canad. J. Math.*, **6** (1954), pp. 316–324.
- [5] Graham, Ron; Zang, Nan, Enumerating split-pair arrangements, *J. Combin. Theory, Ser. A*, **115** (2008), p. 293–303.
- [6] Knuth, Donald, [Whirlpool Permutations](#), preprint, 2020.05.17, 4 pages.
- [7] Lu, Jiayi; Ding, Yuanzhe, A skeleton model to enumerate standard puzzle sequences. preprint, [arxiv:2106.09471](#) (2021), 21 pages.
- [8] Sloane, N. J. A., The On-Line Encyclopedia of Integer Sequences. Published electronically at <http://oeis.org/>.
- [9] Stanley, Richard P., *Enumerative Combinatorics*, vol. 2, Cambridge University Press, 1999.
- [10] Wikipedia, [Catalan number](#), Wikipedia Foundation, *revision July 9, 2021*.
- [11] Wikipedia, [Polyomino](#), Wikipedia Foundation, *revision July 9, 2021*.
- [12] Wikipedia, [Permutation pattern](#), Wikipedia Foundation, *revision July 9, 2021*.
- [13] Wikipedia, [Alternating permutation](#), Wikipedia Foundation, *revision July 9, 2021*.
- [14] Wikipedia, [Young tableaux](#), Wikipedia Foundation, *revision July 9, 2021*.
- [15] Wikipedia, [Ferrers diagram](#), Wikipedia Foundation, *revision July 9, 2021*.

---

2010 *Mathematics Subject Classification*: Primary 05A15, 05A19, 05A05; Secondary 05B50.

*Keywords*: standard puzzles, integer sequences, Fibonacci numbers, tangent numbers, Catalan numbers.