

Intervals in a family of Fibonacci lattices

SLC93

Jean-Luc Baril, Nathanaël Hassler



Overview

- 1 Fibonacci lattices
- 2 Characteristic elements
- 3 Intervals

Lattices and intervals

Definition

A *lattice* is a poset in which each pair of elements admits a *meet* (greatest lower bound) and a *join* (lowest upper bound).

Definition

In a poset (\mathcal{P}, \leq) , an *interval* $[P, Q]$ is a set of the form

$$\{R \in \mathcal{P} \mid P \leq R \leq Q\}.$$

If $[P, Q] = \{P, Q\}$, then this interval is called a *covering*.

Examples of lattices

Examples of lattices enumerated by the Catalan numbers :

- the Stanley lattice [*Stanley*, 1975]
- the Tamari lattice [*Friedman*, *Tamari*, 1967]
- the Kreweras lattice [*Kreweras*, 1972]
- the Phagocyte lattice [*Baril*, *Pallo*, 2006]
- the Pruning-grafting lattice [*Baril*, *Pallo*, 2008]
- the Pyramid lattice [*Baril*, *Kirgizov*, *Naima*, 2023]
- the Ascent lattice [*Baril*, *Bousquet-Mélou*, *Kirgizov*, *Naima*, 2024]

Examples of intervals in the Stanley lattice

The Stanley lattice. The Stanley lattice Stan_n is the lattice on Dyck paths of semilength n where $P \leq Q$ if P is always under Q when we draw them together.




Figure – The Hasse diagram of Stan_3 .

Examples of lattices on Dyck paths

The Stanley lattice. The Stanley lattice Stan_n is the lattice on Dyck paths of semilength n where $P \leq Q$ if P is always under Q when we draw them together.




Figure – The Hasse diagram of Stan_3 .

Examples of intervals in the Stanley lattice

The Stanley lattice. The Stanley lattice Stan_n is the lattice on Dyck paths of semilength n where $P \leq Q$ if P is always under Q when we draw them together.




Figure – The Hasse diagram of Stan_3 .

Enumeration of intervals

Intervals in Stan_n [*De Sainte-Catherine, Viennot, 1986*] :

$$\frac{6(2n)!(2n+2)!}{n!(n+1)!(n+2)!(n+3)!}.$$

Intervals in Tam_n [*Chapoton, 2006*] :

$$\frac{2(4n+1)!}{(n+1)!(3n+2)!}.$$

Linear intervals in both Stan_n and Tam_n [*Chenevrière, 2022*] :

$$\frac{1}{n+1} \binom{2n}{n} + \binom{2n-1}{n-2} + 2 \binom{2n-1}{n+2}.$$

Generalized Fibonacci numbers

Definition

The p -generalized Fibonacci sequences are defined for every $p \geq 2$ by

$$F_n^p = F_{n-1}^p + F_{n-2}^p + \cdots + F_{n-p}^p$$

with initial conditions $F_i^p = 0$ for $i < 0$, and $F_0^p = 1$.

Dyck paths enumerated by the Fibonacci numbers

Definition

For $p \geq 2$, let \mathcal{F}^p (resp. \mathcal{F}_n^p) be the set of Dyck paths (resp. of semilength n) avoiding the patterns DUU and D^{p+1} .

Definition

Let \mathcal{F}^∞ (resp. \mathcal{F}_n^∞) be the set of Dyck paths (resp. of semilength n) avoiding the pattern DUU .

Remark : For any $n \in \mathbb{N}$,

$$\mathcal{F}_n^2 \subseteq \mathcal{F}_n^3 \subseteq \dots \subseteq \mathcal{F}_n^p \subseteq \mathcal{F}_n^{p+1} \subseteq \dots \subseteq \mathcal{F}_n^\infty,$$

$$|\mathcal{F}_n^p| = F_n^p, \quad \text{and} \quad |\mathcal{F}_n^\infty| = 2^{n-1}.$$

Lattice on \mathcal{F}_n^p

Let \leq be the Stanley order.

Definition-Proposition

$\mathbb{F}_n^p = (\mathcal{F}_n^p, \leq)$ and $\mathbb{F}_n^\infty = (\mathcal{F}_n^\infty, \leq)$ are sublattices of the Stanley lattice.

Remark : The cover relation corresponds to transformations $DU \rightarrow UD$.



Lattice on \mathcal{F}_n^p




Figure – The Hasse diagram of \mathbb{F}_5^2 .

Upper covers

Let $F_p(x, y)$ be the generating function where the coefficient of $x^n y^k$ is the number of elements in \mathbb{F}_n^p that have exactly k upper covers.

Theorem

The generating function $F_p(x, y)$ is given by

$$F_p(x, y) = \frac{(1-x)(1+(y-1)x^p)}{1-2x+x^{p+1}-(y-1)(x^2-x^p+x^{p+1}-x^{p+2})}.$$




Figure – A path having 3 upper covers in \mathbb{F}_6^3 .

Coverings

Corollary

The generating function for the number of coverings in \mathbb{F}_n^p , $n \geq 0$, is

$$\partial_y F_p(x, y)|_{y=1} = \frac{(1-x)(x^2 - x^{p+1})(1-x^p)}{(1-2x+x^{p+1})^2}.$$

Corollary

For any $p \geq 2$, the number of meet-irreducible elements in \mathbb{F}_n^p , is given by

$$b_p(n) = \left\lfloor \frac{n^2(p-1)}{2p} \right\rfloor,$$

which also counts the number of edges in the (n, p) -Turán graph.

Boolean intervals

Definition

An interval is said *boolean* if it is isomorphic to the poset of subsets of $[n]$ ordered by inclusion.




Figure – The boolean lattice of size 3.

Boolean intervals

Theorem


The generating function $B_p(x, y)$ for the number of boolean intervals in \mathbb{F}_n^p , with respect to the semilength $n \geq 0$, and the interval height is given by

$$B_p(x, y) = \frac{(1-x)(1+yx^p)}{1-2x+x^{p+1}-y(x^2-x^p+x^{p+1}-x^{p+2})}.$$


Proof. Since \mathbb{F}_n^p is a distributive lattice, we have that

$$B_p(x, y) = F_p(x, 1+y).$$


Extending a Dyck path




Extending a Dyck path in \mathbb{F}_n^∞




Extending a Dyck path in \mathbb{F}_n^∞




Generating intervals in \mathbb{F}_n^∞



Generating intervals in \mathbb{F}_n^∞



Generating intervals in \mathbb{F}_n^∞



Generating intervals in \mathbb{F}_n^∞

Theorem

There is a bijection between intervals in \mathbb{F}_n^∞ and bicolored Motzkin paths of length $n - 1$ in the quarter plane.

Corollary

There are $\binom{2n-1}{n}$ intervals in \mathbb{F}_n^∞ .

Generating intervals in \mathbb{F}_n^∞




Figure – The generation of the interval $[U^2(UD)^3D^2(UD)^3, U^4(UD)^2D^2(UD)^2]$. This interval is thus associated with the bicolored Motzkin path $UF_2UDF_1UF_2$.

Intervals in \mathbb{F}_n^p

Theorem

There is a bijection between intervals in \mathbb{F}_n^p and bicolored Motzkin paths of length $n - 1$ and avoiding the $2^{p+1} - 1$ consecutive patterns of the set $\{F_2, U\}^p \cup \{F_2, D\}^p$.

Corollary

The generating function $J(x)$ for the number of intervals in \mathcal{F}_n^2 is

$$J(x) = \frac{-x^2 + 3x - 1 + \sqrt{x^4 - 2x^3 - x^2 - 2x + 1}}{2x(x^2 - 3x + 1)(x + 1)}.$$

The coefficient of x^n in the series expansion is asymptotically

$$\frac{11 + 5\sqrt{5}}{20} \sqrt{\frac{14\sqrt{5} - 30}{\pi}} \cdot n^{-1/2} \left(\frac{3 + \sqrt{5}}{2} \right)^n.$$

The end

Thank you for your attention !

Structure of the linear intervals I

Definition

An interval is said *linear* when all its elements are pairwise comparable.




Figure – The structures of linear intervals $[P, Q]$ in \mathbb{F}_n^P .

Structure of linear intervals II




Figure – The structures of linear intervals $[P, Q]$ in \mathbb{F}_n^p .

Enumeration of the linear intervals

Corollary

The generating function $L_2(x, y)$ of the number of linear intervals in \mathbb{F}_n^2 with respect to n and the interval height is given by

$$L_2(x, y) = \frac{x^4 y^4 + y^3 x^4 + 1}{1 - x - x^2} + \frac{x^2 y (x^2 - 1) (x^3 y^2 - 1)}{(xy - 1) (x^2 + x - 1)^2 (x^2 y - 1)}.$$

Theorem

Asymptotically, the number of linear intervals in \mathbb{F}_n^p is proportional to the number of coverings.

Catalan words

Definition

A length n *Catalan word* is a word $w_1 \dots w_n$ over the set of non-negative integers, with $w_1 = 0$ and $0 \leq w_i \leq w_{i-1} + 1$ for $i = 2, 3, \dots, n$.

Non-decreasing Catalan words

Proposition

There is a bijection between \mathcal{F}_n^p and the set \mathcal{C}_n^p of length n non-decreasing Catalan words avoiding $p + 1$ consecutive occurrences of the same letter.




Figure – The path $P = U^5 D(UD^3)^2 \in \mathcal{F}_7^\infty$ is associated with the Catalan word $w(P) = 0001112$.

Non-decreasing Catalan words




Figure – The lattice \mathbb{F}_5^3 on the non-decreasing Catalan words of length 5 avoiding 4 consecutive occurrences of the same letter.

Compositions of n

Proposition

There is a bijection between the elements of \mathcal{F}_n^p and the compositions of n with parts in $[1, p]$.




Figure – The path $P = U^5 D(UD^3)^2 \in \mathcal{F}_7^\infty$ is associated with the composition $\lambda(P) = (3, 3, 1)$.

Compositions of n




Figure – The lattice \mathbb{F}_5^3 on the compositions of 5 with parts in $[1, 3]$.

Powerset of $[1, n - 1]$

Proposition

There is a bijection between \mathcal{F}_n^p and the subsets of $[1, n - 1]$ having no p consecutive elements.




Figure – The path $P = U^5 D(UD^3)^2 \in \mathcal{F}_7^\infty$ is associated with the subset $A(P) = \{2, 3, 5, 6\} \subseteq \{1, \dots, 6\}$.

Powerset of $[1, n - 1]$




Figure – The lattice \mathbb{F}_5^3 on the subsets of $[1, 4]$ having no 3 consecutive elements.